

# Rupert Langer

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2150986/publications.pdf>

Version: 2024-02-01

167  
papers

7,735  
citations

36303

51  
h-index

64796

79  
g-index

181  
all docs

181  
docs citations

181  
times ranked

12398  
citing authors

#	ARTICLE	IF	CITATIONS
1	Metabolic tumor constitution is superior to tumor regression grading for evaluating response to neoadjuvant therapy of esophageal adenocarcinoma patients. <i>Journal of Pathology</i> , 2022, 256, 202-213.	4.5	11
2	Interspatial Distribution of Tumor and Immune Cells in Correlation with PD-L1 in Molecular Subtypes of Gastric Cancers. <i>Cancers</i> , 2022, 14, 1736.	3.7	4
3	Spatial Metabolomics Identifies Distinct Tumor-Specific Subtypes in Gastric Cancer Patients. <i>Clinical Cancer Research</i> , 2022, 28, 2865-2877.	7.0	27
4	Benchmarking weakly-supervised deep learning pipelines for whole slide classification in computational pathology. <i>Medical Image Analysis</i> , 2022, 79, 102474.	11.6	64
5	Neoadjuvant chemotherapy improves survival in patients with oesophageal mucinous adenocarcinoma: Post-hoc analysis of the UK MRC OE02 and OE05 trials. <i>European Journal of Cancer</i> , 2022, 170, 140-148.	2.8	1
6	Co-occurrence of malignant neoplasm and Hyperostosis Frontalis Interna in an Iron Age individual from M\u00e4nsingen-Rain (Switzerland): A multi-diagnostic study. <i>International Journal of Paleopathology</i> , 2021, 32, 1-8.	1.4	3
7	Adult form of Langerhans cell histiocytosis with pulmonary and hepatic involvement mimicking malignancy in a patient with chronic hepatitis C infection. <i>Radiology Case Reports</i> , 2021, 16, 327-333.	0.6	0
8	A prognostic score for non-small cell lung cancer resected after neoadjuvant therapy in comparison with the tumor-node-metastases classification and major pathological response. <i>Modern Pathology</i> , 2021, 34, 1333-1344.	5.5	22
9	Frequency and Significance of Pathologic Pulmonary Findings in Postmortem Examinationsâ€”A Single Center Experience before COVID-19. <i>Diagnostics</i> , 2021, 11, 894.	2.6	2
10	Dataset for the reporting of carcinoma of the esophagus in resection specimens: recommendations from the International Collaboration on Cancer Reporting. <i>Human Pathology</i> , 2021, 114, 54-65.	2.0	3
11	Increased LAMP2A levels correlate with a shorter disease-free survival of HER2 negative breast cancer patients and increased breast cancer cell viability. <i>Biochemical and Biophysical Research Communications</i> , 2021, 569, 47-53.	2.1	5
12	Virus-induced senescence is a driver and therapeutic target in COVID-19. <i>Nature</i> , 2021, 599, 283-289.	27.8	195
13	Development and validation of deep learning classifiers to detect Epstein-Barr virus and microsatellite instability status in gastric cancer: a retrospective multicentre cohort study. <i>The Lancet Digital Health</i> , 2021, 3, e654-e664.	12.3	69
14	Varying practices in tumor regression grading of gastrointestinal carcinomas after neoadjuvant therapy: results of an international survey. <i>Modern Pathology</i> , 2020, 33, 676-689.	5.5	31
15	Immunohistochemical analysis of the expression of cancer-associated fibroblast markers in esophageal cancer with and without neoadjuvant therapy. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2020, 476, 725-734.	2.8	15
16	Preservation of Epsteinâ€”Barr virus status and mismatch repair protein status along the metastatic course of gastric cancer. <i>Histopathology</i> , 2020, 76, 740-747.	2.9	13
17	Chaperone-Mediated Autophagy Markers LAMP2A and HSC70 Are Independent Adverse Prognostic Markers in Primary Resected Squamous Cell Carcinomas of the Lung. <i>Oxidative Medicine and Cellular Longevity</i> , 2020, 2020, 1-12.	4.0	16
18	Defense mechanisms to increasing back pressure for hepatic oxygen transport and venous return in porcine fecal peritonitis. <i>American Journal of Physiology - Renal Physiology</i> , 2020, 319, G289-G302.	3.4	4

#	ARTICLE	IF	CITATIONS
19	Analysis of cardiopulmonary findings in COVID-19 fatalities: High incidence of pulmonary artery thrombi and acute suppurative bronchopneumonia. <i>Cardiovascular Pathology</i> , 2020, 49, 107263.	1.6	105
20	Favourable long-term survival of patients with esophageal cancer treated with extended transhiatal esophagectomy combined with en bloc lymphadenectomy: results from a retrospective observational cohort study. <i>BMC Surgery</i> , 2020, 20, 197.	1.3	5
21	Tumor Regression in Lymph Node Metastases of Esophageal Adenocarcinomas after Neoadjuvant Therapy. <i>Gastrointestinal Disorders</i> , 2020, 2, 397-407.	0.8	0
22	Significance of tumour regression in lymph node metastases of gastric and gastroesophageal junction adenocarcinomas. <i>Journal of Pathology: Clinical Research</i> , 2020, 6, 263-272.	3.0	16
23	Myeloid Sarcoma Mimicking Endocarditis: An Autopsy Case. <i>International Journal of Surgical Pathology</i> , 2020, 28, 774-774.	0.8	2
24	Assessing Autophagy in Archived Tissue or How to Capture Autophagic Flux from a Tissue Snapshot. <i>Biology</i> , 2020, 9, 59.	2.8	12
25	Multimodal analysis of formalin-fixed and paraffin-embedded tissue by MALDI imaging and fluorescence in situ hybridization for combined genetic and metabolic analysis. <i>Laboratory Investigation</i> , 2019, 99, 1535-1546.	3.7	10
26	Integrative Clustering in Mass Spectrometry Imaging for Enhanced Patient Stratification. <i>Proteomics - Clinical Applications</i> , 2019, 13, e1800137.	1.6	8
27	Implementation of modern tools in autopsy practice—the way towards contemporary postmortal diagnostics. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2019, 474, 149-158.	2.8	7
28	Impact of age and sex on chemotherapy (CTx) efficacy, toxicity and survival in early oesophagogastric (OG) cancer: A pooled analysis of 3265 patients from four large randomised trials (OE02, OE05, MAGIC) Tj ETQq0 0.0 rgBT /Qverlock 10	0.0	10
29	The Chick Chorioallantoic Membrane (CAM) Assay as a Three-dimensional Model to Study Autophagy in Cancer Cells. <i>Bio-protocol</i> , 2019, 9, e3290.	0.4	5
30	The role of autophagy in HER2-targeted therapy. <i>Swiss Medical Weekly</i> , 2019, 149, w20138.	1.6	13
31	Cancer-Germline Antigen Expression Discriminates Clinical Outcome to CTLA-4 Blockade. <i>Cell</i> , 2018, 173, 624-633.e8.	28.9	113
32	PD-L1 and PD-1 and characterization of tumor-infiltrating lymphocytes in high grade sarcomas of soft tissue — prognostic implications and rationale for immunotherapy. <i>Oncolmmunology</i> , 2018, 7, e1389366.	4.6	72
33	Application of the 8th edition of the AJCC yTNM staging system shows improved prognostication in a single center cohort of esophageal carcinomas. <i>Surgical Oncology</i> , 2018, 27, 100-105.	1.6	12
34	Tumor regression grading of gastrointestinal cancers after neoadjuvant therapy. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2018, 472, 175-186.	2.8	78
35	Adverse prognostic value of PD-L1 expression in primary resected pulmonary squamous cell carcinomas and paired mediastinal lymph node metastases. <i>Modern Pathology</i> , 2018, 31, 101-110.	5.5	38
36	CDX2 in colorectal cancer is an independent prognostic factor and regulated by promoter methylation and histone deacetylation in tumors of the serrated pathway. <i>Clinical Epigenetics</i> , 2018, 10, 120.	4.1	41

#	ARTICLE	IF	CITATIONS
37	Her2-Targeted Therapy Induces Autophagy in Esophageal Adenocarcinoma Cells. <i>International Journal of Molecular Sciences</i> , 2018, 19, 3069.	4.1	23
38	Fatal Measles Virus Infection After Rituximab-Containing Chemotherapy in a Previously Vaccinated Patient. <i>Open Forum Infectious Diseases</i> , 2018, 5, ofy244.	0.9	3
39	Histology of Nivolumab-Induced Thyroiditis. <i>Thyroid</i> , 2018, 28, 1727-1728.	4.5	32
40	Lymphocytic esophagitis: an update on histologic diagnosis, endoscopic findings, and natural history. <i>Annals of the New York Academy of Sciences</i> , 2018, 1434, 185-191.	3.8	14
41	Risk factors for esophageal cancer: emphasis on infectious agents. <i>Annals of the New York Academy of Sciences</i> , 2018, 1434, 319-332.	3.8	25
42	Macroscopic Evaluation of the Trimmed Frozen Block Is a Helpful Tool for Intraoperative Assessment of Resection Margins of Breast Cancer Specimens. <i>International Journal of Surgical Pathology</i> , 2018, 26, 693-700.	0.8	1
43	Reply to the letter "How to standardize the evaluation of tumor regression grading of gastrointestinal cancers after neoadjuvant therapy?" by Dr. Nasierowska-Guttmejer and Dr. Szawlowski, <i>VIAR-D-18-00181. Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2018, 473, 257-258.	2.8	0
44	A specific expression profile of LC3B and p62 is associated with nonresponse to neoadjuvant chemotherapy in esophageal adenocarcinomas. <i>PLoS ONE</i> , 2018, 13, e0197610.	2.5	17
45	MicroRNA expression profiling for the prediction of resistance to neoadjuvant radiochemotherapy in squamous cell carcinoma of the esophagus. <i>Journal of Translational Medicine</i> , 2018, 16, 109.	4.4	34
46	Expression Analysis of Autophagy Related Markers LC3B, p62 and HMGB1 Indicate an Autophagy-Independent Negative Prognostic Impact of High p62 Expression in Pulmonary Squamous Cell Carcinomas. <i>Cancers</i> , 2018, 10, 281.	3.7	15
47	High intratumoural but not peritumoural inflammatory host response is associated with better prognosis in primary resected oesophageal adenocarcinomas. <i>Pathology</i> , 2017, 49, 30-37.	0.6	28
48	Glycine decarboxylase and HIF-1 $\alpha$ expression are negative prognostic factors in primary resected early-stage non-small cell lung cancer. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2017, 470, 323-330.	2.8	22
49	Expression patterns of programmed death-ligand 1 in esophageal adenocarcinomas: comparison between primary tumors and metastases. <i>Cancer Immunology, Immunotherapy</i> , 2017, 66, 777-786.	4.2	20
50	Low co-expression of epidermal growth factor receptor and its chaperone heat shock protein 90 is associated with worse prognosis in primary glioblastoma, IDH-wild-type. <i>Oncology Reports</i> , 2017, 38, 2394-2400.	2.6	5
51	Neoadjuvant cisplatin and fluorouracil versus epirubicin, cisplatin, and capecitabine followed by resection in patients with oesophageal adenocarcinoma (UK MRC OE05): an open-label, randomised phase 3 trial. <i>Lancet Oncology</i> , The, 2017, 18, 1249-1260.	10.7	187
52	Native glycan fragments detected by MALDI-FT-ICR mass spectrometry imaging impact gastric cancer biology and patient outcome. <i>Oncotarget</i> , 2017, 8, 68012-68025.	1.8	34
53	Expression analysis of LC3B and p62 indicates intact activated autophagy is associated with an unfavorable prognosis in colon cancer. <i>Oncotarget</i> , 2017, 8, 54604-54615.	1.8	45
54	Increased intraepithelial CD3+ T-lymphocytes and high PD-L1 expression on tumor cells are associated with a favorable prognosis in esophageal squamous cell carcinoma and allow prognostic immunogenic subgrouping. <i>Oncotarget</i> , 2017, 8, 46756-46768.	1.8	41

#	ARTICLE	IF	CITATIONS
55	Pleomorphic Rhabdomyosarcoma with an Impressive Response to Chemotherapy: Case Report and Review of the Literature. <i>Tumori</i> , 2016, 102, S57-S60.	1.1	4
56	Prognostic relevance of autophagy markers LC3B and p62 in esophageal adenocarcinomas. <i>Oncotarget</i> , 2016, 7, 39241-39255.	1.8	44
57	How Suitable is Matrix-Assisted Laser Desorption/Ionization-Time-of-Flight for Metabolite Imaging from Clinical Formalin-Fixed and Paraffin-Embedded Tissue Samples in Comparison to Matrix-Assisted Laser Desorption/Ionization-Fourier Transform Ion Cyclotron Resonance Mass Spectrometry?. <i>Analytical Chemistry</i> , 2016, 88, 5281-5289.	6.5	24
58	Autophagy and its current relevance to the diagnosis and clinical management of esophageal diseases. <i>Annals of the New York Academy of Sciences</i> , 2016, 1381, 113-121.	3.8	12
59	Macroscopy predicts tumor progression in gastric cancer: A retrospective patho-historical analysis based on Napoleon Bonaparte's autopsy report. <i>Digestive and Liver Disease</i> , 2016, 48, 1378-1385.	0.9	5
60	High-mass-resolution MALDI mass spectrometry imaging of metabolites from formalin-fixed paraffin-embedded tissue. <i>Nature Protocols</i> , 2016, 11, 1428-1443.	12.0	190
61	Detection of a primary tumor in the area of the renal artery with 18F-FDG PET/CT in a patient with metastatic undifferentiated sarcoma and a history of mid-aortic syndrome. <i>Medicine (United States)</i> , 2016, 95, e4622.	1.0	3
62	Depletion of FOXM1 via MET Targeting Underlies Establishment of a DNA Damage-Induced Senescence Program in Gastric Cancer. <i>Clinical Cancer Research</i> , 2016, 22, 5322-5336.	7.0	27
63	Impact of peritumoral and intratumoral budding in esophageal adenocarcinomas. <i>Human Pathology</i> , 2016, 52, 1-8.	2.0	31
64	Necrotizing herpes-simplex virus tonsillitis mimicking peritonsillar abscess. <i>Infection</i> , 2016, 44, 267-268.	4.7	0
65	<i>CD274/PD-L1</i> gene amplification and PD-L1 protein expression are common events in squamous cell carcinoma of the oral cavity. <i>Oncotarget</i> , 2016, 7, 12024-12034.	1.8	141
66	Prognostic value of the autophagy markers LC3 and p62/SQSTM1 in early-stage non-small cell lung cancer. <i>Oncotarget</i> , 2016, 7, 39544-39555.	1.8	93
67	High-resolution MALDI-TOF MS imaging for the analysis of metabolites from formalin-fixed, paraffin-embedded clinical tissue samples. <i>Journal of Pathology</i> , 2015, 237, 123-132.	4.5	123
68	Reliable LC3 and p62 autophagy marker detection in formalin fixed paraffin embedded human tissue by immunohistochemistry. <i>European Journal of Histochemistry</i> , 2015, 59, 2481.	1.5	117
69	Image analysis of immunohistochemistry is superior to visual scoring as shown for patient outcome of esophageal adenocarcinoma. <i>Histochemistry and Cell Biology</i> , 2015, 143, 1-9.	1.7	50
70	Epstein-Barr Virus in Gastro-Esophageal Adenocarcinomas - Single Center Experiences in the Context of Current Literature. <i>Frontiers in Oncology</i> , 2015, 5, 73.	2.8	36
71	Clinical Significance of NOTCH1 and NOTCH2 Expression in Gastric Carcinomas: An Immunohistochemical Study. <i>Frontiers in Oncology</i> , 2015, 5, 94.	2.8	19
72	Influence of Different Neoadjuvant Chemotherapy Regimens on Response, Prognosis, and Complication Rate in Patients with Esophagogastric Adenocarcinoma. <i>Annals of Surgical Oncology</i> , 2015, 22, 905-914.	1.5	14

#	ARTICLE	IF	CITATIONS
73	Tumour border configuration in colorectal cancer: proposal for an alternative scoring system based on the percentage of infiltrating margin. <i>Histopathology</i> , 2015, 67, 464-473.	2.9	19
74	Post-therapeutic response evaluation by a combination of endoscopy and CT scan in esophagogastric adenocarcinoma after chemotherapy: better than its reputation. <i>Gastric Cancer</i> , 2015, 18, 314-325.	5.3	14
75	Neoadjuvant chemotherapy for resectable oesophageal and junctional adenocarcinoma: Results from the UK Medical Research Council randomised OEO5 trial (ISRCTN 01852072).. <i>Journal of Clinical Oncology</i> , 2015, 33, 4002-4002.	1.6	59
76	VE1 immunohistochemistry predicts BRAF V600E mutation status and clinical outcome in colorectal cancer. <i>Oncotarget</i> , 2015, 6, 41453-41463.	1.8	22
77	Adult Pleomorphic Rhabdomyosarcoma: A Multicentre Retrospective Study. <i>Anticancer Research</i> , 2015, 35, 6213-7.	1.1	21
78	Tumor Budding in Upper Gastrointestinal Carcinomas. <i>Frontiers in Oncology</i> , 2014, 4, 216.	2.8	37
79	Heat Shock Protein 90 (HSP90) and Her2 in Adenocarcinomas of the Esophagus. <i>Cancers</i> , 2014, 6, 1382-1393.	3.7	13
80	Assessment of Tumor Regression of Esophageal Adenocarcinomas After Neoadjuvant Chemotherapy. <i>American Journal of Surgical Pathology</i> , 2014, 38, 1551-1556.	3.7	52
81	A Multifactorial Histopathologic Score for the Prediction of Prognosis of Resected Esophageal Adenocarcinomas After Neoadjuvant Chemotherapy. <i>Annals of Surgical Oncology</i> , 2014, 21, 915-921.	1.5	28
82	A retrospective comparative exploratory study on two Methylentetrahydrofolate Reductase (MTHFR) polymorphisms in esophagogastric cancer: the A1298C MTHFR polymorphism is an independent prognostic factor only in neoadjuvantly treated gastric cancer patients. <i>BMC Cancer</i> , 2014, 14, 58.	2.6	17
83	Expression analysis of heat shock protein 90 (HSP90) and Her2 in colon carcinoma. <i>International Journal of Colorectal Disease</i> , 2014, 29, 663-671.	2.2	20
84	Is Preoperative Chemotherapy Followed by Surgery the Appropriate Treatment for Signet Ring Cell Containing Adenocarcinomas of the Esophagogastric Junction and Stomach?. <i>Annals of Surgical Oncology</i> , 2014, 21, 1739-1748.	1.5	86
85	Evaluation of colon cancer histomorphology: a comparison between formalin and PAXgene tissue fixation by an international ring trial. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2014, 465, 509-519.	2.8	24
86	Interim endoscopy results during neoadjuvant therapy for gastric cancer correlate with histopathological response and prognosis. <i>Gastric Cancer</i> , 2014, 17, 478-488.	5.3	20
87	The Impact of Neural Invasion Severity in Gastrointestinal Malignancies. <i>Annals of Surgery</i> , 2014, 260, 900-908.	4.2	85
88	Reply to Letter. <i>Annals of Surgery</i> , 2014, 259, e30.	4.2	1
89	Investigation of IL-23 (p19, p40) and IL-23R identifies nuclear expression of IL-23 p19 as a favorable prognostic factor in colorectal cancer: a retrospective multicenter study of 675 patients. <i>Oncotarget</i> , 2014, 5, 4671-4682.	1.8	10
90	Epidermal growth factor receptor (EGFR) is an independent adverse prognostic factor in esophageal adenocarcinoma patients treated with cisplatin-based neoadjuvant chemotherapy. <i>Oncotarget</i> , 2014, 5, 6620-6632.	1.8	35

#	ARTICLE	IF	CITATIONS
91	Radial Extracorporeal Shock Wave Therapy (rESWT) Induces New Bone Formation in Vivo: Results of an Animal Study in Rabbits. <i>Ultrasound in Medicine and Biology</i> , 2013, 39, 126-133.	1.5	46
92	The Severity of Neural Invasion Is Associated with Shortened Survival in Colon Cancer. <i>Clinical Cancer Research</i> , 2013, 19, 50-61.	7.0	76
93	Loss of p53 in Enterocytes Generates an Inflammatory Microenvironment Enabling Invasion and Lymph Node Metastasis of Carcinogen-Induced Colorectal Tumors. <i>Cancer Cell</i> , 2013, 23, 93-106.	16.8	241
94	Epidermal growth factor receptor, phosphatidylinositol-3-kinase catalytic subunit/PTEN, and KRAS/NRAS/BRAF in primary resected esophageal adenocarcinomas: loss of PTEN is associated with worse clinical outcome. <i>Human Pathology</i> , 2013, 44, 829-836.	2.0	30
95	Clinical response to chemotherapy in oesophageal adenocarcinoma patients is linked to defects in mitochondria. <i>Journal of Pathology</i> , 2013, 230, 410-419.	4.5	71
96	Factors predicting prognosis and recurrence in patients with esophago-gastric adenocarcinoma and histopathological response with less than 10% residual tumor. <i>Langenbeck's Archives of Surgery</i> , 2013, 398, 239-249.	1.9	36
97	Tumor Regression Grading of Gastrointestinal Carcinomas after Neoadjuvant Treatment. <i>Frontiers in Oncology</i> , 2013, 3, 262.	2.8	105
98	Prognostic Implications of the Seventh Edition of the International Union Against Cancer Classification for Patients With Gastric Cancer: The Western Experience of Patients Treated in a Single-Center European Institution. <i>Journal of Clinical Oncology</i> , 2013, 31, 263-271.	1.6	102
99	Association between HSP90 and Her2 in Gastric and Gastroesophageal Carcinomas. <i>PLoS ONE</i> , 2013, 8, e69098.	2.5	25
100	Multiple osteosclerotic lesions in an Iron Age skull from Switzerland (320-250 BC) – an unusual case. <i>Swiss Medical Weekly</i> , 2013, 143, w13819.	1.6	3
101	Stromal cell-associated expression of kallikrein-related peptidase 6 (KLK6) indicates poor prognosis of ovarian cancer patients. <i>Biological Chemistry</i> , 2012, 393, 391-401.	2.5	36
102	Proposal for a Multifactorial Prognostic Score That Accurately Classifies 3 Groups of Gastric Carcinoma Patients With Different Outcomes After Neoadjuvant Chemotherapy and Surgery. <i>Annals of Surgery</i> , 2012, 256, 1002-1007.	4.2	53
103	TFAP2E and DKK4 and Chemoresistance in Colorectal Cancer. <i>New England Journal of Medicine</i> , 2012, 366, 44-53.	27.0	165
104	Molecular Analysis of HER2 Signaling in Human Breast Cancer by Functional Protein Pathway Activation Mapping. <i>Clinical Cancer Research</i> , 2012, 18, 6426-6435.	7.0	110
105	Tumor Classification of Six Common Cancer Types Based on Proteomic Profiling by MALDI Imaging. <i>Journal of Proteome Research</i> , 2012, 11, 1996-2003.	3.7	123
106	MALDI imaging mass spectrometry reveals COX7A2, TAGLN2 and S100-A10 as novel prognostic markers in Barrett's adenocarcinoma. <i>Journal of Proteomics</i> , 2012, 75, 4693-4704.	2.4	90
107	Pan-Histone Deacetylase Inhibitor Panobinostat Sensitizes Gastric Cancer Cells to Anthracyclines via Induction of CITED2. <i>Gastroenterology</i> , 2012, 143, 99-109.e10.	1.3	36
108	High-Grade Supraclavicular Soft Tissue Sarcoma as Secondary Malignancy After Successful Treatment of Acute Myeloid Leukemia: Case Report and Literature Review. <i>Journal of Oral and Maxillofacial Surgery</i> , 2012, 70, 2211-2217.	1.2	1

#	ARTICLE	IF	CITATIONS
109	Evidence of Prognostic Relevant Expression Profiles of Heat-Shock Proteins and Glucose-Regulated Proteins in Oesophageal Adenocarcinomas. <i>PLoS ONE</i> , 2012, 7, e41420.	2.5	25
110	Expression Profiling of Stem Cell-Related Genes in Neoadjuvant-Treated Gastric Cancer: A NOTCH2, GSK3B and $\beta$ -catenin Gene Signature Predicts Survival. <i>PLoS ONE</i> , 2012, 7, e44566.	2.5	35
111	Tissue-based proteomics reveals FXYD3, S100A11 and GSTM3 as novel markers for regional lymph node metastasis in colon cancer. <i>Journal of Pathology</i> , 2012, 228, 459-470.	4.5	107
112	High HSP27 and HSP70 expression levels are independent adverse prognostic factors in primary resected colon cancer. <i>Cellular Oncology (Dordrecht)</i> , 2012, 35, 197-205.	4.4	53
113	Surgical pathology in sub-Saharan Africa—volunteering in Malawi. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2012, 460, 363-370.	2.8	21
114	Adenocarcinomas of the Esophagogastric Junction Are More Likely to Respond to Preoperative Chemotherapy than Distal Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2012, 19, 2108-2118.	1.5	65
115	DNA Repair Gene and MTHFR Gene Polymorphisms as Prognostic Markers in Locally Advanced Adenocarcinoma of the Esophagus or Stomach Treated with Cisplatin and 5-Fluorouracil-Based Neoadjuvant Chemotherapy. <i>Annals of Surgical Oncology</i> , 2011, 18, 2688-2698.	1.5	49
116	Genetic aberrations in primary esophageal melanomas: molecular analysis of c-KIT, PDGFR, KRAS, NRAS and BRAF in a series of 10 cases. <i>Modern Pathology</i> , 2011, 24, 495-501.	5.5	32
117	DNA methyltransferase 1 as a predictive biomarker and potential therapeutic target for chemotherapy in gastric cancer. <i>European Journal of Cancer</i> , 2011, 47, 1817-1825.	2.8	114
118	Discovery of New Molecular Subtypes in Oesophageal Adenocarcinoma. <i>PLoS ONE</i> , 2011, 6, e23985.	2.5	24
119	Histological Assessment of PAXgene Tissue Fixation and Stabilization Reagents. <i>PLoS ONE</i> , 2011, 6, e27704.	2.5	70
120	Two cases of primary pulmonary angiosarcoma as a rare cause of lung haemorrhage. <i>Pathology</i> , 2011, 43, 386-389.	0.6	7
121	How to Classify Adenocarcinomas of the Esophagogastric Junction. <i>American Journal of Surgical Pathology</i> , 2011, 35, 1512-1522.	3.7	56
122	Protein Microarray-based Comparison of HER2, Estrogen Receptor, and Progesterone Receptor Status in Core Biopsies and Surgical Specimens From FFPE Breast Cancer Tissues. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2011, 19, 300-305.	1.2	22
123	Long-term Outcome of 2920 Patients With Cancers of the Esophagus and Esophagogastric Junction. <i>Annals of Surgery</i> , 2011, 253, 689-698.	4.2	132
124	Significance of Histopathological Tumor Regression After Neoadjuvant Chemotherapy in Gastric Adenocarcinomas. <i>Annals of Surgery</i> , 2011, 253, 934-939.	4.2	266
125	Clinical Significance of the Costimulatory Molecule B7-H1 in Barrett Carcinoma. <i>Annals of Thoracic Surgery</i> , 2011, 91, 1025-1031.	1.3	45
126	Molecular Imaging of Proliferation and Glucose Utilization: Utility for Monitoring Response and Prognosis after Neoadjuvant Therapy in Locally Advanced Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2011, 18, 3316-3323.	1.5	58



#	ARTICLE	IF	CITATIONS
127	Biomarker analysis of cetuximab plus oxaliplatin/leucovorin/5-fluorouracil in first-line metastatic gastric and oesophago-gastric junction cancer: results from a phase II trial of the Arbeitsgemeinschaft Internistische Onkologie (AIO). <i>BMC Cancer</i> , 2011, 11, 509.	2.6	58
128	Assessment of ErbB2 (Her2) in oesophageal adenocarcinomas: summary of a revised immunohistochemical evaluation system, bright field double in situ hybridisation and fluorescence in situ hybridisation. <i>Modern Pathology</i> , 2011, 24, 908-916.	5.5	44
129	Tumor-Specific Targeting of Pancreatic Cancer with Shiga Toxin B-Subunit. <i>Molecular Cancer Therapeutics</i> , 2011, 10, 1918-1928.	4.1	49
130	<sup>18</sup> F-FDG PET-CT Guided Salvage Neoadjuvant Radiochemotherapy of Adenocarcinoma of the Esophagogastric Junction: The MUNICON II Trial. <i>Journal of Nuclear Medicine</i> , 2011, 52, 1189-1196.	5.0	167
131	The Severity of Neural Invasion Is a Crucial Prognostic Factor in Rectal Cancer Independent of Neoadjuvant Radiochemotherapy. <i>Annals of Surgery</i> , 2010, 252, 797-804.	4.2	67
132	Peripheral T-cell Lymphoma With Progression to a Clonally Related, Epstein Barr Virus+, Cytotoxic Aggressive T-cell Lymphoma: Evidence for Secondary EBV Infection of an Established Malignant T-cell Clone. <i>American Journal of Surgical Pathology</i> , 2010, 34, 1382-1387.	3.7	14
133	Histone Deacetylase (HDAC) 1 and 2 Expression and Chemotherapy in Gastric Cancer. <i>Annals of Surgical Oncology</i> , 2010, 17, 3336-3343.	1.5	64
134	Successful evaluation of a new animal model using mice for esophageal adenocarcinoma. <i>Langenbeck's Archives of Surgery</i> , 2010, 395, 347-350.	1.9	16
135	The role of the pathologist in tissue banking: European Consensus Expert Group Report. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2010, 456, 449-454.	2.8	79
136	Multidisciplinary Treatment of Aggressive and Rapidly Progressing Biliary Papillomatosis. <i>Digestive Diseases and Sciences</i> , 2010, 55, 3627-3629.	2.3	2
137	High number of CD45RO+ tumor infiltrating lymphocytes is an independent prognostic factor in non-metastasized (stage I/IIA) esophageal adenocarcinoma. <i>BMC Cancer</i> , 2010, 10, 608.	2.6	51
138	High pretherapeutic thymidylate synthetase and MRP1 protein levels are associated with nonresponse to neoadjuvant chemotherapy in oesophageal adenocarcinoma patients. <i>Journal of Surgical Oncology</i> , 2010, 102, 503-508.	1.7	23
139	Expression of class I histone deacetylases (HDAC1 and HDAC2) in oesophageal adenocarcinomas: an immunohistochemical study. <i>Journal of Clinical Pathology</i> , 2010, 63, 994-998.	2.0	30
140	The $\beta$ -Adrenoceptor Agonist GW427353 (Solabegron) Decreases Excitability of Human Enteric Neurons via Release of Somatostatin. <i>Gastroenterology</i> , 2010, 138, 266-274.	1.3	21
141	Proteomic Analysis of PAXgene-Fixed Tissues. <i>Journal of Proteome Research</i> , 2010, 9, 5188-5196.	3.7	67
142	Pathology as the Cornerstone of Human Tissue Banking: European Consensus Expert Group Report. <i>Biopreservation and Biobanking</i> , 2009, 7, 157-160.	1.0	16
143	Imaging of Proliferation in Hepatocellular Carcinoma with the In Vivo Marker <sup>18</sup> F-Fluorothymidine. <i>Journal of Nuclear Medicine</i> , 2009, 50, 1441-1447.	5.0	49
144	Correlation of Matrix Metalloproteinases and Tissue Inhibitors of Matrix Metalloproteinase Expression in Ileal Carcinoids, Lymph Nodes and Liver Metastasis with Prognosis and Survival. <i>Neuroendocrinology</i> , 2009, 89, 66-78.	2.5	16

#	ARTICLE	IF	CITATIONS
145	Novel multiple, monoallelic <i>KRAS</i> mutations at codon 12 and 13. <i>International Journal of Cancer</i> , 2009, 125, 2744-2745.	5.1	4
146	Expression profiling identifies genes that predict recurrence of breast cancer after adjuvant CMF-based chemotherapy. <i>Breast Cancer Research and Treatment</i> , 2009, 118, 45-56.	2.5	42
147	Genome-wide analysis of genetic alterations in Barrett's adenocarcinoma using single nucleotide polymorphism arrays. <i>Laboratory Investigation</i> , 2009, 89, 385-397.	3.7	39
148	Prognostic significance of histopathological tumor regression after neoadjuvant chemotherapy in esophageal adenocarcinomas. <i>Modern Pathology</i> , 2009, 22, 1555-1563.	5.5	101
149	Safety and Effectiveness of Extracorporeal Shockwave Therapy: Results of a Rabbit Model of Chronic Osteomyelitis. <i>Ultrasound in Medicine and Biology</i> , 2009, 35, 595-602.	1.5	14
150	Testicular Metastasis From Adenocarcinoma of the Esophagus. <i>Annals of Thoracic Surgery</i> , 2009, 87, 957-959.	1.3	9
151	Activation of Human Enteric Neurons by Supernatants of Colonic Biopsy Specimens From Patients With Irritable Bowel Syndrome. <i>Gastroenterology</i> , 2009, 137, 1425-1434.	1.3	304
152	Sclerosing angiomatoid nodular transformation of the spleen presenting as a rapidly growing tumour in a patient with rectal cancer. <i>BMJ Case Reports</i> , 2009, 2009, bcr1120081191-bcr1120081191.	0.5	4
153	Interface membrane fibroblasts around aseptically loosened endoprostheses express MMP-13. <i>Journal of Orthopaedic Research</i> , 2008, 26, 143-152.	2.3	10
154	Combined analysis of Rac1, IQGAP1, Tiam1 and E-cadherin expression in gastric cancer. <i>Modern Pathology</i> , 2008, 21, 544-552.	5.5	71
155	Expression and clinical significance of Glucose Regulated Proteins GRP78 (BiP) and GRP94 (GP96) in human adenocarcinomas of the esophagus. <i>BMC Cancer</i> , 2008, 8, 70.	2.6	79
156	Enhanced Activation of Epidermal Growth Factor Receptor Caused by Tumor-Derived E-Cadherin Mutations. <i>Cancer Research</i> , 2008, 68, 707-714.	0.9	72
157	Protein Expression Profiling in Esophageal Adenocarcinoma Patients Indicates Association of Heat-Shock Protein 27 Expression and Chemotherapy Response. <i>Clinical Cancer Research</i> , 2008, 14, 8279-8287.	7.0	54
158	Comparison of Pretherapeutic and Posttherapeutic Expression Levels of Chemotherapy-Associated Genes in Adenocarcinomas of the Esophagus Treated by 5-Fluorouracil and Cisplatin-Based Neoadjuvant Chemotherapy. <i>American Journal of Clinical Pathology</i> , 2007, 128, 191-197.	0.7	27
159	Significance of HER2 Low-Level Copy Gain in Barrett's Cancer: Implications for Fluorescence In situ Hybridization Testing in Tissues. <i>Clinical Cancer Research</i> , 2007, 13, 5115-5123.	7.0	51
160	Prediction of Response to Neoadjuvant Chemotherapy in Carcinomas of the Upper Gastrointestinal Tract. , 2007, 176, 33-36.		9
161	C609T polymorphism of the NAD(P)H:quinone oxidoreductase I gene does not significantly affect susceptibility for esophageal adenocarcinoma. <i>International Journal of Cancer</i> , 2005, 113, 506-508.	5.1	15
162	Esophageal (Barrett's) adenocarcinoma is not associated with Epstein-Barr virus infection: An analysis of 162 cases. <i>International Journal of Cancer</i> , 2005, 117, 698-700.	5.1	11

#	ARTICLE	IF	CITATIONS
163	Association of Pretherapeutic Expression of Chemotherapy-Related Genes with Response to Neoadjuvant Chemotherapy in Barrett Carcinoma. <i>Clinical Cancer Research</i> , 2005, 11, 7462-7469.	7.0	68
164	Coexpression of Cyclooxygenases (COX-1, COX-2) and Vascular Endothelial Growth Factors (VEGF-A), Tj ETQq0 0 0 rgBT /Overlock 10 Tf	0.9	100
165	Serotonin Excites Neurons in the Human Submucous Plexus via 5-HT3 Receptors. <i>Gastroenterology</i> , 2005, 128, 1317-1326.	1.3	81
166	Array-based comparative genomic hybridization for the detection of DNA sequence copy number changes in Barrett's adenocarcinoma. <i>Journal of Pathology</i> , 2004, 203, 780-788.	4.5	56
167	Neoadjuvant Radiation in High-Grade Soft-Tissue Sarcomas. <i>American Journal of Surgical Pathology</i> , 0, Publish Ahead of Print, .	3.7	3