
Kyo Yamasu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2151399/publications.pdf Version: 2024-02-01



KVO YAMASU

#	Article	IF	CITATIONS
1	Formation of the Adult Rudiment of Sea Urchins Is Influenced by Thyroid Hormones. Developmental Biology, 1994, 161, 1-11.	2.0	80
2	gbx2 Homeobox gene is required for the maintenance of the isthmic region in the zebrafish embryonic brain. Developmental Dynamics, 2003, 228, 433-450.	1.8	52
3	Initial specification of the epibranchial placode in zebrafish embryos depends on the fibroblast growth factor signal. Developmental Dynamics, 2007, 236, 564-571.	1.8	50
4	Function of a sea urchin egg Src family kinasein initiating Ca2+ release at fertilization. Developmental Biology, 2003, 256, 367-378.	2.0	47
5	Induction of metamorphosis in the sand dollar Peronella japonica by thyroid hormones. Development Growth and Differentiation, 1998, 40, 307-312.	1.5	41
6	Conservative Segregation of Tetrameric Units of H3 and H4 Histones during Nuclesome Replication. Journal of Biochemistry, 1990, 107, 15-20.	1.7	37
7	The roles of the FGF signal in zebrafish embryos analyzed using constitutive activation and dominant-negative suppression of different FGF receptors. Mechanisms of Development, 2009, 126, 1-17.	1.7	34
8	Expression of the FGF receptor 2 gene (fgfr2) during embryogenesis in the zebrafish Danio rerio. Mechanisms of Development, 2002, 119, S173-S178.	1.7	33
9	Genomic organization, alternative splicing, and multiple regulatory regions of the zebrafish fgf8 gene. Development Growth and Differentiation, 2006, 48, 447-462.	1.5	27
10	FGF receptor gene expression and its regulation by FGF signaling during early zebrafish development. Genesis, 2010, 48, 707-716.	1.6	27
11	Autoregulatory loop and retinoic acid repression regulate <i>pou2/pou5f1</i> gene expression in the zebrafish embryonic brain. Developmental Dynamics, 2008, 237, 1373-1388.	1.8	26
12	Optical interrogation of neuronal circuitry in zebrafish using genetically encoded voltage indicators. Scientific Reports, 2018, 8, 6048.	3.3	24
13	Identification of ephrin-A3 and novel genes specific to the midbrain-MHB in embryonic zebrafish by ordered differential display. Mechanisms of Development, 2001, 107, 83-96.	1.7	21
14	The Protein Tyrosine Kinases of the Sea Urchin Anthocidaris crassispina. Zoological Science, 1997, 14, 941-946.	0.7	20
15	Transcription of fgf8 is regulated by activating and repressive cis-elements at the midbrain–hindbrain boundary in zebrafish embryos. Developmental Biology, 2008, 316, 471-486.	2.0	19
16	Gbx2 functions as a transcriptional repressor to regulate the specification and morphogenesis of the mid–hindbrain junction in a dosage- and stage-dependent manner. Mechanisms of Development, 2013, 130, 532-552.	1.7	19
17	Expression of a src-type protein tyrosine kinase gene, AcSrc1, in the sea urchin embryo. Development Growth and Differentiation, 1999, 41, 19-28.	1.5	18
18	Pou2, a class V POU-type transcription factor in zebrafish, regulates dorsoventral patterning and convergent extension movement at different blastula stages. Mechanisms of Development, 2012, 129, 219-235.	1.7	18

Kyo Yamasu

#	Article	IF	CITATIONS
19	Three enhancer regions regulate gbx2 gene expression in the isthmic region during zebrafish development. Mechanisms of Development, 2006, 123, 907-924.	1.7	17
20	Functional organization of DNA elements regulating SM30alpha, a spicule matrix gene of sea urchin embryos. Development Growth and Differentiation, 1999, 41, 81-91.	1.5	16
21	Molecular Cloning of a cDNA that Encodes the Precursor to Several Exogastrula-inducing Peptides, Epidermal-growth-factor-related Polypeptides of the Sea Urchin Anthocidaris crassispina. FEBS Journal, 1995, 228, 515-523.	0.2	16
22	Mesendoderm specification depends on the function of <scp>P</scp> ou2, the class V <scp>POU</scp> â€ŧype transcription factor, during zebrafish embryogenesis. Development Growth and Differentiation, 2012, 54, 686-701.	1.5	15
23	Comprehensive analysis of target genes in zebrafish embryos reveals gbx2 involvement in neurogenesis. Developmental Biology, 2017, 430, 237-248.	2.0	15
24	Fractionation of newly replicated nucleosomes by density labeling and rate zonal centrifugation for the analysis of the deposition sites of newly synthesized nucleosomal core histones. FEBS Journal, 1985, 150, 575-580.	0.2	13
25	Association of the sea urchin EGF-related peptide, EGIP-D, with fasciclin I-related ECM proteins from the sea urchin Anthocidaris crassispina. Development Growth and Differentiation, 1999, 41, 483-494.	1.5	12
26	Structure of the zebrafish fasciclin I-related extracellular matrix protein (βig-h3) and its characteristic expression during embryogenesis. Gene Expression Patterns, 2003, 3, 331-336.	0.8	11
27	Optical measurement of neuronal activity in the developing cerebellum of zebrafish using voltage-sensitive dye imaging. NeuroReport, 2018, 29, 1349-1354.	1.2	11
28	Maternal Exogastrula-Inducing Peptides (EGIPs) and Their Changes during Development in the Sea Urchin Anthocidaris crassispina. Development Growth and Differentiation, 1992, 34, 661-668.	1.5	10
29	Localization of an Exogastrula-Inducing Peptide (EGIP) in Embryos of the Sea Urchin Anthocidaris crassispina. (Exogastrula-inducing peptide (EGIP)/gastrulation/acidic vesicle/sea) Tj ETQq1 1 0.784314 rgBT /Ove	erlo r.Is 10 7	f 5ø 337 Td (≀
30	Binding Properties of Thyroxine to Nuclear Extract from Sea Urchin Larvae. Zoological Science, 2012, 29, 79-82.	0.7	9
31	Early development of the enteric nervous system visualized by using a new transgenic zebrafish line harboring a regulatory region for choline acetyltransferase a (chata) gene. Gene Expression Patterns, 2018, 28, 12-21.	0.8	9
32	Deadenylation by the <scp>CCR</scp> 4â€ <scp>NOT</scp> complex contributes to the turnover of <i>hairy</i> â€related <scp>mRNA</scp> s in the zebrafish segmentation clock. FEBS Letters, 2018, 592, 3388-3398.	2.8	9
33	Transcriptional autoregulation of zebrafish <i>tbx6</i> is required for somite segmentation. Development (Cambridge), 2019, 146, .	2.5	9
34	Functional roles of the Ripply-mediated suppression of segmentation gene expression at the anterior presomitic mesoderm in zebrafish. Mechanisms of Development, 2018, 152, 21-31.	1.7	8
35	A Protein That Binds an Exogastrula-Inducing Peptide, EGIP-D, in the Hyaline Layer of Sea Urchin Embryos. (exogastrula-inducing peptide (EGIP)/binding protein/hyaline layer/sea) Tj ETQq1 1 0.784314 rgBT /Ove	erlacta 10 ⁻	Tf 570 97 Td (u
36	The role of gastrulation brain homeobox 2 (gbx2) in the development of the ventral telencephalon in zebrafish embryos. Differentiation, 2018, 99, 28-40.	1.9	7

Kyo Yamasu

#	Article	IF	CITATIONS
37	Reassembly of Nucleosomal Histone Octamers during Replication of Chromatin1. Journal of Biochemistry, 1987, 101, 1041-1049.	1.7	6
38	Retinoic acid-dependent establishment of positional information in the hindbrain was conserved during vertebrate evolution. Developmental Biology, 2011, 350, 154-168.	2.0	6
39	Posterior–anterior gradient of zebrafish hes6 expression in the presomitic mesoderm is established by the combinatorial functions of the downstream enhancer and 3′UTR. Developmental Biology, 2016, 409, 543-554.	2.0	6
40	Involvement of an Oct4-related PouV gene, pou5f3/pou2, in neurogenesis in the early neural plate of zebrafish embryos. Developmental Biology, 2020, 457, 30-42.	2.0	6
41	Enhancer activity-based identification of functional enhancers using zebrafish embryos. Genomics, 2016, 108, 102-107.	2.9	5
42	In vitro analysis of the transcriptional regulatory mechanism of zebrafish pou5f3. Experimental Cell Research, 2018, 364, 28-41.	2.6	5
43	Molecular Cloning of a cDNA that Encodes the Precursor to Several Exogastrula-inducing Peptides, Epidermal-growth-factor-related Polypeptides of the Sea Urchin Anthocidaris crassispina. FEBS Journal, 1995, 228, 515-523.	0.2	4
44	Purification of EGIP-D-Binding Protein from the Embryos of the Sea Urchin Anthocidaris crassispina. Zoological Science, 1997, 14, 931-934.	0.7	4
45	Cloning and characterization of cDNA for syndecan core protein in sea urchin embryos. Development Growth and Differentiation, 2000, 42, 449-458.	1.5	4
46	Genomic organization of the gene that encodes the precursor to EGF-related peptides, exogastrula-inducing peptides, of the sea urchin Anthocidaris crassispina. Biochimica Et Biophysica Acta Gene Regulatory Mechanisms, 2002, 1574, 311-320.	2.4	4
47	Role of somite patterning in the formation of Weberian apparatus and pleural rib in zebrafish. Journal of Anatomy, 2020, 236, 622-629.	1.5	4
48	Role of syndecan in the elongation of postoral arms in sea urchin larvae. Development Growth and Differentiation, 2002, 44, 45-53.	1.5	3
49	Expression of the Gene for Translation Elongation Factor 1α-Related Protein during Development of the Sea Urchin Anthocidaris crassispina. Zoological Science, 1999, 16, 785-792.	0.7	2
50	Characterization of the upstream region that regulates the transcription of the gene for the precursor to EGF-related peptides, exogastrula-inducing peptides, of the sea urchin Anthocidaris crassispina. Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology, 2003, 136, 15-26.	1.6	2
51	Involvement of Oct4â€ŧype transcription factor Pou5f3 in posterior spinal cord formation in zebrafish embryos. Development Growth and Differentiation, 2021, 63, 306-322.	1.5	2
52	Conservation of the Dimeric Unit of H2A and H2B Histones during the Replication Cycle. Experimental Cell Research, 1993, 207, 226-229.	2.6	1
53	4D imaging identifies dynamic migration and the fate of gbx2-expressing cells in the brain primordium of zebrafish. Neuroscience Letters, 2019, 690, 112-119.	2.1	1
54	A globin-family protein, Cytoglobin 1, is involved in the development of neural crest-derived tissues and organs in zebrafish. Developmental Biology, 2021, 472, 1-17.	2.0	1

#	Article	IF	CITATIONS
55	FGF receptor gene expression and its regulation by FGF signaling during early zebrafish development. Genesis, 2010, 48, spcone-spcone.	1.6	0