

# Sarah Palmer Short

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2206229/publications.pdf>

Version: 2024-02-01

23  
papers

993  
citations

516710

16  
h-index

610901

24  
g-index

25  
all docs

25  
docs citations

25  
times ranked

1774  
citing authors

#	ARTICLE	IF	CITATIONS
1	MTG16 regulates colonic epithelial differentiation, colitis, and tumorigenesis by repressing E protein transcription factors. <i>JCI Insight</i> , 2022, 7, .	5.0	9
2	Colonic Epithelial-Derived Selenoprotein P Is the Source for Antioxidant-Mediated Protection in Colitis-Associated Cancer. <i>Gastroenterology</i> , 2021, 160, 1694-1708.e3.	1.3	33
3	Serine Threonine Kinase 17A Maintains the Epithelial State in Colorectal Cancer Cells. <i>Molecular Cancer Research</i> , 2019, 17, 882-894.	3.4	10
4	Blood vessel epicardial substance reduces LRP6 receptor and cytoplasmic $\beta$ -catenin levels to modulate Wnt signaling and intestinal homeostasis. <i>Carcinogenesis</i> , 2019, 40, 1086-1098.	2.8	11
5	Kaiso is required for MTG16-dependent effects on colitis-associated carcinoma. <i>Oncogene</i> , 2019, 38, 5091-5106.	5.9	10
6	Colorectal Cancer and Metabolism. <i>Current Colorectal Cancer Reports</i> , 2018, 14, 226-241.	0.5	88
7	Roles for selenium and selenoprotein P in the development, progression, and prevention of intestinal disease. <i>Free Radical Biology and Medicine</i> , 2018, 127, 26-35.	2.9	69
8	BVES is required for maintenance of colonic epithelial integrity in experimental colitis by modifying intestinal permeability. <i>Mucosal Immunology</i> , 2018, 11, 1363-1374.	6.0	18
9	BVES regulates c-Myc stability via PP2A and suppresses colitis-induced tumourigenesis. <i>Gut</i> , 2017, 66, 852-862.	12.1	43
10	Using 3D Organoid Cultures to Model Intestinal Physiology and Colorectal Cancer. <i>Current Colorectal Cancer Reports</i> , 2017, 13, 183-191.	0.5	10
11	Selenoproteins and oxidative stress-induced inflammatory tumorigenesis in the gut. <i>Cellular and Molecular Life Sciences</i> , 2017, 74, 607-616.	5.4	57
12	Selenoproteins in Tumorigenesis and Cancer Progression. <i>Advances in Cancer Research</i> , 2017, 136, 49-83.	5.0	76
13	p120-Catenin is an obligate haploinsufficient tumor suppressor in intestinal neoplasia. <i>Journal of Clinical Investigation</i> , 2017, 127, 4462-4476.	8.2	19
14	BVES Regulates Intestinal Stem Cell Programs and Intestinal Crypt Viability after Radiation. <i>Stem Cells</i> , 2016, 34, 1626-1636.	3.2	23
15	Selenoprotein P in colitis-associated carcinoma. <i>Molecular and Cellular Oncology</i> , 2016, 3, e1075094.	0.7	8
16	Selenoprotein P influences colitis-induced tumorigenesis by mediating stemness and oxidative damage. <i>Journal of Clinical Investigation</i> , 2015, 125, 2646-2660.	8.2	87
17	The transcriptional corepressor MTGR1 regulates intestinal secretory lineage allocation. <i>FASEB Journal</i> , 2015, 29, 786-795.	0.5	13
18	Phase II trial of bortezomib plus doxorubicin in hepatocellular carcinoma (E6202): a trial of the Eastern Cooperative Oncology Group. <i>Investigational New Drugs</i> , 2014, 32, 1017-1027.	2.6	27

#	ARTICLE	IF	CITATIONS
19	RAF265 Inhibits the Growth of Advanced Human Melanoma Tumors. <i>Clinical Cancer Research</i> , 2012, 18, 2184-2198.	7.0	61
20	Kaiso Directs the Transcriptional Corepressor MTG16 to the Kaiso Binding Site in Target Promoters. <i>PLoS ONE</i> , 2012, 7, e51205.	2.5	22
21	Adenoma Formation following Limited Ablation of p120-Catenin in the Mouse Intestine. <i>PLoS ONE</i> , 2011, 6, e19880.	2.5	39
22	Cytokine Receptor CXCR4 Mediates Estrogen-Independent Tumorigenesis, Metastasis, and Resistance to Endocrine Therapy in Human Breast Cancer. <i>Cancer Research</i> , 2011, 71, 603-613.	0.9	140
23	p120-catenin is essential for maintenance of barrier function and intestinal homeostasis in mice. <i>Journal of Clinical Investigation</i> , 2010, 120, 1824-1835.	8.2	119