

Dorte Herzke

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2212687/publications.pdf>

Version: 2024-02-01

129
papers

9,395
citations

50276

46
h-index

40979

93
g-index

136
all docs

136
docs citations

136
times ranked

7339
citing authors

#	ARTICLE	IF	CITATIONS
1	Information Requirements under the Essential-Use Concept: PFAS Case Studies. <i>Environmental Science & Technology</i> , 2022, 56, 6232-6242.	10.0	32
2	Correspondence regarding the Perspective "Addressing the importance of microplastic particles as vectors for long-range transport of chemical contaminants: perspective in relation to prioritizing research and regulatory actions". <i>Microplastics and Nanoplastics</i> , 2022, 2, .	8.8	1
3	A Bad Start in Life? Maternal Transfer of Legacy and Emerging Poly- and Perfluoroalkyl Substances to Eggs in an Arctic Seabird. <i>Environmental Science & Technology</i> , 2022, 56, 6091-6102.	10.0	33
4	Bioaccumulation of Per and Polyfluoroalkyl Substances in Antarctic Breeding South Polar Skuas (<i>Catharacta maccormicki</i>) and Their Prey. <i>Frontiers in Marine Science</i> , 2022, 9, .	2.5	4
5	Temporal Trends of Organochlorine and Perfluorinated Contaminants in a Terrestrial Raptor in Northern Europe Over 34 years (1986-2019). <i>Environmental Toxicology and Chemistry</i> , 2022, 41, 1508-1519.	4.3	9
6	First documentation of plastic ingestion in the arctic glaucous gull (<i>Larus hyperboreus</i>). <i>Science of the Total Environment</i> , 2022, 834, 155340.	8.0	7
7	Ecosystem specific accumulation of organohalogenated compounds: A comparison between adjacent freshwater and terrestrial avian predators. <i>Environmental Research</i> , 2022, 212, 113455.	7.5	3
8	A schematic sampling protocol for contaminant monitoring in raptors. <i>Ambio</i> , 2021, 50, 95-100.	5.5	28
9	Trophic and fitness correlates of mercury and organochlorine compound residues in egg-laying Antarctic petrels. <i>Environmental Research</i> , 2021, 193, 110518.	7.5	14
10	Ingested plastics in northern fulmars (<i>Fulmarus glacialis</i>): A pathway for polybrominated diphenyl ether (PBDE) exposure?. <i>Science of the Total Environment</i> , 2021, 778, 146313.	8.0	28
11	Oceanic long-range transport of organic additives present in plastic products: an overview. <i>Environmental Sciences Europe</i> , 2021, 33, .	5.5	43
12	Addressing Urgent Questions for PFAS in the 21st Century. <i>Environmental Science & Technology</i> , 2021, 55, 12755-12765.	10.0	17
13	Moving forward in microplastic research: A Norwegian perspective. <i>Environment International</i> , 2021, 157, 106794.	10.0	29
14	Finding essentiality feasible: common questions and misinterpretations concerning the "essential-use" concept. <i>Environmental Sciences: Processes and Impacts</i> , 2021, 23, 1079-1087.	3.5	16
15	Maternal-Child Exposures to Persistent Organic Pollutants in Dhaka, Bangladesh. <i>Exposure and Health</i> , 2020, 12, 79-87.	4.9	7
16	Pelagic vs Coastal "Key Drivers of Pollutant Levels in Barents Sea Polar Bears with Contrasted Space-Use Strategies. <i>Environmental Science & Technology</i> , 2020, 54, 985-995.	10.0	18
17	Are Fluoropolymers Really of Low Concern for Human and Environmental Health and Separate from Other PFAS?. <i>Environmental Science & Technology</i> , 2020, 54, 12820-12828.	10.0	149
18	A novel use of the leukocyte coping capacity assay to assess the immunomodulatory effects of organohalogenated contaminants in avian wildlife. <i>Environment International</i> , 2020, 142, 105861.	10.0	9

#	ARTICLE	IF	CITATIONS
19	The high persistence of PFAS is sufficient for their management as a chemical class. <i>Environmental Sciences: Processes and Impacts</i> , 2020, 22, 2307-2312.	3.5	125
20	Exposure to PFAS is Associated with Telomere Length Dynamics and Demographic Responses of an Arctic Top Predator. <i>Environmental Science & Technology</i> , 2020, 54, 10217-10226.	10.0	30
21	An overview of the uses of per- and polyfluoroalkyl substances (PFAS). <i>Environmental Sciences: Processes and Impacts</i> , 2020, 22, 2345-2373.	3.5	632
22	Car Tire Crumb Rubber: Does Leaching Produce a Toxic Chemical Cocktail in Coastal Marine Systems?. <i>Frontiers in Environmental Science</i> , 2020, 8, .	3.3	76
23	The NORMAN Association and the European Partnership for Chemicals Risk Assessment (PARC): let's cooperate!. <i>Environmental Sciences Europe</i> , 2020, 32, .	5.5	46
24	Strategies for grouping per- and polyfluoroalkyl substances (PFAS) to protect human and environmental health. <i>Environmental Sciences: Processes and Impacts</i> , 2020, 22, 1444-1460.	3.5	126
25	Spatiotemporal Analysis of Perfluoroalkyl Substances in White-Tailed Eagle (<i>Haliaeetus</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tt 50 <i>Technology</i> , 2020, 54, 5011-5020.	10.0	17
26	Contaminants, prolactin and parental care in an Arctic seabird: Contrasted associations of perfluoroalkyl substances and organochlorine compounds with egg-turning behavior. <i>General and Comparative Endocrinology</i> , 2020, 291, 113420.	1.8	14
27	White-tailed eagle (<i>Haliaeetus albicilla</i>) feathers from Norway are suitable for monitoring of legacy, but not emerging contaminants. <i>Science of the Total Environment</i> , 2019, 647, 525-533.	8.0	40
28	Levels and trends of poly- and perfluoroalkyl substances in the Arctic environment – An update. <i>Emerging Contaminants</i> , 2019, 5, 240-271.	4.9	117
29	White-Tailed Eagle (<i>Haliaeetus albicilla</i>) Body Feathers Document Spatiotemporal Trends of Perfluoroalkyl Substances in the Northern Environment. <i>Environmental Science & Technology</i> , 2019, 53, 12744-12753.	10.0	45
30	Seabird-Transported Contaminants Are Reflected in the Arctic Tundra, But Not in Its Soil-Dwelling Springtails (Collembola). <i>Environmental Science & Technology</i> , 2019, 53, 12835-12845.	10.0	11
31	Integrated exposure assessment of northern goshawk (<i>Accipiter gentilis</i>) nestlings to legacy and emerging organic pollutants using non-destructive samples. <i>Environmental Research</i> , 2019, 178, 108678.	7.5	25
32	Individual variability in contaminants and physiological status in a resident Arctic seabird species. <i>Environmental Pollution</i> , 2019, 249, 191-199.	7.5	20
33	The concept of essential use for determining when uses of PFASs can be phased out. <i>Environmental Sciences: Processes and Impacts</i> , 2019, 21, 1803-1815.	3.5	125
34	Bird feathers as a biomonitor for environmental pollutants: Prospects and pitfalls. <i>TrAC - Trends in Analytical Chemistry</i> , 2019, 118, 223-226.	11.4	78
35	Snow buntings (<i>Plectrophenax nivealis</i>) as bio-indicators for exposure differences to legacy and emerging persistent organic pollutants from the Arctic terrestrial environment on Svalbard. <i>Science of the Total Environment</i> , 2019, 667, 638-647.	8.0	26
36	Potential Effect of Migration Strategy on Pollutant Occurrence in Eggs of Arctic Breeding Barnacle Geese (<i>Branta leucopsis</i>). <i>Environmental Science & Technology</i> , 2019, 53, 5427-5435.	10.0	21

#	ARTICLE	IF	CITATIONS
37	Plastic litter in the European Arctic: What do we know?. <i>Emerging Contaminants</i> , 2019, 5, 308-318.	4.9	79
38	Plasma concentrations of organohalogenated contaminants in white-tailed eagle nestlings – The role of age and diet. <i>Environmental Pollution</i> , 2019, 246, 527-534.	7.5	30
39	Higher plasma oxidative damage and lower plasma antioxidant defences in an Arctic seabird exposed to longer perfluoroalkyl acids. <i>Environmental Research</i> , 2019, 168, 278-285.	7.5	52
40	What is the effect of phasing out long-chain per- and polyfluoroalkyl substances on the concentrations of perfluoroalkyl acids and their precursors in the environment? A systematic review. <i>Environmental Evidence</i> , 2018, 7, .	2.7	132
41	Persistent organic pollutants and organophosphate esters in feathers and blood plasma of adult kittiwakes (<i>Rissa tridactyla</i>) from Svalbard – associations with body condition and thyroid hormones. <i>Environmental Research</i> , 2018, 164, 158-164.	7.5	18
42	DNA damage in Arctic seabirds: Baseline, sensitivity to a genotoxic stressor, and association with organohalogen contaminants. <i>Environmental Toxicology and Chemistry</i> , 2018, 37, 1084-1091.	4.3	13
43	Zürich Statement on Future Actions on Per- and Polyfluoroalkyl Substances (PFASs). <i>Environmental Health Perspectives</i> , 2018, 126, 84502.	6.0	91
44	Developmental Toxicity of Perfluorooctanesulfonate (PFOS) and Its Chlorinated Polyfluoroalkyl Ether Sulfonate Alternative F-53B in the Domestic Chicken. <i>Environmental Science & Technology</i> , 2018, 52, 12859-12867.	10.0	60
45	Organochlorines, perfluoroalkyl substances, mercury, and egg incubation temperature in an Arctic seabird: Insights from data loggers. <i>Environmental Toxicology and Chemistry</i> , 2018, 37, 2881-2894.	4.3	11
46	Blood clinical-chemical parameters and feeding history in growing Japanese quail (<i>Coturnix</i>). <i>Toxicological and Environmental Chemistry</i> , 2017, 99, 938-952.	1.2	3
47	Contaminants and energy expenditure in an Arctic seabird: Organochlorine pesticides and perfluoroalkyl substances are associated with metabolic rate in a contrasted manner. <i>Environmental Research</i> , 2017, 157, 118-126.	7.5	45
48	Geographical Differences in Dietary Exposure to Perfluoroalkyl Acids between Manufacturing and Application Regions in China. <i>Environmental Science & Technology</i> , 2017, 51, 5747-5755.	10.0	39
49	Dissimilar effects of organohalogenated compounds on thyroid hormones in glaucous gulls. <i>Environmental Research</i> , 2017, 158, 350-357.	7.5	29
50	Estimating human exposure to perfluoroalkyl acids via solid food and drinks: Implementation and comparison of different dietary assessment methods. <i>Environmental Research</i> , 2017, 158, 269-276.	7.5	25
51	Anti-parasite treatment and blood biochemistry in raptor nestlings. <i>Canadian Journal of Zoology</i> , 2017, 95, 685-693.	1.0	0
52	Per- and polyfluoroalkyl substances in plasma and feathers of nestling birds of prey from northern Norway. <i>Environmental Research</i> , 2017, 158, 277-285.	7.5	26
53	Perfluorinated substances and telomeres in an Arctic seabird: Cross-sectional and longitudinal approaches. <i>Environmental Pollution</i> , 2017, 230, 360-367.	7.5	56
54	Contamination of ivory gulls (<i>Pagophila eburnea</i>) at four colonies in Svalbard in relation to their trophic behaviour. <i>Polar Biology</i> , 2017, 40, 917-929.	1.2	13

#	ARTICLE	IF	CITATIONS
55	Temporal variation in circulating concentrations of organochlorine pollutants in a pelagic seabird breeding in the high Arctic. <i>Environmental Toxicology and Chemistry</i> , 2017, 36, 442-448.	4.3	16
56	Persistent organic pollutant levels and the importance of source proximity in Baltic and Svalbard breeding common eiders. <i>Environmental Toxicology and Chemistry</i> , 2016, 35, 1526-1533.	4.3	13
57	Antioxidant Responses in Relation to Persistent Organic Pollutants and Metals in a Low- and a High-Exposure Population of Seabirds. <i>Environmental Science & Technology</i> , 2016, 50, 4817-4825.	10.0	14
58	DNA double-strand breaks in incubating female common eiders (<i>Somateria mollissima</i>): Comparison between a low and a high polluted area. <i>Environmental Research</i> , 2016, 151, 297-303.	7.5	12
59	Levels, Isomer Profiles, and Estimated Riverine Mass Discharges of Perfluoroalkyl Acids and Fluorinated Alternatives at the Mouths of Chinese Rivers. <i>Environmental Science & Technology</i> , 2016, 50, 11584-11592.	10.0	186
60	Exposure to per- and polyfluoroalkyl substances through the consumption of fish from lakes affected by aqueous film-forming foam emissions – A combined epidemiological and exposure modeling approach. <i>The SAMINOR 2 Clinical Study. Environment International</i> , 2016, 94, 272-282.	10.0	34
61	Spatial and temporal trends in perfluoroalkyl substances (PFASs) in ringed seals (<i>Pusa hispida</i>) from Svalbard. <i>Environmental Pollution</i> , 2016, 214, 230-238.	7.5	37
62	Exposure to oxychlorane is associated with shorter telomeres in arctic breeding kittiwakes. <i>Science of the Total Environment</i> , 2016, 563-564, 125-130.	8.0	47
63	Tracking pan-continental trends in environmental contamination – Using sentinel raptors – what types of samples should we use?. <i>Ecotoxicology</i> , 2016, 25, 777-801.	2.4	149
64	Negligible Impact of Ingested Microplastics on Tissue Concentrations of Persistent Organic Pollutants in Northern Fulmars off Coastal Norway. <i>Environmental Science & Technology</i> , 2016, 50, 1924-1933.	10.0	215
65	Organohalogenated contaminants in white-tailed eagle (<i>Haliaeetus albicilla</i>) nestlings: An assessment of relationships to immunoglobulin levels, telomeres and oxidative stress. <i>Science of the Total Environment</i> , 2016, 539, 337-349.	8.0	55
66	What is the effect of phasing out long-chain per- and polyfluoroalkyl substances on the concentrations of perfluoroalkyl acids and their precursors in the environment? A systematic review protocol. <i>Environmental Evidence</i> , 2015, 4, .	2.7	40
67	Perfluoroalkyl substance concentrations in a terrestrial raptor: Relationships to environmental conditions and individual traits. <i>Environmental Toxicology and Chemistry</i> , 2015, 34, 184-191.	4.3	21
68	Survival rate and breeding outputs in a high Arctic seabird exposed to legacy persistent organic pollutants and mercury. <i>Environmental Pollution</i> , 2015, 200, 1-9.	7.5	75
69	Occurrence of perfluorinated alkylated substances in cereals, salt, sweets and fruit items collected in four European countries. <i>Chemosphere</i> , 2015, 129, 179-185.	8.2	38
70	Increased adrenal responsiveness and delayed hatching date in relation to polychlorinated biphenyl exposure in Arctic-breeding black-legged kittiwakes (<i>Rissa tridactyla</i>). <i>General and Comparative Endocrinology</i> , 2015, 219, 165-172.	1.8	24
71	Are imported consumer products an important diffuse source of PFASs to the Norwegian environment?. <i>Environmental Pollution</i> , 2015, 198, 223-230.	7.5	51
72	Perfluoroalkyl substances detected in the world's southernmost marine mammal, the Weddell seal (<i>Lophoceros pinnatus</i>)	7.5	26

#	ARTICLE	IF	CITATIONS
73	Enantiomer-Selective and Quantitative Trace Analysis of Selected Persistent Organic Pollutants (POP) in Traditional Food from Western Greenland. <i>Journal of Toxicology and Environmental Health - Part A: Current Issues</i> , 2014, 77, 616-627.	2.3	14
74	Endocrine and Fitness Correlates of Long-Chain Perfluorinated Carboxylates Exposure in Arctic Breeding Black-Legged Kittiwakes. <i>Environmental Science & Technology</i> , 2014, 48, 13504-13510.	10.0	64
75	The stress of being contaminated? Adrenocortical function and reproduction in relation to persistent organic pollutants in female black legged kittiwakes. <i>Science of the Total Environment</i> , 2014, 476-477, 553-560.	8.0	36
76	Polychlorinated biphenyls (PCBs), polybrominated diphenyl ethers (PBDEs) and perfluorinated alkylated substances (PFASs) in traditional seafood items from western Greenland. <i>Environmental Science and Pollution Research</i> , 2014, 21, 4741-4750.	5.3	29
77	Levels and trends of PBDEs and HBCDs in the global environment: Status at the end of 2012. <i>Environment International</i> , 2014, 65, 147-158.	10.0	346
78	Spatial and temporal distribution of chiral pesticides in <i>Calanus</i> spp. from three Arctic fjords. <i>Environmental Pollution</i> , 2014, 192, 154-161.	7.5	18
79	Integument colouration in relation to persistent organic pollutants and body condition in arctic breeding black-legged kittiwakes (<i>Rissa tridactyla</i>). <i>Science of the Total Environment</i> , 2014, 470-471, 248-254.	8.0	18
80	Effect of Body Condition on Tissue Distribution of Perfluoroalkyl Substances (PFASs) in Arctic Fox (<i>Vulpes lagopus</i>). <i>Environmental Science & Technology</i> , 2014, 48, 11654-11661.	10.0	43
81	DNA double-strand breaks in relation to persistent organic pollutants in a fasting seabird. <i>Ecotoxicology and Environmental Safety</i> , 2014, 106, 68-75.	6.0	17
82	Perfluorinated alkylated substances in vegetables collected in four European countries; occurrence and human exposure estimations. <i>Environmental Science and Pollution Research</i> , 2013, 20, 7930-7939.	5.3	76
83	Occurrence of perfluoroalkyl substances (PFASs) in various food items of animal origin collected in four European countries. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2013, 30, 1918-1932.	2.3	71
84	Perfluoroalkyl substances in soft tissues and tail feathers of Belgian barn owls (<i>Tyto alba</i>) using statistical methods for left-censored data to handle non-detects. <i>Environment International</i> , 2013, 52, 9-16.	10.0	45
85	Ecological and spatial factors drive intra- and interspecific variation in exposure of subarctic predatory bird nestlings to persistent organic pollutants. <i>Environment International</i> , 2013, 57-58, 25-33.	10.0	28
86	Antiparasite treatments reduce humoral immunity and impact oxidative status in raptor nestlings. <i>Ecology and Evolution</i> , 2013, 3, 5157-5166.	1.9	20
87	Dietary exposure to selected perfluoroalkyl acids (PFAAs) in four European regions. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2013, 30, 2141-2151.	2.3	59
88	Plasma concentrations of organohalogenated pollutants in predatory bird nestlings: Associations to growth rate and dietary tracers. <i>Environmental Toxicology and Chemistry</i> , 2013, 32, 2520-2527.	4.3	33
89	The structure of the fire fighting foam surfactant Forafac®1157 and its biological and photolytic transformation products. <i>Chemosphere</i> , 2012, 89, 869-875.	8.2	117
90	Temporal Dynamics of Circulating Persistent Organic Pollutants in a Fasting Seabird under Different Environmental Conditions. <i>Environmental Science & Technology</i> , 2012, 46, 10287-10294.	10.0	36

#	ARTICLE	IF	CITATIONS
91	Blood plasma clinicalâ€“chemical parameters as biomarker endpoints for organohalogen contaminant exposure in Norwegian raptor nestlings. <i>Ecotoxicology and Environmental Safety</i> , 2012, 80, 76-83.	6.0	48
92	Perfluoroalkyl and polyfluoroalkyl substances (PFASs) in consumer products in Norway â€“ A pilot study. <i>Chemosphere</i> , 2012, 88, 980-987.	8.2	215
93	Relationships between POPs and baseline corticosterone levels in black-legged kittiwakes (<i>Rissa</i>) Tj ETQq1 1 0.784314 rgBT /Overlock	7.5	33
94	Temporal trends and spatial differences of perfluoroalkylated substances in livers of harbor porpoise (<i>Phocoena phocoena</i>) populations from Northern Europe, 1991â€“2008. <i>Science of the Total Environment</i> , 2012, 419, 216-224.	8.0	30
95	Temporal Trends and Pattern of Polyfluoroalkyl Compounds in Tawny Owl (<i>Strix aluco</i>) Eggs from Norway, 1986â€“2009. <i>Environmental Science & Technology</i> , 2011, 45, 8090-8097.	10.0	69
96	Impacts of Climate and Feeding Conditions on the Annual Accumulation (1986â€“2009) of Persistent Organic Pollutants in a Terrestrial Raptor. <i>Environmental Science & Technology</i> , 2011, 45, 7542-7547.	10.0	21
97	The search for alternative aqueous film forming foams (AFFF) with a low environmental impact: Physiological and transcriptomic effects of two ForafacÂ® fluorosurfactants in turbot. <i>Aquatic Toxicology</i> , 2011, 104, 168-176.	4.0	58
98	Novel brominated flame retardants: A review of their analysis, environmental fate and behaviour. <i>Environment International</i> , 2011, 37, 532-556.	10.0	1,188
99	A first evaluation of the usefulness of feathers of nestling predatory birds for non-destructive biomonitoring of persistent organic pollutants. <i>Environment International</i> , 2011, 37, 622-630.	10.0	73
100	Differences between Arctic and Atlantic fjord systems on bioaccumulation of persistent organic pollutants in zooplankton from Svalbard. <i>Science of the Total Environment</i> , 2011, 409, 2783-2795.	8.0	50
101	Environmental pollutants in the Swedish marine ecosystem, with special emphasis on polybrominated diphenyl ethers (PBDE). <i>Chemosphere</i> , 2011, 82, 1286-1292.	8.2	27
102	Influence of season, location, and feeding strategy on bioaccumulation of halogenated organic contaminants in Arctic marine zooplankton. <i>Environmental Toxicology and Chemistry</i> , 2011, 30, 77-87.	4.3	45
103	Seasonality in contaminant accumulation in Arctic marine pelagic food webs using trophic magnification factor as a measure of bioaccumulation. <i>Environmental Toxicology and Chemistry</i> , 2011, 30, 1026-1035.	4.3	71
104	Brominated flame retardants in the Arctic environment â€” trends and new candidates. <i>Science of the Total Environment</i> , 2010, 408, 2885-2918.	8.0	632
105	Fluctuating wing asymmetry and hepatic concentrations of persistent organic pollutants are associated in European shag (<i>Phalacrocorax aristotelis</i>) chicks. <i>Science of the Total Environment</i> , 2010, 408, 578-585.	8.0	29
106	Current Levels and Trends of Brominated Flame Retardants in the Environment. <i>Handbook of Environmental Chemistry</i> , 2010, , 123-140.	0.4	5
107	Relationships between organohalogen contaminants and blood plasma clinicalâ€“chemical parameters in chicks of three raptor species from Northern Norway. <i>Ecotoxicology and Environmental Safety</i> , 2010, 73, 7-17.	6.0	52
108	Strongly increasing blood concentrations of lipid-soluble organochlorines in high arctic common eiders during incubation fast. <i>Chemosphere</i> , 2010, 79, 320-325.	8.2	59

#	ARTICLE	IF	CITATIONS
109	Bioaccumulation of Brominated Flame Retardants. Handbook of Environmental Chemistry, 2010, , 141-185.	0.4	5
110	Salmon Farms as a Source of Organohalogenated Contaminants in Wild Fish. Environmental Science & Technology, 2010, 44, 8736-8743.	10.0	29
111	Perfluorinated and other persistent halogenated organic compounds in European shag (<i>Phalacrocorax aristotelis</i>) and common eider (<i>Somateria mollissima</i>) from Norway: A suburban to remote pollutant gradient. Science of the Total Environment, 2009, 408, 340-348.	8.0	88
112	Brominated flame retardants in the European chemicals policy of REACH—Regulation and determination in materials. Journal of Chromatography A, 2009, 1216, 320-333.	3.7	198
113	Perfluorinated, brominated, and chlorinated contaminants in a population of lesser black-backed gulls (<i>Larus fuscus</i>). Environmental Toxicology and Chemistry, 2008, 27, 1383-1392.	4.3	36
114	Monitoring of Raptors and Their Contamination Levels in Norway. Ambio, 2008, 37, 420-424.	5.5	17
115	Perfluorinated and chlorinated pollutants as predictors of demographic parameters in an endangered seabird. Environmental Pollution, 2008, 156, 417-424.	7.5	51
116	Detailed analysis of polybrominated biphenyl congeners in bird eggs from Norway. Environmental Pollution, 2008, 156, 1204-1210.	7.5	27
117	Levels and trends of HBCD and BDEs in the European and Asian environments, with some information for other BFRs. Chemosphere, 2008, 73, 223-241.	8.2	234
118	Natural and man-made organobromine compounds in marine biota from Central Norway. Environment International, 2007, 33, 17-26.	10.0	38
119	PERFLUORINATED, BROMINATED AND CHLORINATED CONTAMINANTS IN A POPULATION OF LESSER BLACK-BACKED GULLS. Environmental Toxicology and Chemistry, 2007, preprint, 1.	4.3	2
120	The BEEP Stavanger Workshop: Mesocosm exposures. Aquatic Toxicology, 2006, 78, S5-S12.	4.0	25
121	Levels and trends of brominated flame retardants in the European environment. Chemosphere, 2006, 64, 187-208.	8.2	720
122	Analytical strategies for successful enantioselective separation of atropisomeric polybrominated biphenyls 132 and 149 in environmental samples. Journal of Chromatography A, 2005, 1063, 193-199.	3.7	15
123	Determination of the enantiomer fraction of PBB 149 by gas chromatography/electron capture negative ionization tandem mass spectrometry in the selected reaction monitoring mode. Rapid Communications in Mass Spectrometry, 2005, 19, 3719-3723.	1.5	11
124	Two Trace Analytical Methods for Determination of Hydroxylated PCBs and Other Halogenated Phenolic Compounds in Eggs from Norwegian Birds of Prey. Analytical Chemistry, 2004, 76, 441-452.	6.5	98
125	Polychlorinated camphenes (toxaphenes), polybrominated diphenylethers and other halogenated organic pollutants in glaucous gull (<i>Larus hyperboreus</i>) from Svalbard and Bj�rn�ya (Bear Island). Environmental Pollution, 2003, 121, 293-300.	7.5	66
126	BFR—governmental testing programme. Environment International, 2003, 29, 781-792.	10.0	53

#	ARTICLE	IF	CITATIONS
127	Kinetics and organotropy of some polyfluorinated dibenzo-p-dioxins and dibenzofurans (PFDD/PDFD) in rats. <i>Life Sciences</i> , 2002, 71, 1475-1486.	4.3	12
128	Organochlorines in egg samples from Norwegian birds of prey: Congener-, isomer- and enantiomer specific considerations. <i>Science of the Total Environment</i> , 2002, 291, 59-71.	8.0	72
129	Identification of toxaphene congeners in bird eggs by combining quadrupole NICI-mS and ion trap EI-MS/MS. <i>Journal of Separation Science</i> , 2002, 25, 453-461.	2.5	6