

Diego Rubiales

List of Publications by Year
in descending order

Source: <https://exaly.com/author-pdf/2224483/publications.pdf>

Version: 2024-02-01

366
papers

14,092
citations

23567
58
h-index

48315
88
g-index

376
all docs

376
docs citations

376
times ranked

7764
citing authors

#	ARTICLE	IF	CITATIONS
1	Nanotechnology for parasitic plant control. Pest Management Science, 2009, 65, 540-545.	3.4	347
2	Nanoparticle penetration and transport in living pumpkin plants: in situsubcellular identification. BMC Plant Biology, 2009, 9, 45.	3.6	331
3	Nanoparticles as Smart Treatment-delivery Systems in Plants: Assessment of Different Techniques of Microscopy for their Visualization in Plant Tissues. Annals of Botany, 2008, 101, 187-195.	2.9	303
4	Biotechnology approaches to overcome biotic and abiotic stress constraints in legumes. Euphytica, 2006, 147, 1-24.	1.2	214
5	Cereal landraces for sustainable agriculture. A review. Agronomy for Sustainable Development, 2010, 30, 237-269.	5.3	197
6	Integrated pest management in faba bean. Field Crops Research, 2010, 115, 308-318.	5.1	174
7	Pea (<i>Pisum sativum</i> L.) in the Genomic Era. Agronomy, 2012, 2, 74-115.	3.0	172
8	Absorption and translocation to the aerial part of magnetic carbon-coated nanoparticles through the root of different crop plants. Journal of Nanobiotechnology, 2010, 8, 26.	9.1	159
9	Screening techniques and sources of resistance to foliar diseases caused by major necrotrophic fungi in grain legumes. Euphytica, 2006, 147, 223-253.	1.2	154
10	Biology and Management of Weedy Root Parasites. , 2007, , 267-349.		154
11	Achievements and Challenges in Legume Breeding for Pest and Disease Resistance. Critical Reviews in Plant Sciences, 2015, 34, 195-236.	5.7	153
12	Screening techniques and sources of resistance against parasitic weeds in grain legumes. Euphytica, 2006, 147, 187-199.	1.2	137
13	Potentially durable resistance mechanisms in plants to specialised fungal pathogens. Euphytica, 2002, 124, 201-216.	1.2	136
14	Lathyrus improvement for resistance against biotic and abiotic stresses: From classical breeding to marker assisted selection. Euphytica, 2006, 147, 133-147.	1.2	133
15	Faba bean breeding for disease resistance. Field Crops Research, 2010, 115, 297-307.	5.1	128
16	Characterization of <i>Lr34</i> , a Major Gene Conferring Nonhypersensitive Resistance to Wheat Leaf Rust. Plant Disease, 1995, 79, 1208.	1.4	128
17	Recognition of root exudates by seeds of broomrape (<i>Orobanche</i> and <i>Phelipanche</i>) species. Annals of Botany, 2009, 103, 423-431.	2.9	110
18	Innovations in parasitic weeds management in legume crops. A review. Agronomy for Sustainable Development, 2012, 32, 433-449.	5.3	109

#	ARTICLE	IF	CITATIONS
19	Orobanche crenata resistance and avoidance in pea (Pisum spp.) operate at different developmental stages of the parasite. Weed Research, 2005, 45, 379-387.	1.7	107
20	Faba bean breeding for resistance against biotic stresses: Towards application of marker technology. Euphytica, 2006, 147, 67-80.	1.2	104
21	Mapping of quantitative trait loci controlling broomrape (Orobanche crenataForsk.) resistance in faba bean (Vicia fabaL.). Genome, 2002, 45, 1057-1063.	2.0	103
22	Revisiting strategies for reducing the seedbank of <i>Orobanche</i> and <i>Phelipanche</i> spp.. Weed Research, 2009, 49, 23-33.	1.7	103
23	Isolate and organ-specific QTLs for ascochyta blight resistance in faba bean (Vicia faba L).. Theoretical and Applied Genetics, 2004, 108, 1071-1078.	3.6	94
24	Interaction between Orobanche crenata and its Host Legumes: Unsuccessful Haustorial Penetration and Necrosis of the Developing Parasite. Annals of Botany, 2005, 95, 935-942.	2.9	93
25	The role of strigolactones in host specificity of<i>Orobanche</i>and<i>Phelipanche</i>seed germination. Seed Science Research, 2011, 21, 55-61.	1.7	92
26	A proteomic approach to study pea (Pisum sativum) responses to powdery mildew (Erysiphe pisi). Proteomics, 2006, 6, S163-S174.	2.2	90
27	Screening techniques and sources of resistance to rusts and mildews in grain legumes. Euphytica, 2006, 147, 255-272.	1.2	90
28	Macroscopic and Histological Characterisation of Genes er1 and er2 for Powdery Mildew Resistance in Pea. European Journal of Plant Pathology, 2006, 115, 309-321.	1.7	89
29	Plant resistance to parasitic plants: molecular approaches to an old foe. New Phytologist, 2007, 173, 703-712.	7.3	89
30	Genetic mapping of QTLs controlling horticultural traits in diploid roses. Theoretical and Applied Genetics, 2005, 111, 511-520.	3.6	88
31	Characterization of resistance in chickpea to crenate broomrape (Orobanche crenata). Weed Science, 2003, 51, 702-707.	1.5	86
32	Response to Mycosphaerella pinodes in a germplasm collection of Pisum spp. Plant Breeding, 2005, 124, 313-315.	1.9	86
33	Identification of a New Gene for Resistance to Powdery Mildew in Pisum fulvum, a Wild Relative of Pea. Breeding Science, 2007, 57, 181-184.	1.9	84
34	Powdery mildew control in pea. A review. Agronomy for Sustainable Development, 2012, 32, 401-409.	5.3	84
35	Genetic basis of qualitative and quantitative resistance to powdery mildew in wheat: from consensus regions to candidate genes. BMC Genomics, 2013, 14, 562.	2.8	84
36	A proteomic approach to studying plant response to crenate broomrape (Orobanche crenata) in pea (Pisum sativum). Phytochemistry, 2004, 65, 1817-1828.	2.9	83

#	ARTICLE	IF	CITATIONS
37	Intercropping with cereals reduces infection by <i>Orobanche crenata</i> in legumes. <i>Crop Protection</i> , 2007, 26, 1166-1172.	2.1	83
38	Broomrape management in faba bean. <i>Field Crops Research</i> , 2010, 115, 319-328.	5.1	79
39	Reduced nitric oxide levels during drought stress promote drought tolerance in barley and is associated with elevated polyamine biosynthesis. <i>Scientific Reports</i> , 2017, 7, 13311.	3.3	79
40	Histological Characterization of Resistance to <i>Uromyces viciae-fabae</i> in Faba Bean. <i>Phytopathology</i> , 2002, 92, 294-299.	2.2	78
41	Crenate broomrape (<i>Orobanche crenata</i>) infection in field pea cultivars. <i>Crop Protection</i> , 2003, 22, 865-872.	2.1	78
42	Identification of RAPD markers linked to the Uvf-1 gene conferring hypersensitive resistance against rust (<i>Uromyces viciae-fabae</i>) in <i>Vicia faba</i> L.. <i>Theoretical and Applied Genetics</i> , 2003, 107, 353-358.	3.6	77
43	Identification of genes differentially expressed in a resistant reaction to <i>Mycosphaerella pinodes</i> in pea using microarray technology. <i>BMC Genomics</i> , 2011, 12, 28.	2.8	77
44	Protein cross-linking, peroxidase and β -1,3-endoglucanase involved in resistance of pea against <i>Orobanche crenata</i> . <i>Journal of Experimental Botany</i> , 2006, 57, 1461-1469.	4.8	75
45	Mucilage production during the incompatible interaction between <i>Orobanche crenata</i> and <i>Vicia sativa</i> . <i>Journal of Experimental Botany</i> , 2006, 57, 931-942.	4.8	74
46	<i>Lathyrus</i> diversity: available resources with relevance to crop improvement – <i>L. sativus</i> and <i>L. cicera</i> as case studies. <i>Annals of Botany</i> , 2014, 113, 895-908.	2.9	74
47	A metabolomic study in oats (<i>Avena sativa</i>) highlights a drought tolerance mechanism based upon salicylate signalling pathways and the modulation of carbon, antioxidant and photooxidative metabolism. <i>Plant, Cell and Environment</i> , 2015, 38, 1434-1452.	5.7	73
48	Introduction: Legumes in Sustainable Agriculture. <i>Critical Reviews in Plant Sciences</i> , 2015, 34, 2-3.	5.7	73
49	Breeding approaches for crenate broomrape (<i>Orobanche crenata</i> Forsk.) management in pea (<i>Pisum sativum</i> L.). <i>Pest Management Science</i> , 2009, 65, 553-559.	3.4	71
50	Quantum Dot and Superparamagnetic Nanoparticle Interaction with Pathogenic Fungi: Internalization and Toxicity Profile. <i>ACS Applied Materials & Interfaces</i> , 2014, 6, 9100-9110.	8.0	71
51	Parasitic plant management in sustainable agriculture. <i>Weed Research</i> , 2009, 49, 1-5.	1.7	69
52	Fatty Acid Profile Changes During Gradual Soil Water Depletion in Oats Suggests a Role for Jasmonates in Coping With Drought. <i>Frontiers in Plant Science</i> , 2018, 9, 1077.	3.6	69
53	Parasitic plants, wild relatives and the nature of resistance. <i>New Phytologist</i> , 2003, 160, 459-461.	7.3	68
54	Acibenzolar-S-methyl-induced resistance to sunflower rust (<i>Puccinia helianthi</i>) is associated with an enhancement of coumarins on foliar surface. <i>Physiological and Molecular Plant Pathology</i> , 2002, 60, 155-162.	2.5	65

#	ARTICLE	IF	CITATIONS
55	Model legumes contribute to faba bean breeding. <i>Field Crops Research</i> , 2010, 115, 253-269.	5.1	64
56	Mapping of quantitative trait loci for resistance to <i>Mycosphaerella pinodes</i> in <i>Pisum sativum</i> subsp. <i>syriacum</i> . <i>Molecular Breeding</i> , 2008, 21, 439-454.	2.1	62
57	Low Strigolactone Root Exudation: A Novel Mechanism of Broomrape (<i>Orobanche</i> and <i>Tj ETQq1</i> 1 0.784314 rgBT /Overlock 18 Chemistry, 2014, 62, 7063-7071.	5.2	62
58	Locating genes associated with <i>Ascochyta fabae</i> resistance in <i>Vicia faba</i> . <i>Australian Journal of Agricultural Research</i> , 2003, 54, 85.	1.5	61
59	Search for Resistance to Crenate Broomrape (<i>Orobanche crenata</i> Forsk.) in Pea Germplasm. <i>Genetic Resources and Crop Evolution</i> , 2005, 52, 853-861.	1.6	60
60	Understanding <i>Orobanche</i> and <i>Phelipanche</i> "host plant interactions and developing resistance. <i>Weed Research</i> , 2009, 49, 8-22.	1.7	60
61	Identification of quantitative trait loci for specific mechanisms of resistance to <i>Orobanche crenata</i> Forsk. in pea (<i>Pisum sativum</i> L.). <i>Molecular Breeding</i> , 2010, 25, 259-272.	2.1	60
62	Regiolone and Isosclerone, Two Enantiomeric Phytotoxic Naphthalenone Pentaketides: Computational Assignment of Absolute Configuration and Its Relationship with Phytotoxic Activity. <i>European Journal of Organic Chemistry</i> , 2011, 2011, 5564-5570.	2.4	60
63	Identification and validation of RAPD and SCAR markers linked to the gene <i>Er3</i> conferring resistance to <i>Erysiphe pisi</i> DC in pea. <i>Molecular Breeding</i> , 2008, 22, 193-200.	2.1	59
64	A High-Density Integrated DArTseq SNP-Based Genetic Map of <i>Pisum fulvum</i> and Identification of QTLs Controlling Rust Resistance. <i>Frontiers in Plant Science</i> , 2018, 9, 167.	3.6	58
65	Physical and Chemical Barriers in Root Tissues Contribute to Quantitative Resistance to <i>Fusarium oxysporum</i> f. sp. <i>pisi</i> in Pea. <i>Frontiers in Plant Science</i> , 2018, 9, 199.	3.6	58
66	Characterization of new sources of resistance to <i>Uromyces viciae-fabae</i> in a germplasm collection of <i>Vicia faba</i> . <i>Plant Pathology</i> , 2000, 49, 389-395.	2.4	57
67	Colonisation of field pea roots by arbuscular mycorrhizal fungi reduces <i>Orobanche</i> and <i>Phelipanche</i> species seed germination. <i>Weed Research</i> , 2010, 50, 262-268.	1.7	57
68	Characterization of <i>Lr46</i> , a Gene Conferring Partial Resistance to Wheat Leaf Rust. <i>Hereditas</i> , 2004, 135, 111-114.	1.4	56
69	Identification and characterization of sources of resistance to <i>Erysiphe pisi</i> Syd. in <i>Pisum</i> spp.. <i>Plant Breeding</i> , 2007, 126, 113-119.	1.9	56
70	Proteomics: a promising approach to study biotic interaction in legumes. A review. <i>Euphytica</i> , 2006, 147, 37-47.	1.2	55
71	Host plant resistance against broomrapes (<i>Orobanche</i> spp.): defence reactions and mechanisms of resistance. <i>Annals of Applied Biology</i> , 2008, 152, 131-141.	2.5	55
72	Genetic Diversity and Population Structure Among Oat Cultivars and Landraces. <i>Plant Molecular Biology Reporter</i> , 2013, 31, 1305-1314.	1.8	55

#	ARTICLE	IF	CITATIONS
73	Mechanism and molecular markers associated with rust resistance in a chickpea interspecific cross (<i>Cicer arietinum</i> Å— <i>Cicer reticulatum</i>). <i>European Journal of Plant Pathology</i> , 2008, 121, 43-53.	1.7	54
74	Mapping of quantitative trait loci controlling partial resistance against rust incited by <i>Uromyces pisi</i> (Pers.) Wint. in a <i>Pisum fulvum</i> L. intraspecific cross. <i>Euphytica</i> , 2010, 175, 151-159.	1.2	54
75	Locating quantitative trait loci associated with <i>Orobanche crenata</i> resistance in pea. <i>Weed Research</i> , 2004, 44, 323-328.	1.7	53
76	Identification and multi-environment validation of resistance to <i>Ascochyta fabae</i> in faba bean (<i>Vicia</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	8.1	53
77	Identification of quantitative trait loci and candidate genes for specific cellular resistance responses against <i>Didymella pinodes</i> in pea. <i>Plant Cell Reports</i> , 2014, 33, 1133-1145.	5.6	53
78	Resistance against barley leaf rust (<i>Puccinia hordei</i>) in West-European spring barley germplasm. <i>Agronomy for Sustainable Development</i> , 2000, 20, 769-782.	0.8	53
79	<i>Medicago truncatula</i> as a Model for Nonhost Resistance in Legume-Parasitic Plant Interactions. <i>Plant Physiology</i> , 2007, 145, 437-449.	4.8	52
80	Peagol and peagoldione, two new strigolactone-like metabolites isolated from pea root exudates. <i>Tetrahedron Letters</i> , 2009, 50, 6955-6958.	1.4	52
81	A detailed evaluation method to identify sources of quantitative resistance to <i>Fusarium oxysporum</i> f. sp. <i>pisi</i> race 2 within a <i>Pisum</i> spp. germplasm collection. <i>Plant Pathology</i> , 2012, 61, 532-542.	2.4	52
82	Adaptation of spring faba bean types across European climates. <i>Field Crops Research</i> , 2013, 145, 1-9.	5.1	52
83	The Effect of <i>Orobanche crenata</i> Infection Severity in Faba Bean, Field Pea, and Grass Pea Productivity. <i>Frontiers in Plant Science</i> , 2016, 7, 1409.	3.6	52
84	Characterization of mechanisms of resistance against <i>Didymella pinodes</i> in <i>Pisum</i> spp.. <i>European Journal of Plant Pathology</i> , 2013, 135, 761-769.	1.7	51
85	<i>Didymella pinodes</i> and its management in field pea: Challenges and opportunities. <i>Field Crops Research</i> , 2013, 148, 61-77.	5.1	51
86	Quantitative Trait Loci Associated to Drought Adaptation in Pea (<i>Pisum sativum</i> L.). <i>Plant Molecular Biology Reporter</i> , 2015, 33, 1768-1778.	1.8	51
87	Resistance to broomrape (<i>Orobanche crenata</i>) in faba bean (<i>Vicia faba</i>): cell wall changes associated with prehaustorial defensive mechanisms. <i>Annals of Applied Biology</i> , 2007, 151, 89-98.	2.5	50
88	Control of <i>Orobanche crenata</i> in legumes intercropped with fenugreek (<i>Trigonella foenum-graecum</i>). <i>Crop Protection</i> , 2008, 27, 653-659.	2.1	50
89	Identification and multi-environment validation of resistance to <i>Botrytis fabae</i> in <i>Vicia faba</i> . <i>Field Crops Research</i> , 2009, 114, 84-90.	5.1	50
90	Identification of common genomic regions controlling resistance to <i>Mycosphaerella pinodes</i> , earliness and architectural traits in different pea genetic backgrounds. <i>Euphytica</i> , 2011, 182, 43-52.	1.2	50

#	ARTICLE	IF	CITATIONS
91	Avoidance of rust infection by some genotypes of <i>Hordeum chilense</i> due to their relative inability to induce the formation of appressoria. <i>Physiological and Molecular Plant Pathology</i> , 1996, 49, 89-101.	2.5	49
92	Infection Structures of Host-Specialized Isolates of <i>Uromyces viciae-fabae</i> and of Other Species of <i>Uromyces</i> Infecting Leguminous Crops. <i>Plant Disease</i> , 2005, 89, 17-22.	1.4	49
93	Trigoxazonane, a monosubstituted trioxazonane from <i>Trigonella foenum-graecum</i> root exudate, inhibits <i>Orobanche crenata</i> seed germination. <i>Phytochemistry</i> , 2007, 68, 2487-2492.	2.9	49
94	Polyphenols, Including the New Peapolyphenols Aa ⁺ C, from Pea Root Exudates Stimulate <i>Orobanche foetida</i> Seed Germination. <i>Journal of Agricultural and Food Chemistry</i> , 2010, 58, 2902-2907.	5.2	49
95	Faba bean adaptation to autumn sowing under European climates. <i>Agronomy for Sustainable Development</i> , 2012, 32, 727-734.	5.3	49
96	Intercropping reduces <i>Mycosphaerella pinodes</i> severity and delays upward progress on the pea plant. <i>Crop Protection</i> , 2010, 29, 744-750.	2.1	48
97	Inter-cropping with berseem clover (<i>Trifolium alexandrinum</i>) reduces infection by <i>Orobanche crenata</i> in legumes. <i>Crop Protection</i> , 2010, 29, 867-871.	2.1	48
98	Future Prospects for Ascochyta Blight Resistance Breeding in Cool Season Food Legumes. <i>Frontiers in Plant Science</i> , 2012, 3, 27.	3.6	48
99	Induction of systemic acquired resistance against rust, ascochyta blight and broomrape in faba bean by exogenous application of salicylic acid and benzothiadiazole. <i>Crop Protection</i> , 2012, 34, 65-69.	2.1	48
100	Infection of chickpea (<i>Cicer arietinum</i>) by crenate broomrape (<i>Orobanche crenata</i>) as influenced by sowing date and weather conditions. <i>Agronomy for Sustainable Development</i> , 2003, 23, 359-362.	0.8	48
101	Identification of sources of resistance to crenate broomrape (<i>Orobanche crenata</i>) in Spanish lentil (<i>Lens culinaris</i>) germplasm. <i>Weed Research</i> , 2008, 48, 85-94.	1.7	47
102	Understanding pea resistance mechanisms in response to <i>Fusarium oxysporum</i> through proteomic analysis. <i>Phytochemistry</i> , 2015, 115, 44-58.	2.9	47
103	Variation Among and Within Populations of the Parasitic Weed <i>Orobanche crenata</i> from Spain and Israel Revealed by Inter Simple Sequence Repeat Markers. <i>Phytopathology</i> , 2002, 92, 1262-1266.	2.2	46
104	Characterization of wheat DArT markers: genetic and functional features. <i>Molecular Genetics and Genomics</i> , 2012, 287, 741-753.	2.1	46
105	Genetic Relationships among <i>Orobanche</i> Species as Revealed by RAPD Analysis. <i>Annals of Botany</i> , 2003, 91, 637-642.	2.9	45
106	Resistance against broomrapes (<i>Orobanche</i> and <i>Phelipanche</i> spp.) in faba bean (<i>Vicia faba</i>) based in low induction of broomrape seed germination. <i>Euphytica</i> , 2012, 186, 897-905.	1.2	45
107	Two-Dimensional Electrophoresis Based Proteomic Analysis of the Pea (<i>Pisum sativum</i>) in Response to <i>Mycosphaerella pinodes</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2010, 58, 12822-12832.	5.2	44
108	Confirmation that the <i>Er3</i> gene, conferring resistance to <i>Erysiphe pisi</i> in pea, is a different gene from <i>er1</i> and <i>er2</i> genes. <i>Plant Breeding</i> , 2011, 130, 281-282.	1.9	44

#	ARTICLE	IF	CITATIONS
109	Identification and multi-environment validation of resistance against broomrapes (<i>Orobanche</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10	5.1	44
110	Adaptation of oat (<i>Avena sativa</i>) cultivars to autumn sowings in Mediterranean environments. Field Crops Research, 2014, 156, 111-122.	5.1	44
111	Sources of Resistance to Crenate Broomrape Among Species of <i>Vicia</i> . Plant Disease, 2005, 89, 23-27.	1.4	43
112	Effect of Fungal and Plant Metabolites on Broomrapes (<i>Orobanche</i> and <i>Phelipanche</i> spp.) Seed Germination and Radicle Growth. Journal of Agricultural and Food Chemistry, 2014, 62, 10485-10492.	5.2	43
113	Genome-wide association study for crown rust (<i>Puccinia coronata</i> f. sp. <i>avenae</i>) and powdery mildew (<i>Blumeria graminis</i> f. sp. <i>avenae</i>) resistance in an oat (<i>Avena sativa</i>) collection of commercial varieties and landraces. Frontiers in Plant Science, 2015, 6, 103.	3.6	43
114	<i>Lathyrus sativus</i> transcriptome resistance response to <i>Ascochyta lathyri</i> investigated by deepSuperSAGE analysis. Frontiers in Plant Science, 2015, 6, 178.	3.6	43
115	Tritordeum: Triticale's New Brother Cereal. Developments in Plant Breeding, 1996, , 57-72.	0.2	43
116	Genetic diversity in <i>Orobanche crenata</i> populations from southern Spain. Theoretical and Applied Genetics, 2001, 103, 1108-1114.	3.6	42
117	Characterization of resistance response of pea (<i>Pisum</i> spp.) against rust (<i>Uromyces pisi</i>). Plant Breeding, 2009, 128, 665-670.	1.9	42
118	Genetic analysis of durable resistance against leaf rust in durum wheat. Molecular Breeding, 2009, 24, 25-39.	2.1	41
119	Soyasapogenol B and <i>trans</i> - Δ^2 -dehydrocampesterol from common vetch (<i>Vicia sativa</i> L.) root exudates stimulate broomrape seed germination. Pest Management Science, 2011, 67, 1015-1022.	3.4	41
120	Genome-wide identification and comparison of legume MLO gene family. Scientific Reports, 2016, 6, 32673.	3.3	41
121	Variation in resistance to <i>Orobanche crenata</i> in species of <i>Cicer</i> . Weed Research, 2004, 44, 27-32.	1.7	40
122	Resistance to broomrape in wild lentils (<i>Lens</i> spp.). Plant Breeding, 2009, 128, 266-270.	1.9	40
123	Benzothiadiazole and BABA improve resistance to <i>Uromyces pisi</i> (Pers.) Wint. in <i>Pisum sativum</i> L. with an enhancement of enzymatic activities and total phenolic content. European Journal of Plant Pathology, 2010, 128, 483-493.	1.7	40
124	Agronomic, breeding, and biotechnological approaches to parasitic plant management through manipulation of germination stimulant levels in agricultural soils. Botany, 2011, 89, 813-826.	1.0	40
125	Identification of the Main Toxins Isolated from <i>Fusarium oxysporum</i> f. sp. <i>pisi</i> Race 2 and Their Relation with Isolates' Pathogenicity. Journal of Agricultural and Food Chemistry, 2014, 62, 2574-2580.	5.2	40
126	Identification of resistance to <i>Uromyces pisi</i> (Pers.) Wint. in <i>Pisum</i> spp. germplasm. Field Crops Research, 2009, 114, 198-203.	5.1	39

#	ARTICLE	IF	CITATIONS
127	Chemical control of faba bean rust (<i>Uromyces viciae-fabae</i>). <i>Crop Protection</i> , 2011, 30, 907-912.	2.1	39
128	Effects of crop mixtures on chocolate spot development on faba bean grown in mediterranean climates. <i>Crop Protection</i> , 2011, 30, 1015-1023.	2.1	39
129	QTLs for <i>Orobanche</i> spp. resistance in faba bean: identification and validation across different environments. <i>Molecular Breeding</i> , 2013, 32, 909-922.	2.1	39
130	Field response of <i>Lathyrus cicera</i> germplasm to crenate broomrape (<i>Orobanche crenata</i>). <i>Field Crops Research</i> , 2009, 113, 321-327.	5.1	38
131	<i>Hordeum chilense</i> resistance to powdery mildew and its potential use in cereal breeding. <i>Euphytica</i> , 1993, 67, 215-220.	1.2	37
132	Effect of sowing date and host resistance on the establishment and development of <i>Orobanche crenata</i> in faba bean and common vetch. <i>Weed Research</i> , 2004, 44, 282-288.	1.7	37
133	Crenate broomrape control in pea by foliar application of benzothiadiazole (BTH). <i>Phytoparasitica</i> , 2004, 32, 21-29.	1.2	37
134	Identification of QTLs for powdery mildew and scald resistance in barley. <i>Euphytica</i> , 2006, 151, 421-429.	1.2	37
135	Identification by suppression subtractive hybridization and expression analysis of <i>Medicago truncatula</i> putative defence genes in response to <i>Orobanche crenata</i> parasitization. <i>Physiological and Molecular Plant Pathology</i> , 2007, 70, 49-59.	2.5	37
136	Inhibition of <i>Orobanche crenata</i> Seed Germination and Radicle Growth by Allelochemicals Identified in Cereals. <i>Journal of Agricultural and Food Chemistry</i> , 2013, 61, 9797-9803.	5.2	37
137	Allelic diversity in the transcriptomes of contrasting rust-infected genotypes of <i>Lathyrus sativus</i> , a lasting resource for smart breeding. <i>BMC Plant Biology</i> , 2014, 14, 376.	3.6	37
138	Genetic analysis of root morphological traits in wheat. <i>Molecular Genetics and Genomics</i> , 2015, 290, 785-806.	2.1	37
139	Resistance to broomrape species (<i>Orobanche</i> spp.) in common vetch (<i>Vicia sativa</i> L.). <i>Crop Protection</i> , 2009, 28, 7-12.	2.1	36
140	First Report of <i>Orobanche foetida</i> on Common Vetch (<i>Vicia sativa</i>) in Morocco. <i>Plant Disease</i> , 2005, 89, 528-528.	1.4	36
141	Characterization of resistance to powdery mildew (<i>Erysiphe pisi</i>) in a germplasm collection of <i>Lathyrus sativus</i> . <i>Plant Breeding</i> , 2006, 125, 308-310.	1.9	35
142	Host differentiation in <i>Orobanche foetida</i> Poir. <i>Flora: Morphology, Distribution, Functional Ecology of Plants</i> , 2007, 202, 201-208.	1.2	35
143	The resistance to leaf rust and powdery mildew of recombinant lines of barley (<i>Hordeum vulgare</i> L.) derived from <i>H. v. vulgare</i> × <i>H. v. bulbosum</i> crosses. <i>Plant Breeding</i> , 2007, 126, 259-267.	1.9	35
144	Variability of interactions between barrel medic (<i>Medicago truncatula</i>) genotypes and <i>Orobanche</i> species. <i>Annals of Applied Biology</i> , 2008, 153, 117-126.	2.5	35

#	ARTICLE	IF	CITATIONS
145	Resistance to powdery mildew (<i>Blumeria graminis</i> f.sp. <i>avenae</i>) in oat seedlings and adult plants. <i>Plant Pathology</i> , 2011, 60, 846-856.	2.4	35
146	Proteomic Analysis of Pea (<i>Pisum sativum</i> L.) Response During Compatible and Incompatible Interactions with the Pea Aphid (<i>Acyrtosiphon pisum</i> H.). <i>Plant Molecular Biology Reporter</i> , 2014, 32, 697-718.	1.8	35
147	Molecular and cytogenetic characterization of a common wheat- <i>Agropyron cristatum</i> chromosome translocation conferring resistance to leaf rust. <i>Euphytica</i> , 2015, 201, 89-95.	1.2	35
148	Genetic diversity of Moroccan populations of <i>Orobancha foetida</i> : evolving from parasitising wild hosts to crop plants. <i>Weed Research</i> , 2008, 48, 179-186.	1.7	34
149	Transferability of molecular markers from major legumes to <i>Lathyrus</i> spp. for their application in mapping and diversity studies. <i>Molecular Biology Reports</i> , 2014, 41, 269-283.	2.3	34
150	Prehaustorial Resistance against Alfalfa Rust (<i>Uromyces striatus</i>) in <i>Medicago truncatula</i> . <i>European Journal of Plant Pathology</i> , 2004, 110, 239-243.	1.7	33
151	Constitutive Coumarin Accumulation on Sunflower Leaf Surface Prevents Rust Germ Tube Growth and Appressorium Differentiation. <i>Crop Science</i> , 2007, 47, 1119-1124.	1.8	33
152	Fenugreek root exudates show species-specific stimulation of <i>Orobancha</i> seed germination. <i>Weed Research</i> , 2008, 48, 163-168.	1.7	33
153	Stimulation of Seed Germination of <i>Orobancha</i> Species by Ophiobolin A and Fusicoccin Derivatives. <i>Journal of Agricultural and Food Chemistry</i> , 2008, 56, 8343-8347.	5.2	33
154	Differential response of pea (<i>Pisum sativum</i>) to rusts incited by <i>Uromyces viciae-fabae</i> and <i>U. pisi</i> . <i>Crop Protection</i> , 2009, 28, 980-986.	2.1	33
155	Identification and characterization of partial resistance to rust in a germplasm collection of <i>Lathyrus sativus</i> L.. <i>Plant Breeding</i> , 2009, 128, 495-500.	1.9	33
156	Proteomic analysis by two-dimensional differential in gel electrophoresis (2D DIGE) of the early response of <i>Pisum sativum</i> to <i>Orobancha crenata</i> . <i>Journal of Experimental Botany</i> , 2012, 63, 107-119.	4.8	33
157	Identification and mapping of quantitative trait loci for leaf rust resistance derived from a tetraploid wheat <i>Triticum dicoccum</i> accession. <i>Molecular Breeding</i> , 2014, 34, 1659-1675.	2.1	33
158	A fertile amphiploid between diploid wheat (<i>Triticum tauschii</i>) and crested wheatgrass (<i>Agropyron cristatum</i>). <i>Genome</i> , 1999, 42, 519-524.	2.0	32
159	Morphology and AFLP markers suggest three <i>Hordeum chilense</i> ecotypes that differ in avoidance to rust fungi. <i>Canadian Journal of Botany</i> , 2001, 79, 204-213.	1.1	32
160	Pathogenic Specialization of <i>Puccinia triticina</i> in Andalusia from 1998 to 2000. <i>Journal of Phytopathology</i> , 2005, 153, 344-349.	1.0	32
161	Transformation and regeneration of the holoparasitic plant <i>Phelipanche aegyptiaca</i> . <i>Plant Methods</i> , 2011, 7, 36.	4.3	32
162	Fusarium Wilt Management in Legume Crops. <i>Agronomy</i> , 2020, 10, 1073.	3.0	32

#	ARTICLE	IF	CITATIONS
163	Morphology and AFLP markers suggest three <i>Hordeum chilense</i> ecotypes that differ in avoidance to rust fungi. Canadian Journal of Botany, 2001, 79, 204-213.	1.1	32
164	Higher rust resistance and similar yield of oat landraces versus cultivars under high temperature and drought. Agronomy for Sustainable Development, 2017, 37, 1.	5.3	31
165	Quantitative Analysis of Target Peptides Related to Resistance Against <i>Ascochyta</i> Blight (<i>Peyronellaea pinodes</i>) in Pea. Journal of Proteome Research, 2020, 19, 1000-1012.	3.7	31
166	Grain Yield Stability of Cereal-Legume Intercrops Is Greater Than Sole Crops in More Productive Conditions. Agriculture (Switzerland), 2021, 11, 255.	3.1	31
167	Comparative proteomic analysis of BTH and BABA-induced resistance in pea (<i>Pisum sativum</i>) toward infection with pea rust (<i>Uromyces pisi</i>). Journal of Proteomics, 2012, 75, 5189-5205.	2.4	30
168	Pinolide, a New Nonenolide Produced by <i>Didymella pinodes</i> , the Causal Agent of <i>Ascochyta</i> Blight on <i>Pisum sativum</i> . Journal of Agricultural and Food Chemistry, 2012, 60, 5273-5278.	5.2	30
169	Identification of resistance to <i>Ascochyta fabae</i> in <i>Vicia faba</i> germplasm. Plant Breeding, 2001, 120, 529-531.	1.9	29
170	Characterization of the <i>Orobanche-Medicago truncatula</i> association for studying early stages of the parasite-host interaction. Weed Research, 2004, 44, 218-223.	1.7	29
171	Characterization of Resistance Mechanisms to <i>Erysiphe pisi</i> in <i>Medicago truncatula</i>. Phytopathology, 2007, 97, 1049-1053.	2.2	29
172	New Chemical Clues for Broomrape-Sunflower Host-Parasite Interactions: Synthesis of Guaianestrinolactones. Journal of Agricultural and Food Chemistry, 2009, 57, 5853-5864.	5.2	29
173	Escape and true resistance to crenate broomrape (<i>Orobanche crenata</i> Forsk.) in grass pea (<i>Lathyrus</i>) Tj ETQq1 1 0.784314 rgBT /Overl	5.1	29
174	<i>Erysiphe trifolii</i> is able to overcome <i>er1</i> and <i>Er3</i> , but not <i>er2</i> , resistance genes in pea. European Journal of Plant Pathology, 2013, 136, 557-563.	1.7	29
175	Induction of Haustorium Development by <i>Sphaeropsidones</i> in Radicles of the Parasitic Weeds <i>Striga</i> and <i>Orobanche</i>. A Structure-Activity Relationship Study. Journal of Agricultural and Food Chemistry, 2016, 64, 5188-5196.	5.2	29
176	Legume breeding for rust resistance: lessons to learn from the model <i>Medicago truncatula</i> . Euphytica, 2011, 180, 89-98.	1.2	28
177	Screening faba bean for chocolate spot resistance: evaluation methods and effects of age of host tissue and temperature. European Journal of Plant Pathology, 2012, 132, 443-453.	1.7	28
178	Clarification on Host Range of <i>Didymella pinodes</i> the Causal Agent of Pea <i>Ascochyta</i> Blight. Frontiers in Plant Science, 2016, 7, 592.	3.6	28
179	Natural Variation in Portuguese Common Bean Germplasm Reveals New Sources of Resistance Against <i>Fusarium oxysporum</i> f. sp. <i>phaseoli</i> and Resistance-Associated Candidate Genes. Phytopathology, 2020, 110, 633-647.	2.2	28
180	Omics resources and omics-enabled approaches for achieving high productivity and improved quality in pea (<i>Pisum sativum</i> L.). Theoretical and Applied Genetics, 2021, 134, 755-776.	3.6	28

#	ARTICLE	IF	CITATIONS
181	Low Appressorium Formation by Rust Fungi on <i>Hordeum chilense</i> Lines. <i>Phytopathology</i> , 1992, 82, 1007.	2.2	28
182	Disease resistance in pea (<i>Pisum sativum</i> L.) types for autumn sowings in Mediterranean environments - a review. <i>Czech Journal of Genetics and Plant Breeding</i> , 2009, 45, 135-142.	0.8	27
183	Botrytone, a New Naphthalenone Pentaketide Produced by <i>Botrytis fabae</i> , the Causal Agent of Chocolate Spot Disease on <i>Vicia faba</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2011, 59, 9201-9206.	5.2	27
184	Identification of Sources of Quantitative Resistance to <i>Fusarium oxysporum</i> f. sp. <i>medicaginis</i> in <i>Medicago truncatula</i> . <i>Plant Disease</i> , 2014, 98, 667-673.	1.4	27
185	Models, Developments, and Perspectives of Mutual Legume Intercropping. <i>Advances in Agronomy</i> , 2015, 130, 337-419.	5.2	27
186	Gene expression profiling of <i>Medicago truncatula</i> roots in response to the parasitic plant <i>Orobancha crenata</i> . <i>Weed Research</i> , 2009, 49, 66-80.	1.7	26
187	Induction of Systemic Acquired Resistance in Pea against Rust (<i>Uromyces pisi</i>) by Exogenous Application of Biotic and Abiotic Inducers. <i>Journal of Phytopathology</i> , 2010, 158, 30-34.	1.0	26
188	BTH and BABA induce resistance in pea against rust (<i>Uromyces pisi</i>) involving differential phytoalexin accumulation. <i>Planta</i> , 2015, 242, 1095-1106.	3.2	26
189	Histopathology of the infection on resistant and susceptible lentil accessions by two contrasting pathotypes of <i>Fusarium oxysporum</i> f.sp. <i>lentis</i> . <i>European Journal of Plant Pathology</i> , 2017, 148, 53-63.	1.7	26
190	Understanding photothermal interactions will help expand production range and increase genetic diversity of lentil (<i>Lens culinaris</i> Medik.). <i>Plants People Planet</i> , 2021, 3, 171-181.	3.3	26
191	Population genetics in weedy species of <i>Orobancha</i> . <i>Australasian Plant Pathology</i> , 2009, 38, 228.	1.0	25
192	Ryecyanatines A and B and ryecarbonitrilines A and B, substituted cyanatophenol, cyanatobenzo[1,3]dioxole, and benzo[1,3]dioxolecarbonitriles from rye (<i>Secale cereale</i> L.) root exudates: Novel metabolites with allelopathic activity on <i>Orobancha</i> seed germination and radicle growth. <i>Phytochemistry</i> , 2015, 109, 57-65.	2.9	25
193	QTLs for ascochyta blight resistance in faba bean (<i>Vicia faba</i> L.): validation in field and controlled conditions. <i>Crop and Pasture Science</i> , 2016, 67, 216.	1.5	25
194	Legume Breeding for the Agroecological Transition of Global Agri-Food Systems: A European Perspective. <i>Frontiers in Plant Science</i> , 2021, 12, 782574.	3.6	25
195	Antifungal Activity of a New Phenolic Compound from Capitulum of a Head Rot-resistant Sunflower Genotype. <i>Journal of Chemical Ecology</i> , 2007, 33, 2245-2253.	1.8	24
196	Pre and posthaustorial resistance to rusts in <i>Lathyrus cicera</i> L.. <i>Euphytica</i> , 2009, 165, 27-34.	1.2	24
197	Identification of Genes Involved in Resistance to <i>Didymella pinodes</i> in Pea by deepSuperSAGE Transcriptome Profiling. <i>Plant Molecular Biology Reporter</i> , 2014, 32, 258-269.	1.8	24
198	Identification of quantitative trait loci involved in resistance to <i>Pseudomonas syringae</i> pv. <i>syringae</i> in pea (<i>Pisum sativum</i> L.). <i>Euphytica</i> , 2012, 186, 805-812.	1.2	23

#	ARTICLE	IF	CITATIONS
199	Genetic resistance to powdery mildew in common bean. <i>Euphytica</i> , 2012, 186, 875-882.	1.2	23
200	Lentisone, a New Phytotoxic Anthraquinone Produced by <i>Ascochyta lentis</i> , the Causal Agent of <i>Ascochyta</i> Blight in <i>Lens culinaris</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2013, 61, 7301-7308.	5.2	23
201	Complexation of sesquiterpene lactones with cyclodextrins: synthesis and effects on their activities on parasitic weeds. <i>Organic and Biomolecular Chemistry</i> , 2017, 15, 6500-6510.	2.8	23
202	Defence reactions of <i>Hordeum chilense</i> accessions to three formae speciales of cereal powdery mildew fungi. <i>Canadian Journal of Botany</i> , 2000, 78, 1561-1570.	1.1	23
203	The reaction of <i>x Tritordeum</i> and its <i>Triticum</i> spp. and <i>Hordeum chilense</i> parents to rust diseases. <i>Euphytica</i> , 1991, 54, 75-81.	1.2	22
204	Genetic Variation Among and Within <i>Uromyces</i> Species Infecting Legumes. <i>Journal of Phytopathology</i> , 2008, 156, 419-424.	1.0	22
205	Identification and characterization of sources of resistance in <i>Avena sativa</i> , <i>A. abyssinica</i> and <i>A. sativa</i> germplasm against a pathotype of <i>Puccinia coronata</i> f.sp. <i>avenae</i> with virulence against the <i>Pc94</i> resistance gene. <i>Plant Pathology</i> , 2012, 61, 315-322.	2.4	22
206	Response of vetches (<i>Vicia</i> spp.) to specialized forms of <i>Uromyces vicia-fabae</i> and to <i>Uromyces pisi</i> . <i>Crop Protection</i> , 2013, 46, 38-43.	2.1	22
207	Identification of pathotypes in <i>Fusarium oxysporum</i> f.sp. <i>lentis</i> . <i>European Journal of Plant Pathology</i> , 2016, 144, 539-549.	1.7	22
208	Identification and multi-environment validation of resistance to pea weevil (<i>Bruchus pisorum</i>) in <i>Pisum</i> germplasm. <i>Journal of Pest Science</i> , 2018, 91, 505-514.	3.7	22
209	Pisatin involvement in the variation of inhibition of <i>Fusarium oxysporum</i> f. sp. <i>pisi</i> spore germination by root exudates of <i>Pisum</i> spp. germplasm. <i>Plant Pathology</i> , 2018, 67, 1046-1054.	2.4	22
210	Resistance to <i>Septoria tritici</i> in <i>Hordeum chilense</i> x <i>Triticum</i> spp. Amphiploids. <i>Plant Breeding</i> , 1992, 109, 281-286.	1.9	21
211	Resistance reaction to powdery mildew (<i>Erysiphe pisi</i>) in a germplasm collection of <i>Lathyrus cicera</i> from Iberian origin. <i>Genetic Resources and Crop Evolution</i> , 2007, 54, 1517-1521.	1.6	21
212	Characterisation of resistance to crenate broomrape (<i>Orobancha crenata</i> Forsk.) in <i>Lathyrus cicera</i> L.. <i>Euphytica</i> , 2010, 173, 77-84.	1.2	21
213	Parasitic plant infection is partially controlled through symbiotic pathways. <i>Weed Research</i> , 2010, 50, 76-82.	1.7	21
214	Phylogenetic Analysis of <i>Uromyces</i> Species Infecting Grain and Forage Legumes by Sequence analysis of Nuclear Ribosomal Internal Transcribed Spacer Region. <i>Journal of Phytopathology</i> , 2011, 159, 137-145.	1.0	21
215	Multiple-disease resistance in <i>Vicia faba</i> : Multi-environment field testing for identification of combined resistance to rust and chocolate spot. <i>Field Crops Research</i> , 2011, 124, 59-65.	5.1	21
216	Characterization of Resistance Mechanisms in Faba Bean (<i>Vicia faba</i>) against Broomrape Species (<i>Orobancha</i> and <i>Phelipanche</i> spp.). <i>Frontiers in Plant Science</i> , 2016, 7, 1747.	3.6	21

#	ARTICLE	IF	CITATIONS
217	Phthalimideâ€derived strigolactone mimics as germinating agents for seeds of parasitic weeds. Pest Management Science, 2016, 72, 2069-2081.	3.4	21
218	Er3 gene, conferring resistance to powdery mildew in pea, is located in pea LGIV. Euphytica, 2018, 214, 1.	1.2	21
219	Penetration resistance to Erysiphe pisi in pea mediated by er1 gene is associated with protein cross-linking but not with callose apposition or hypersensitive response. Euphytica, 2015, 201, 381-387.	1.2	20
220	Antifeedant activity of long-chain alcohols, and fungal and plant metabolites against pea aphid (<i>Acyrtosiphon pisum</i>) as potential biocontrol strategy. Natural Product Research, 2019, 33, 2471-2479.	1.8	20
221	Faba bean. , 2021, , 452-481.		20
222	Resistance to leaf rust in cultivars of bread wheat and durum wheat grown in Spain. Plant Breeding, 2007, 126, 13-18.	1.9	19
223	Inheritance of resistance to Mycosphaerella pinodes in two wild accessions of Pisum. European Journal of Plant Pathology, 2007, 119, 53-58.	1.7	19
224	Twoâ€dimensional gel electrophoresisâ€based proteomic analysis of the <i>Medicago truncatula</i> â€rust (<i>Uromyces striatus</i>) interaction. Annals of Applied Biology, 2010, 157, 243-257.	2.5	19
225	First genetic linkage map of Lathyrus cicera based on RNA sequencing-derived markers: Key tool for genetic mapping of disease resistance. Horticulture Research, 2018, 5, 45.	6.3	19
226	Editorial: Advances in Ascochyta Research. Frontiers in Plant Science, 2018, 9, 22.	3.6	19
227	Cereal Landraces for Sustainable Agriculture. , 2011, , 147-186.		19
228	A fertile amphiploid between diploid wheat (&Triticum tauschii&) and crested wheatgrass (&Agropyron cristatum&). Genome, 1999, 42, 519-524.	2.0	19
229	Advances in disease and pest resistance in faba bean. Theoretical and Applied Genetics, 2022, 135, 3735-3756.	3.6	19
230	Targeting sources of drought tolerance within an Avena spp. collection through multivariate approaches. Planta, 2012, 236, 1529-1545.	3.2	18
231	Identification and characterization of resistance to rust (<i>Uromyces ciceris-arietini</i> (Grognot) Jacz.) Tj ETQq1 1 0.784314 rgBT /Overload	1.2	18
232	Clarification on rust species potentially infecting pea (<i>Pisum sativum</i> L.) crop and host range of <i>Uromyces pisi</i> (Pers.) Wint. Crop Protection, 2012, 37, 65-70.	2.1	18
233	Identification of preâ€and posthaustorial resistance to rust (<i>Uromyces viciae-fabae</i>) in lentil (<i>Lens culinaris</i>) germplasm. Plant Breeding, 2013, 132, 676-680.	1.9	18
234	Differences in Crenate Broomrape Parasitism Dynamics on Three Legume Crops Using a Thermal Time Model. Frontiers in Plant Science, 2016, 7, 1910.	3.6	18

#	ARTICLE	IF	CITATIONS
235	Inhibition of early development stages of rust fungi by the two fungal metabolites cyclopaldic acid and <i>epi</i> -epoformin. Pest Management Science, 2017, 73, 1161-1168.	3.4	18
236	Identification of quantitative trait loci (QTL) controlling resistance to pea weevil (<i>Bruchus pisorum</i>) in a high-density integrated DArTseq SNP-based genetic map of pea. Scientific Reports, 2020, 10, 33.	3.3	18
237	Diseases and their management.. , 2007, , 497-519.		18
238	Avirulence factors corresponding to barley genes Pa3 and Pa7 which confer resistance against <i>Puccinia hordei</i> in rust fungi other than <i>P. hordei</i> . Physiological and Molecular Plant Pathology, 1994, 45, 321-331.	2.5	17
239	QTL mapping provides evidence for lack of association of the avoidance of leaf rust in <i>Hordeum chilense</i> with stomata density. Theoretical and Applied Genetics, 2003, 106, 1283-1292.	3.6	17
240	Differential Effects of Phenylalanine Ammonia Lyase, Cinnamyl Alcohol Dehydrogenase, and Energetic Metabolism Inhibition on Resistance of Appropriate Host and Nonhost Cereals to Rust Interactions. Phytopathology, 2007, 97, 1578-1583.	2.2	17
241	Transcription factor profiling leading to the identification of putative transcription factors involved in the <i>Medicago truncatula</i> - <i>Uromyces striatus</i> interaction. Theoretical and Applied Genetics, 2010, 121, 1311-1321.	3.6	17
242	Differential gene transcript accumulation in peas in response to powdery mildew (<i>Erysiphe pisi</i>) attack. Euphytica, 2014, 198, 13-28.	1.2	17
243	Resistance reaction of <i>Medicago truncatula</i> genotypes to <i>Fusarium oxysporum</i> : effect of plant age, substrate and inoculation method. Crop and Pasture Science, 2015, 66, 506.	1.5	17
244	Abnormal germling development by brown rust and powdery mildew on cer barley mutants. Hereditas, 2004, 135, 271-276.	1.4	16
245	Identification of resistance to <i>Fusarium oxysporum</i> f.sp. <i>lentis</i> in Spanish lentil germplasm. European Journal of Plant Pathology, 2015, 143, 399-405.	1.7	16
246	Free polyamine and polyamine regulation during pre-penetration and penetration resistance events in oat against crown rust (<i>Puccinia coronata</i> f. sp. <i>avenae</i>). Plant Pathology, 2016, 65, 392-401.	2.4	16
247	Legumes in sustainable agriculture. Crop and Pasture Science, 2017, 68, i.	1.5	16
248	Title is missing!. Euphytica, 2000, 115, 221-224.	1.2	15
249	<i>Uromyces Viciae-fabae</i> Haustorium Formation in Susceptible and Resistant Faba Bean Lines. European Journal of Plant Pathology, 2003, 109, 71-73.	1.7	15
250	A Fertile Amphiploid between Durum Wheat (<i>Triticum Turgidum</i>) and the \tilde{A} -Agroticum Amphiploid (<i>Agropyron cristatum</i> \tilde{A} - T. Tauschii). Hereditas, 2004, 135, 183-186.	1.4	15
251	Histological responses in <i>Hordeum chilense</i> to brown and yellow rust fungi. Plant Pathology, 1992, 41, 611-617.	2.4	15
252	Legume breeding for broomrape resistance. Czech Journal of Genetics and Plant Breeding, 2014, 50, 144-150.	0.8	15

#	ARTICLE	IF	CITATIONS
253	Legume Crops and Biotrophic Pathogen Interactions: A Continuous Cross-Talk of a Multilayered Array of Defense Mechanisms. <i>Plants</i> , 2020, 9, 1460.	3.5	15
254	Avoidance of leaf rust fungi in wild relatives of cultivated cereals. <i>Euphytica</i> , 1996, 87, 1-6.	1.2	14
255	Occurrence of <i>Didymella fabae</i> , the Teleomorph of <i>Ascochyta fabae</i> , on Faba Bean Straw in Spain. <i>Journal of Phytopathology</i> , 2002, 150, 146-148.	1.0	14
256	Registration of RIL58â€¦LC72/Cr5, a Chickpea Germplasm Line with Rust and <i>Ascochyta</i> Blight Resistance. <i>Crop Science</i> , 2006, 46, 2331-2332.	1.8	14
257	Effects of Phenylpropanoid and Energetic Metabolism Inhibition on Faba Bean Resistance Mechanisms to Rust. <i>Phytopathology</i> , 2007, 97, 60-65.	2.2	14
258	Reaction of tritordeum to <i>Fusarium culmorum</i> and <i>Septoria nodorum</i> . <i>Euphytica</i> , 1996, 88, 165-174.	1.2	13
259	Mildew-resistant mutants induced in North American two- and six-rowed malting barley cultivars. <i>Theoretical and Applied Genetics</i> , 2003, 107, 1278-1287.	3.6	13
260	Meiotic Pairing in a Trigeneric Hybrid <i>Triticum Tauschii</i> - <i>Agropyron Cristatum</i> - <i>Hordeum Chilense</i> . <i>Hereditas</i> , 1998, 129, 113-118.	1.4	13
261	Resistance to rusts (<i>Uromyces pisi</i> and <i>U. viciae-fabae</i>) in pea. <i>Czech Journal of Genetics and Plant Breeding</i> , 2014, 50, 135-143.	0.8	13
262	Assessment of field pea (<i>Pisum sativum</i> L.) grain yield, aerial biomass and flowering date stability in Mediterranean environments. <i>Crop and Pasture Science</i> , 2017, 68, 915.	1.5	13
263	Identification of tolerance to metribuzin and imazethapyr herbicides in faba bean. <i>Crop Science</i> , 2021, 61, 2593-2611.	1.8	13
264	Development of Pea Breeding Lines with Resistance to <i>Orobanche crenata</i> Derived from Pea Landraces and Wild <i>Pisum</i> spp.. <i>Agronomy</i> , 2021, 11, 36.	3.0	13
265	Role of partial resistance to <i>Puccinia hordei</i> in barley in the defence of barley to inappropriate rust fungi. <i>Physiological and Molecular Plant Pathology</i> , 1994, 45, 219-228.	2.5	12
266	Prospects for Exploitation of Disease Resistance from <i>Hordeum Chilense</i> in Cultivated Cereals. <i>Hereditas</i> , 2004, 135, 161-169.	1.4	12
267	Extent and pattern of genetic differentiation within and between European populations of <i>Phelipanche ramosa</i> revealed by amplified fragment length polymorphism analysis. <i>Weed Research</i> , 2009, 49, 48-55.	1.7	12
268	Differential response of pea (<i>Pisum sativum</i>) to <i>Orobanche crenata</i> , <i>Orobanche foetida</i> and <i>Phelipanche aegyptiaca</i> . <i>Crop Protection</i> , 2012, 31, 27-30.	2.1	12
269	Rapid and Efficient Estimation of Pea Resistance to the Soil-Borne Pathogen <i>Fusarium oxysporum</i> by Infrared Imaging. <i>Sensors</i> , 2015, 15, 3988-4000.	3.8	12
270	Labelâ€¦free quantitative proteomic analysis of tolerance to drought in <i>Pisum sativum</i> . <i>Proteomics</i> , 2016, 16, 2776-2787.	2.2	12

#	ARTICLE	IF	CITATIONS
271	Identification and multi-environment validation of resistance to rust (<i>Uromyces viciae-fabae</i>) in <i>Vicia faba</i> . <i>Crop and Pasture Science</i> , 2017, 68, 1013.	1.5	12
272	Multi-Environmental Trials Reveal Genetic Plasticity of Oat Agronomic Traits Associated With Climate Variable Changes. <i>Frontiers in Plant Science</i> , 2018, 9, 1358.	3.6	12
273	A diversity of resistance sources to <i>Fusarium oxysporum</i> f. sp. <i>pisii</i> found within grass pea germplasm. <i>Plant and Soil</i> , 2021, 463, 19-38.	3.7	12
274	Crop Diversification to Control Powdery Mildew in Pea. <i>Agronomy</i> , 2021, 11, 690.	3.0	12
275	Diseases and their management.. , 2009, , 262-281.		12
276	Short Communication Resistance to common bunt in <i>Hordeum chilense</i> X <i>Triticum</i> spp. Amphiploids. <i>Plant Breeding</i> , 1996, 115, 416-418.	1.9	11
277	Defence reactions of <i>Hordeum chilense</i> accessions to three formae speciales of cereal powdery mildew fungi. <i>Canadian Journal of Botany</i> , 2000, 78, 1561-1570.	1.1	11
278	Comparative proteomic analysis of <i>Orobanche</i> and <i>Phelipanche</i> species inferred from seed proteins. <i>Weed Research</i> , 2009, 49, 81-87.	1.7	11
279	Temperature and water stress during conditioning and incubation phase affecting <i>Orobanche crenata</i> seed germination and radicle growth. <i>Frontiers in Plant Science</i> , 2015, 6, 408.	3.6	11
280	High productivity of dry pea genotypes resistant to crenate broomrape in Mediterranean environments. <i>Agronomy for Sustainable Development</i> , 2017, 37, 1.	5.3	11
281	Editorial: Advances in Parasitic Weed Research. <i>Frontiers in Plant Science</i> , 2018, 9, 236.	3.6	11
282	Identification of potential candidate genes controlling pea aphid tolerance in a <i>Pisum fulvum</i> high-density integrated DArTseq SNP-based genetic map. <i>Pest Management Science</i> , 2020, 76, 1731-1742.	3.4	11
283	Disclosing the Nutritional Quality Diversity of Portuguese Common Beansâ€”The Missing Link for Their Effective Use in Protein Quality Breeding Programs. <i>Agronomy</i> , 2021, 11, 221.	3.0	11
284	Effects of crop mixtures on rust development on faba bean grown in Mediterranean climates. <i>Crop Protection</i> , 2021, 146, 105686.	2.1	11
285	First Report of Crenate Broomrape (<i>Orobanche crenata</i>) on Lentil (<i>Lens culinaris</i>) and Common Vetch (<i>Vicia sativa</i>) in Salamanca Province, Spain. <i>Plant Disease</i> , 2008, 92, 1368-1368.	1.4	11
286	Adaptability and Stability of Faba Bean (<i>Vicia faba</i> L.) Accessions under Diverse Environments and Herbicide Treatments. <i>Plants</i> , 2022, 11, 251.	3.5	11
287	Searching for Abiotic Tolerant and Biotic Stress Resistant Wild Lentils for Introgression Breeding Through Predictive Characterization. <i>Frontiers in Plant Science</i> , 2022, 13, 817849.	3.6	11
288	Histology of the infection of tritordeum and its parents by cereal brown rusts. <i>Plant Pathology</i> , 1993, 42, 93-99.	2.4	10

#	ARTICLE	IF	CITATIONS
289	Title is missing!. Euphytica, 1999, 109, 157-159.	1.2	10
290	Search for Partial Resistance to Leaf Rust in a Collection of Ancient Spanish Wheats. Hereditas, 2004, 135, 193-197.	1.4	10
291	Genetic diversity in two variants of <i>Orobanche gracilis</i> Sm. [var. <i>gracilis</i> and var. <i>deludens</i> (Beck) A. Pujadas] (Orobanchaceae) from different regions of Spain. Electronic Journal of Biotechnology, 2007, 10, 0-0.	2.2	10
292	Characterization of Resistance Mechanisms to Powdery Mildew (<i>Erysiphe betae</i>) in Beet (<i>Beta</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 62	2.2	10
293	Identification of resistance to rust (<i>Uromyces appendiculatus</i>) and powdery mildew (<i>Erysiphe diffusa</i>) in Portuguese common bean germplasm. Plant Breeding, 2013, 132, 654-657.	1.9	10
294	Resistance to rust and powdery mildew in <i>Lathyrus</i> crops. Czech Journal of Genetics and Plant Breeding, 2014, 50, 116-122.	0.8	10
295	Changes in polyamine profile in host and non-host oat-powdery mildew interactions. Phytochemistry Letters, 2014, 8, 207-212.	1.2	10
296	Transcriptional profiling of <i>Medicago truncatula</i> during <i>Erysiphe pisi</i> infection. Frontiers in Plant Science, 2015, 6, 517.	3.6	10
297	Identification and characterisation of antixenosis and antibiosis to pea aphid (<i>Acyrtosiphon</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 2.5 10	2.5	10
298	Resistance to Anthracnose (<i>Colletotrichum lentis</i> , Race 0) in <i>Lens</i> spp. Germplasm. Agronomy, 2020, 10, 1799.	3.0	10
299	Partial Resistance Against <i>Erysiphe pisi</i> and <i>E. trifolii</i> Under Different Genetic Control in <i>Lathyrus cicera</i> : Outcomes from a Linkage Mapping Approach. Plant Disease, 2020, 104, 2875-2884.	1.4	10
300	Shared and tailored common bean transcriptomic responses to combined fusarium wilt and water deficit. Horticulture Research, 2021, 8, 149.	6.3	10
301	Advances in pea breeding. Burleigh Dodds Series in Agricultural Science, 2019, , 575-606.	0.2	10
302	Genomic regions associated with herbicide tolerance in a worldwide faba bean (<i>Vicia faba</i> L.) collection. Scientific Reports, 2022, 12, 158.	3.3	10
303	Grass Pea (<i>Lathyrus sativus</i> L.)-A Sustainable and Resilient Answer to Climate Challenges. Agronomy, 2022, 12, 1324.	3.0	10
304	A Fertile Amphiploid between a Wild Barley (<i>Hordeum chilense</i>) and Crested Wheatgrass (<i>Agropyron cristatum</i>). International Journal of Plant Sciences, 1999, 160, 783-786.	1.3	9
305	Impact of fungal and plant metabolites application on early development stages of pea powdery mildew. Pest Management Science, 2019, 75, 2464-2473.	3.4	9
306	Adaptation of Grass Pea (<i>Lathyrus sativus</i>) to Mediterranean Environments. Agronomy, 2020, 10, 1295.	3.0	9

#	ARTICLE	IF	CITATIONS
307	Pea Breeding Lines Adapted to Autumn Sowings in Broomrape Prone Mediterranean Environments. <i>Agronomy</i> , 2021, 11, 769.	3.0	9
308	Metabolomics profile responses to changing environments in a common bean (<i>Phaseolus vulgaris</i> L.) germplasm collection. <i>Food Chemistry</i> , 2022, 370, 131003.	8.2	9
309	Genomic Constitution and Expression of Disease Resistance in <i>Agropyron cristatum</i> * Durum Wheat Derivatives. <i>Breeding Science</i> , 2007, 57, 17-21.	1.9	9
310	Title is missing!. <i>Euphytica</i> , 2001, 122, 369-372.	1.2	8
311	Effect of Host Plant Resistance on Haustorium Formation in Cereal Rust Fungi. <i>Journal of Phytopathology</i> , 2004, 152, 381-382.	1.0	8
312	Partial Resistance to Leaf Rust in a Collection of Ancient Spanish Barleys. <i>Hereditas</i> , 2004, 135, 199-203.	1.4	8
313	Identification of a new pathotype of <i>Puccinia hordei</i> with virulence for the resistance gene Rph7. <i>European Journal of Plant Pathology</i> , 2006, 116, 103-106.	1.7	8
314	Plant Defense Responses in <i>Medicago truncatula</i> Unveiled by Microarray Analysis. <i>Plant Molecular Biology Reporter</i> , 2015, 33, 569-583.	1.8	8
315	Editorial: Advances in Legume Research. <i>Frontiers in Plant Science</i> , 2018, 9, 501.	3.6	8
316	Flower and Pod Source Influence on Pea Weevil (<i>Bruchus pisorum</i>) Oviposition Capacity and Preference. <i>Frontiers in Plant Science</i> , 2019, 10, 491.	3.6	8
317	Genetic diversity and structure of <i>Fusarium oxysporum</i> f.sp. <i>lentis</i> isolates from Iran, Syria and Algeria. <i>European Journal of Plant Pathology</i> , 2019, 153, 1019-1029.	1.7	8
318	Broomrape Threat to Agriculture. <i>Outlooks on Pest Management</i> , 2020, 31, 141-145.	0.2	8
319	The <i>MLO1</i> powdery mildew susceptibility gene in <i>Lathyrus</i> species: The power of high-density linkage maps in comparative mapping and syntenic analysis. <i>Plant Genome</i> , 2021, 14, e20090.	2.8	8
320	Inhibition of Spore Germination and Appressorium Formation of Rust Species by Plant and Fungal Metabolites. <i>Natural Product Communications</i> , 2016, 11, 1343-1347.	0.5	8
321	Hybrids Between <i>Hordeum vulgare</i> and Tetra-, Hexa-, and Octoploid Tritordeums (Amphiploid H.) Tj ETQq1 1 0.784314 rgBT / Overlock 1	1.4	7
322	Identification and characterisation of resistance against rust (<i>Puccinia allii</i>) in garlic (<i>Allium</i> sp.) germplasm. <i>Annals of Applied Biology</i> , 2011, 159, 93-98.	2.5	7
323	Unveiling common responses of <i>Medicago truncatula</i> to appropriate and inappropriate rust species. <i>Frontiers in Plant Science</i> , 2014, 5, 618.	3.6	7
324	Detection of partial resistance quantitative trait loci against <i>Didymella pinodes</i> in <i>Medicago truncatula</i> . <i>Molecular Breeding</i> , 2014, 33, 589-599.	2.1	7

#	ARTICLE	IF	CITATIONS
325	Characterization of Transcription Factors Following Expression Profiling of <i>Medicago truncatula</i> – <i>Botrytis</i> spp. Interactions. <i>Plant Molecular Biology Reporter</i> , 2014, 32, 1030-1040.	1.8	7
326	Cytoskeleton reorganization/disorganization is a key feature of induced inaccessibility for defence to successive pathogen attacks. <i>Molecular Plant Pathology</i> , 2017, 18, 662-671.	4.2	7
327	Application of Crop Growth Models to Assist Breeding for Intercropping: Opportunities and Challenges. <i>Frontiers in Plant Science</i> , 2022, 13, 720486.	3.6	7
328	Association Mapping of <i>Lathyrus sativus</i> Disease Response to <i>Uromyces pisi</i> Reveals Novel Loci Underlying Partial Resistance. <i>Frontiers in Plant Science</i> , 2022, 13, 842545.	3.6	7
329	Agronomic Performance of Broomrape Resistant and Susceptible Faba Bean Accession. <i>Agronomy</i> , 2022, 12, 1421.	3.0	7
330	Determinate Faba Bean Young Pod Response to Glyphosate and Crenate Broomrape (<i>Orobanche</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50	0.9	6
331	Search for partial resistance against <i>Puccinia hordei</i> in barley landraces from the Fertile Crescent. <i>Plant Breeding</i> , 2006, 125, 343-346.	1.9	6
332	Las leguminosas grano en la agricultura española y europea. <i>Arbor</i> , 2016, 192, a311.	0.3	6
333	Allelopathy for Parasitic Plant Management. <i>Natural Product Communications</i> , 2018, 13, 1934578X1801300.	0.5	6
334	Broomrape as a Major Constraint for Grass Pea (<i>Lathyrus sativus</i>) Production in Mediterranean Rain-Fed Environments. <i>Agronomy</i> , 2020, 10, 1931.	3.0	6
335	Heat Waves and Broomrape Are the Major Constraints for Lentil Cultivation in Southern Spain. <i>Agronomy</i> , 2021, 11, 1871.	3.0	6
336	Screening for Resistance to Leaf Rust (<i>Puccinia hordei</i>) in a Collection of Spanish Barleys. <i>Breeding Science</i> , 2006, 56, 173-177.	1.9	6
337	The contribution of <i>Hordeum chilense</i> to partial resistance of tritordeum to wheat brown rust. <i>Euphytica</i> , 1992, 59, 129-133.	1.2	5
338	Identification and expression analysis of a MYB family transcription factor in the parasitic plant <i>Orobanche ramosa</i> . <i>Annals of Applied Biology</i> , 2007, 150, 123-130.	2.5	5
339	Resistance of <i>Hordeum chilense</i> against loose smuts of wheat and barley (<i>Ustilago tritici</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 5	1.9	5
340	Grass Pea. <i>Handbook of Plant Breeding</i> , 2015, , 251-265.	0.1	5
341	Inhibition of Spore Germination and Appressorium Formation of Rust Species by Plant and Fungal Metabolites. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	5
342	Grass pea and pea phylogenetic relatedness reflected at <i>Fusarium oxysporum</i> host range. <i>Crop Protection</i> , 2021, 141, 105495.	2.1	5

#	ARTICLE	IF	CITATIONS
343	First Report of Crenate Broomrape (<i>Orobancha crenata</i>) on White Lupine (<i>Lupinus albus</i>) Growing in Alkaline Soils in Spain and Egypt. Plant Disease, 2009, 93, 970-970.	1.4	5
344	The legume manifesto: (Net)workers on Fabaceae, unite!. Ratarstvo I Povrtarstvo, 2011, 48, 253-258.	0.5	5
345	Grass pea natural variation reveals oligogenic resistance to <i>Fusarium oxysporum</i> f. sp. <i>pisi</i> . Plant Genome, 2021, 14, e20154.	2.8	5
346	Portuguese Common Bean Natural Variation Helps to Clarify the Genetic Architecture of the Legume's Nutritional Composition and Protein Quality. Plants, 2022, 11, 26.	3.5	5
347	Anthraquinones and their analogues as potential biocontrol agents of rust and powdery mildew diseases of field crops. Pest Management Science, 2022, , .	3.4	5
348	Response of Vicia species to Ascochyta fabae and Uromyces viciae-fabae. Czech Journal of Genetics and Plant Breeding, 2014, 50, 109-115.	0.8	4
349	Compromised Photosynthetic Electron Flow and H ₂ O ₂ Generation Correlate with Genotype-Specific Stomatal Dysfunctions during Resistance against Powdery Mildew in Oats. Frontiers in Plant Science, 2016, 7, 1660.	3.6	4
350	Characterization of the Resistance to Powdery Mildew and Leaf Rust Carried by the Bread Wheat Cultivar Victo. International Journal of Molecular Sciences, 2021, 22, 3109.	4.1	4
351	Inheritance of resistance to Mycosphaerella pinodes in two wild accessions of Pisum. , 2007, , 53-58.		4
352	Development of Quantitative Real-Time PCR Assays to Quantify Erysiphe pisi and Erysiphe trifolii and Its Implementation for Monitoring Their Relative Prevalence in Pea Crops in Spain and Tunisia. Agronomy, 2022, 12, 334.	3.0	4
353	Evaluation of performance and stability of new sources for tolerance to post-emergence herbicides		

#	ARTICLE	IF	CITATIONS
361	Abiotic and Biotic Stresses Interaction in Fabaceae Plants. Contributions from the Grain Legumes/Soilborne Vascular Diseases/Drought Stress Triangle. , 2020, , 237-260.		2
362	Advances in understanding plant root response to weedy root parasites. Burleigh Dodds Series in Agricultural Science, 2021, , 215-230.	0.2	1
363	Identification of resistance to tan spot (<i>Pyrenophora tritici-repentis</i>) in <i>Hordeum chilense</i> and its expression in its amphiploids with diploid, tetraploid and hexaploid <i>Triticum</i> species. Plant Breeding, 2012, 131, 579-583.	1.9	0
364	Potential of Tritordeum as a New Cereal Crop. , 1994, , 443-445.		0
365	Developing pest- and disease-resistant cultivars of grain legumes. Burleigh Dodds Series in Agricultural Science, 2018, , 155-176.	0.2	0
366	Aleksandar Mikić, the legume (re)searcher. , 0, , .		0