

# Fotios Loupakis

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2227424/publications.pdf>

Version: 2024-02-01

352  
papers

16,972  
citations

20797

60  
h-index

17580

121  
g-index

361  
all docs

361  
docs citations

361  
times ranked

19752  
citing authors

#	ARTICLE	IF	CITATIONS
1	Epstein-Barr virus associated gastric dysplasia: a new rare entity?. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2022, 480, 939-944.	1.4	3
2	Complete pathological response of colorectal peritoneal metastases in Lynch syndrome after immunotherapy case report: is a paradigm shift in cytoreductive surgery needed?. <i>BMC Gastroenterology</i> , 2022, 22, 17.	0.8	4
3	Systematic review of randomised clinical trials and observational studies for patients with RAS wild-type or BRAF-mutant metastatic and/or unresectable colorectal cancer. <i>Critical Reviews in Oncology/Hematology</i> , 2022, 173, 103646.	2.0	3
4	Prognostic impact of FGFR2/3 alterations in patients with biliary tract cancers receiving systemic chemotherapy: the BITCOIN study. <i>European Journal of Cancer</i> , 2022, 166, 165-175.	1.3	17
5	Genetic variants involved in the cGAS-STING pathway predict outcome in patients with metastatic colorectal cancer: Data from FIRE-3 and TRIBE trials. <i>European Journal of Cancer</i> , 2022, 172, 22-30.	1.3	3
6	Outcome of patients with colorectal cancer undergoing lung metastases resection: a single-institution retrospective analysis. <i>Tumori</i> , 2021, 107, 46-54.	0.6	2
7	Hurrying up but not rushing, acting and not reacting, good sense and not common sense: Open thoughts and reasonable doubts on COVID-19 vaccination strategies in cancer patients. <i>Critical Reviews in Oncology/Hematology</i> , 2021, 167, 103271.	2.0	1
8	Encorafenib Plus Cetuximab as a New Standard of Care for Previously Treated BRAF V600E-Mutant Metastatic Colorectal Cancer: Updated Survival Results and Subgroup Analyses from the BEACON Study. <i>Journal of Clinical Oncology</i> , 2021, 39, 273-284.	0.8	254
9	Clinical significance of enterocyte-specific gene polymorphisms as candidate markers of oxaliplatin-based treatment for metastatic colorectal cancer. <i>Pharmacogenomics Journal</i> , 2021, 21, 285-295.	0.9	3
10	Synaptophysin expression in mutated advanced colorectal cancers identifies a new subgroup of tumours with worse prognosis. <i>European Journal of Cancer</i> , 2021, 146, 145-154.	1.3	8
11	RNA-Binding Protein Polymorphisms as Novel Biomarkers to Predict Outcomes of Metastatic Colorectal Cancer: A Meta-analysis from TRIBE, FIRE-3, and MAVERICC. <i>Molecular Cancer Therapeutics</i> , 2021, 20, 1153-1160.	1.9	1
12	NUTM1-rearranged colorectal sarcoma: a clinicopathologically and genetically distinctive malignant neoplasm with a poor prognosis. <i>Modern Pathology</i> , 2021, 34, 1547-1557.	2.9	24
13	RAS as a positive predictive biomarker: focus on lung and colorectal cancer patients. <i>European Journal of Cancer</i> , 2021, 146, 74-83.	1.3	29
14	MicroRNAs as Predictive Biomarkers of Resistance to Targeted Therapies in Gastrointestinal Tumors. <i>Biomedicines</i> , 2021, 9, 318.	1.4	7
15	Molecular profiling of appendiceal serrated lesions, polyps and mucinous neoplasms: a single-centre experience. <i>Journal of Cancer Research and Clinical Oncology</i> , 2021, 147, 1897-1904.	1.2	7
16	The Role of p53 Expression in Patients with RAS/BRAF Wild-Type Metastatic Colorectal Cancer Receiving Irinotecan and Cetuximab as Later Line Treatment. <i>Targeted Oncology</i> , 2021, 16, 517-527.	1.7	7
17	Trastuzumab deruxtecan (DS-8201) in patients with HER2-expressing metastatic colorectal cancer (DESTINY-CRC01): a multicentre, open-label, phase 2 trial. <i>Lancet Oncology</i> , The, 2021, 22, 779-789.	5.1	234
18	Random survival forests identify pathways with polymorphisms predictive of survival in KRAS mutant and KRAS wild-type metastatic colorectal cancer patients. <i>Scientific Reports</i> , 2021, 11, 12191.	1.6	3

#	ARTICLE	IF	CITATIONS
19	Germ line polymorphisms of genes involved in pluripotency transcription factors predict efficacy of cetuximab in metastatic colorectal cancer. <i>European Journal of Cancer</i> , 2021, 150, 133-142.	1.3	1
20	Detection of Molecular Residual Disease Using Personalized Circulating Tumor DNA Assay in Patients With Colorectal Cancer Undergoing Resection of Metastases. <i>JCO Precision Oncology</i> , 2021, 5, 1166-1177.	1.5	55
21	A Real-World Application of Liquid Biopsy in Metastatic Colorectal Cancer: The Poseidon Study. <i>Cancers</i> , 2021, 13, 5128.	1.7	6
22	Association of CLDN18 Protein Expression with Clinicopathological Features and Prognosis in Advanced Gastric and Gastroesophageal Junction Adenocarcinomas. <i>Journal of Personalized Medicine</i> , 2021, 11, 1095.	1.1	42
23	PD-L1 expression in gastroesophageal dysplastic lesions. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2020, 477, 151-156.	1.4	24
24	Polymorphisms within Immune Regulatory Pathways Predict Cetuximab Efficacy and Survival in Metastatic Colorectal Cancer Patients. <i>Cancers</i> , 2020, 12, 2947.	1.7	4
25	A polymorphism in the cachexia-associated gene INHBA predicts efficacy of regorafenib in patients with refractory metastatic colorectal cancer. <i>PLoS ONE</i> , 2020, 15, e0239439.	1.1	5
26	Practical considerations in the use of regorafenib in metastatic colorectal cancer. <i>Therapeutic Advances in Medical Oncology</i> , 2020, 12, 175883592095686.	1.4	16
27	Immunogenic cell death pathway polymorphisms for predicting oxaliplatin efficacy in metastatic colorectal cancer. , 2020, 8, e001714.		23
28	TP53 Mutation Analysis in Gastric Cancer and Clinical Outcomes of Patients with Metastatic Disease Treated with Ramucirumab/Paclitaxel or Standard Chemotherapy. <i>Cancers</i> , 2020, 12, 2049.	1.7	11
29	Impact of Pre-Analytical Factors on MSI Test Accuracy in Mucinous Colorectal Adenocarcinoma: A Multi-Assay Concordance Study. <i>Cells</i> , 2020, 9, 2019.	1.8	30
30	Efficacy and Safety of Immune Checkpoint Inhibitors in Patients with Microsatellite Instability-High End-Stage Cancers and Poor Performance Status Related to High Disease Burden. <i>Oncologist</i> , 2020, 25, 803-809.	1.9	26
31	The Pan-Immune-Inflammation Value is a new prognostic biomarker in metastatic colorectal cancer: results from a pooled-analysis of the Valentino and TRIBE first-line trials. <i>British Journal of Cancer</i> , 2020, 123, 403-409.	2.9	93
32	Glycolytic competence in gastric adenocarcinomas negatively impacts survival outcomes of patients treated with salvage paclitaxel-ramucirumab. <i>Gastric Cancer</i> , 2020, 23, 1064-1074.	2.7	5
33	Prognostic impact of immune-microenvironment in colorectal liver metastases resected after triplets plus a biologic agent: A pooled analysis of five prospective trials. <i>European Journal of Cancer</i> , 2020, 135, 78-88.	1.3	10
34	Combination of variations in inflammation- and endoplasmic reticulum-associated genes as putative biomarker for bevacizumab response in KRAS wild-type colorectal cancer. <i>Scientific Reports</i> , 2020, 10, 9778.	1.6	5
35	KRAS G12C Metastatic Colorectal Cancer: Specific Features of a New Emerging Target Population. <i>Clinical Colorectal Cancer</i> , 2020, 19, 219-225.	1.0	45
36	Upfront FOLFOXIRI plus bevacizumab and reintroduction after progression versus mFOLFOX6 plus bevacizumab followed by FOLFIRI plus bevacizumab in the treatment of patients with metastatic colorectal cancer (TRIBE2): a multicentre, open-label, phase 3, randomised, controlled trial. <i>Lancet Oncology</i> , The, 2020, 21, 497-507.	5.1	196

#	ARTICLE	IF	CITATIONS
37	Cancer care during the spread of coronavirus disease 2019 (COVID-19) in Italy: young oncologistsâ€™ perspective. <i>ESMO Open</i> , 2020, 5, e000759.	2.0	161
38	Single Nucleotide Polymorphisms in MiRNA Binding Sites of Nucleotide Excision Repair-Related Genes Predict Clinical Benefit of Oxaliplatin in FOLFOXIRI Plus Bevacizumab: Analysis of the TRIBE Trial. <i>Cancers</i> , 2020, 12, 1742.	1.7	4
39	Prediction of Benefit from Checkpoint Inhibitors in Mismatch Repair Deficient Metastatic Colorectal Cancer: Role of Tumor Infiltrating Lymphocytes. <i>Oncologist</i> , 2020, 25, 481-487.	1.9	77
40	The heterogeneous clinical and pathological landscapes of metastatic Braf-mutated colorectal cancer. <i>Cancer Cell International</i> , 2020, 20, 30.	1.8	63
41	A polymorphism within the R-spondin 2 gene predicts outcome in metastatic colorectal cancer patients treated with FOLFIRI/bevacizumab: data from FIRE-3 and TRIBE trials. <i>European Journal of Cancer</i> , 2020, 131, 89-97.	1.3	9
42	Investigating the concordance in molecular subtypes of primary colorectal tumors and their matched synchronous liver metastasis. <i>International Journal of Cancer</i> , 2020, 147, 2303-2315.	2.3	14
43	Retreatment With Anti-EGFR Antibodies in Metastatic Colorectal Cancer Patients: A Multi-institutional Analysis. <i>Clinical Colorectal Cancer</i> , 2020, 19, 191-199.e6.	1.0	20
44	Thyroid hormones ratio is a major prognostic marker in advanced metastatic colorectal cancer: Results from the phase III randomised CORRECT trial. <i>European Journal of Cancer</i> , 2020, 133, 66-73.	1.3	19
45	Encorafenib plus cetuximab with or without binimetinib for BRAF V600E metastatic colorectal cancer: Updated survival results from a randomized, three-arm, phase III study versus choice of either irinotecan or FOLFIRI plus cetuximab (BEACON CRC). <i>Journal of Clinical Oncology</i> , 2020, 38, 4001-4001.	0.8	35
46	Novel Common Genetic Susceptibility Loci for Colorectal Cancer. <i>Journal of the National Cancer Institute</i> , 2019, 111, 146-157.	3.0	129
47	Tumor mutation burden: from comprehensive mutational screening to the clinic. <i>Cancer Cell International</i> , 2019, 19, 209.	1.8	116
48	Pathological Tumor Regression Grade Classifications in Gastrointestinal Cancers: Role on Patientsâ€™ Prognosis. <i>International Journal of Surgical Pathology</i> , 2019, 27, 816-835.	0.4	8
49	A validated prognostic classifier for BRAF-mutated metastatic colorectal cancer: the BRAF BeCool study. <i>European Journal of Cancer</i> , 2019, 118, 121-130.	1.3	51
50	Aryl hydrocarbon receptor nuclear translocator-like (ARNTL/BMAL1) is associated with bevacizumab resistance in colorectal cancer via regulation of vascular endothelial growth factor A. <i>EBioMedicine</i> , 2019, 45, 139-154.	2.7	36
51	High Circulating Methylated DNA Is a Negative Predictive and Prognostic Marker in Metastatic Colorectal Cancer Patients Treated With Regorafenib. <i>Frontiers in Oncology</i> , 2019, 9, 622.	1.3	22
52	Correlation between p53 expression and clinical outcome in RAS/BRAF wild type metastatic colorectal cancer patients receiving later-line irinotecan-cetuximab. <i>Annals of Oncology</i> , 2019, 30, v226.	0.6	0
53	Genetic variants in the one-carbon metabolism pathway to predict outcome in patients with metastatic colorectal cancer (mCRC): Data from TRIBE and FIRE-3 phase III trials. <i>Annals of Oncology</i> , 2019, 30, v763-v764.	0.6	0
54	Treatment with checkpoint inhibitors in a metastatic colorectal cancer patient with molecular and immunohistochemical heterogeneity in MSI/dMMR status. <i>Journal of Clinical Oncology</i> , 2019, 37, 297.		24

#	ARTICLE	IF	CITATIONS
55	PTEN in Colorectal Cancer: Shedding Light on Its Role as Predictor and Target. <i>Cancers</i> , 2019, 11, 1765.	1.7	54
56	CK7 and consensus molecular subtypes as major prognosticators in V600EBRAF mutated metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2019, 121, 593-599.	2.9	24
57	Negative Hyperselection of Patients With <i>RAS</i> and <i>BRAF</i> Wild-Type Metastatic Colorectal Cancer Who Received Panitumumab-Based Maintenance Therapy. <i>Journal of Clinical Oncology</i> , 2019, 37, 3099-3110.	0.8	65
58	Relationship Between Tumor Response and Tumor-Related Symptoms in <i>RAS</i> Wild-Type Metastatic Colorectal Cancer: Retrospective Analyses From 3 Panitumumab Trials. <i>Clinical Colorectal Cancer</i> , 2019, 18, 245-256.e5.	1.0	2
59	Encorafenib, Binimetinib, and Cetuximab in <i>BRAF</i> V600E-Mutated Colorectal Cancer. <i>New England Journal of Medicine</i> , 2019, 381, 1632-1643.	13.9	918
60	Early modifications of circulating microRNAs levels in metastatic colorectal cancer patients treated with regorafenib. <i>Pharmacogenomics Journal</i> , 2019, 19, 455-464.	0.9	5
61	Shared heritability and functional enrichment across six solid cancers. <i>Nature Communications</i> , 2019, 10, 431.	5.8	88
62	An overview on clinical, pathological and molecular features of lung metastases from colorectal cancer. <i>Expert Review of Respiratory Medicine</i> , 2019, 13, 635-644.	1.0	7
63	Quantitative evidence for early metastatic seeding in colorectal cancer. <i>Nature Genetics</i> , 2019, 51, 1113-1122.	9.4	315
64	Claudin-18 expression in oesophagogastric adenocarcinomas: a tissue microarray study of 523 molecularly profiled cases. <i>British Journal of Cancer</i> , 2019, 121, 257-263.	2.9	53
65	Impact of polymorphisms within genes involved in regulating DNA methylation in patients with metastatic colorectal cancer enrolled in three independent, randomised, open-label clinical trials: a meta-analysis from TRIBE, MAVERICC and FIRE-3. <i>European Journal of Cancer</i> , 2019, 111, 138-147.	1.3	4
66	AMPK variant, a candidate of novel predictor for chemotherapy in metastatic colorectal cancer: A meta-analysis using TRIBE, MAVERICC and FIRE3. <i>International Journal of Cancer</i> , 2019, 145, 2082-2090.	2.3	4
67	Class 1, 2, and 3 <i>BRAF</i> -Mutated Metastatic Colorectal Cancer: A Detailed Clinical, Pathologic, and Molecular Characterization. <i>Clinical Cancer Research</i> , 2019, 25, 3954-3961.	3.2	67
68	Benefit from anti-EGFRs in <i>RAS</i> and <i>BRAF</i> wild-type metastatic transverse colon cancer: a clinical and molecular proof of concept study. <i>ESMO Open</i> , 2019, 4, e000489.	2.0	14
69	Ramucirumab: the long and winding road toward being an option for mCRC treatment. <i>Expert Opinion on Biological Therapy</i> , 2019, 19, 399-409.	1.4	6
70	Chemotherapeutic and antiangiogenic drugs beyond tumor progression in colon cancer: Evaluation of the effects of switched schedules and related pharmacodynamics. <i>Biochemical Pharmacology</i> , 2019, 164, 94-105.	2.0	14
71	DPYD*6 plays an important role in fluoropyrimidine toxicity in addition to DPYD*2A and c.2846A>T: a comprehensive analysis in 1254 patients. <i>Pharmacogenomics Journal</i> , 2019, 19, 556-563.	0.9	35
72	Another Chapter of the Right Versus Left Story : Is Primary Tumor Location a Prognostic Feature in <i>RAS</i> Mutant Metastatic Colorectal Cancer?. <i>Oncologist</i> , 2019, 24, e77-e79.	1.9	3

#	ARTICLE	IF	CITATIONS
73	Prognostic Effect of Adenosine-related Genetic Variants in Metastatic Colorectal Cancer Treated With Bevacizumab-based Chemotherapy. <i>Clinical Colorectal Cancer</i> , 2019, 18, e8-e19.	1.0	12
74	miR-224 Is Significantly Upregulated and Targets Caspase-3 and Caspase-7 During Colorectal Carcinogenesis. <i>Translational Oncology</i> , 2019, 12, 282-291.	1.7	14
75	Safety and Tolerability of Anti-Angiogenic Protein Kinase Inhibitors and Vascular-Disrupting Agents in Cancer: Focus on Gastrointestinal Malignancies. <i>Drug Safety</i> , 2019, 42, 159-179.	1.4	18
76	Safety and Tolerability of c-MET Inhibitors in Cancer. <i>Drug Safety</i> , 2019, 42, 211-233.	1.4	76
77	BRCA1 genetic variant to predict survival in metastatic colorectal cancer (mCRC) patients (pts) treated with FOLFIRI/bevacizumab (bev): Results from phase III TRIBE and FIRE-3 trials.. <i>Journal of Clinical Oncology</i> , 2019, 37, 3145-3145.	0.8	3
78	Targeted therapies in metastatic gastric cancer: Current knowledge and future perspectives. <i>World Journal of Gastroenterology</i> , 2019, 25, 5773-5788.	1.4	69
79	Th17 cell pathway-related genetic variants in metastatic colorectal cancer: A meta-analysis using TRIBE, MAVERICC, and FIRE-3.. <i>Journal of Clinical Oncology</i> , 2019, 37, 594-594.	0.8	0
80	Polymorphisms in the telomerase complex to predict outcome in patients (pts) with metastatic colorectal cancer (mCRC): Data from TRIBE and FIRE-3 phase III trials.. <i>Journal of Clinical Oncology</i> , 2019, 37, 566-566.	0.8	0
81	Genetic variations within the CD40L immune stimulating gene predict outcome for mCRC patients treated with first-line FOLFIRI/bevacizumab: Data from FIRE-3 and TRIBE.. <i>Journal of Clinical Oncology</i> , 2019, 37, 558-558.	0.8	4
82	Genetic variants in RNA binding protein (RBP) to predict outcome in metastatic colorectal cancer (mCRC): Data from FIRE-3, TRIBE, and MAVERICC trials.. <i>Journal of Clinical Oncology</i> , 2019, 37, 3545-3545.	0.8	0
83	Polymorphisms in the dopamine (DA) signaling to predict outcome in patients (pts) with metastatic colorectal cancer (mCRC): Data from TRIBE, MAVERICC, and FIRE-3 phase III trials.. <i>Journal of Clinical Oncology</i> , 2019, 37, 3048-3048.	0.8	1
84	Abstract 1342: Polymorphisms in genes involved in mitophagy pathway predict clinical outcome in patients (pts) with metastatic colorectal cancer (mCRC): Data from TRIBE and FIRE3 phase III trials. <i>Cancer Research</i> , 2019, 79, 1342-1342.	0.4	1
85	Pharmacokinetic analysis of metronomic capecitabine in refractory metastatic colorectal cancer patients. <i>Investigational New Drugs</i> , 2018, 36, 709-714.	1.2	8
86	Primary tumor sidedness and benefit from FOLFOXIRI plus bevacizumab as initial therapy for metastatic colorectal cancer. Retrospective analysis of the TRIBE trial by GONO. <i>Annals of Oncology</i> , 2018, 29, 1528-1534.	0.6	83
87	Prognostic Value of ACVRL1 Expression in Metastatic Colorectal Cancer Patients Receiving First-line Chemotherapy With Bevacizumab: Results From the Triplet Plus Bevacizumab (TRIBE) Study. <i>Clinical Colorectal Cancer</i> , 2018, 17, e471-e488.	1.0	12
88	The role of tumor angiogenesis as a therapeutic target in colorectal cancer. <i>Expert Review of Anticancer Therapy</i> , 2018, 18, 251-266.	1.1	41
89	Gene Polymorphisms in the CCL5/CCR5 Pathway as a Genetic Biomarker for Outcome and Handâ€™Foot Skin Reaction in Metastatic Colorectal Cancer Patients Treated With Regorafenib. <i>Clinical Colorectal Cancer</i> , 2018, 17, e395-e414.	1.0	25
90	A Polymorphism within the Vitamin D Transporter Gene Predicts Outcome in Metastatic Colorectal Cancer Patients Treated with FOLFIRI/Bevacizumab or FOLFIRI/Cetuximab. <i>Clinical Cancer Research</i> , 2018, 24, 784-793.	3.2	23

#	ARTICLE	IF	CITATIONS
91	Loss of Chromosome 18q11.2-q12.1 Is Predictive for Survival in Patients With Metastatic Colorectal Cancer Treated With Bevacizumab. <i>Journal of Clinical Oncology</i> , 2018, 36, 2052-2060.	0.8	26
92	The DISTINCTIVE study: A biologically enriched phase II study of second-line folfiri/afllbercept in proSpecTively stratified, anti-EGFR resistaNt, metastatic coloreCTal cancer patlents with RAS Validated wild typE status - Trial in progress. <i>Annals of Oncology</i> , 2018, 29, v82.	0.6	3
93	Impact of primary tumour location on efficacy of bevacizumab plus chemotherapy in metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2018, 119, 1451-1455.	2.9	19
94	Copy number load predicts outcome of metastatic colorectal cancer patients receiving bevacizumab combination therapy. <i>Nature Communications</i> , 2018, 9, 4112.	5.8	55
95	LONG-NONCODING RNAs in gastroesophageal cancers. <i>Non-coding RNA Research</i> , 2018, 3, 195-212.	2.4	39
96	Assessment of intratumor immune-microenvironment in colorectal cancers with extranodal extension of nodal metastases. <i>Cancer Cell International</i> , 2018, 18, 131.	1.8	7
97	Precision medicine in cholangiocarcinoma. <i>Translational Gastroenterology and Hepatology</i> , 2018, 3, 40-40.	1.5	61
98	Potential role of PIN1 genotypes in predicting benefit from oxaliplatin-based and irinotecan-based treatment in patients with metastatic colorectal cancer. <i>Pharmacogenomics Journal</i> , 2018, 18, 623-632.	0.9	8
99	Cancer Immunotherapy and Identification of Prognostic and Predictive Biomarkers. <i>BioMed Research International</i> , 2018, 2018, 1-2.	0.9	3
100	First-line FOLFOX plus panitumumab followed by 5-FU/LV plus panitumumab or single-agent panitumumab as maintenance therapy in patients with RAS wild-type metastatic colorectal cancer (mCRC): The VALENTINO study. <i>Annals of Oncology</i> , 2018, 29, v106.	0.6	0
101	The PANDA study: a randomized phase II study of first-line FOLFOX plus panitumumab versus 5FU plus panitumumab in RAS and BRAF wild-type elderly metastatic colorectal cancer patients. <i>BMC Cancer</i> , 2018, 18, 98.	1.1	17
102	Noninferiority of three monthsversussix months of oxaliplatinâ€based adjuvant chemotherapy for resected colon cancer. How shouldIDEAfindings affect clinical practice?. <i>International Journal of Cancer</i> , 2018, 143, 2342-2350.	2.3	7
103	Prognostic Value of Thyroid Hormone Ratios in Patients With Advanced Metastatic Colorectal Cancer Treated With Regorafenib: TheÂTOREADOR Study. <i>Clinical Colorectal Cancer</i> , 2018, 17, e601-e615.	1.0	18
104	NOS2 polymorphisms in prediction of benefit from first-line chemotherapy in metastatic colorectal cancer patients. <i>PLoS ONE</i> , 2018, 13, e0193640.	1.1	5
105	Polymorphism in the circadian clock pathway to predict outcome in patients (pts) with metastatic colorectal cancer (mCRC): Data from TRIBE and FIRE-3 phase III trials.. <i>Journal of Clinical Oncology</i> , 2018, 36, 3576-3576.	0.8	2
106	Clinico-pathological and molecular characterisation of BRAF mutant metastatic colorectal cancer (mCRC): Are all mutations created equal?. <i>Journal of Clinical Oncology</i> , 2018, 36, 3590-3590.	0.8	4
107	Clinical prognostic score of BRAF V600E mutated (BM) metastatic colorectal cancer (mCRC): Results from the â€œBRAF, BeCoolâ€platform.. <i>Journal of Clinical Oncology</i> , 2018, 36, 639-639.	0.8	2
108	Polymorphism in cancer-associated fibroblasts (CAFs) related genes and clinical outcome in metastatic colorectal cancer (mCRC) patients (pts) enrolled in two independent randomized phase III trials: TRIBE and FIRE-3.. <i>Journal of Clinical Oncology</i> , 2018, 36, 645-645.	0.8	1

#	ARTICLE	IF	CITATIONS
109	Matrix metalloproteinase-related gene polymorphisms to predict efficacy of regorafenib in patients with metastatic colorectal cancer.. Journal of Clinical Oncology, 2018, 36, 692-692.	0.8	1
110	Efficacy outcomes with bevacizumab added to chemotherapy (bev+CT) compared with chemotherapy alone (CT) in left- and right-sided tumors in metastatic colorectal cancer (mCRC).. Journal of Clinical Oncology, 2018, 36, 726-726.	0.8	4
111	<i>DPYD</i> and <i>UGT1A1</i> genotyping to predict adverse events during first-line FOLFIRI or FOLFOXIRI plus bevacizumab in metastatic colorectal cancer. Oncotarget, 2018, 9, 7859-7866.	0.8	25
112	The impact of Tfh cell/ B cell pathway-related genetic variants in metastatic colorectal cancer patients with bevacizumab-based chemotherapy.. Journal of Clinical Oncology, 2018, 36, 651-651.	0.8	0
113	Single nucleotide polymorphisms in miRNA binding sites of nucleotide excision repair-related genes to predict clinical benefit of oxaliplatin in FOLFOXIRI plus bevacizumab in TRIBE trial.. Journal of Clinical Oncology, 2018, 36, 663-663.	0.8	0
114	Polymorphisms in beta-defensin pathways and clinical outcomes in metastatic colorectal cancer patients treated with FOLFIRI-bevacizumab in two randomized phase III trials.. Journal of Clinical Oncology, 2018, 36, 662-662.	0.8	3
115	Genetic variants in methylation and demethylation pathways to predict clinical outcome in metastatic colorectal cancer (mCRC) patients (pts) treated with first-line FOLFIRI/Bev: Data from TRIBE and FIRE-3 trials.. Journal of Clinical Oncology, 2018, 36, 646-646.	0.8	0
116	Histopathologic response and growth patterns of colorectal cancer liver metastases (CRCLM) in patients treated with triplets plus bevacizumab (bev) or anti-EGFRs.. Journal of Clinical Oncology, 2018, 36, 636-636.	0.8	0
117	Clinical significance of enterocyte-specific gene polymorphisms as candidate marker of oxaliplatin-based treatment for metastatic colorectal cancer.. Journal of Clinical Oncology, 2018, 36, 12066-12066.	0.8	0
118	Genetic variants within the glucocorticoids related genes to predict outcome in patients with metastatic colorectal cancer (mCRC).. Journal of Clinical Oncology, 2018, 36, 12098-12098.	0.8	0
119	The impact of Th17 cell pathway-related genetic variants in metastatic colorectal cancer patients treated with bevacizumab-based chemotherapy.. Journal of Clinical Oncology, 2018, 36, e15578-e15578.	0.8	0
120	Genetic variation in TET3 and survival in metastatic colorectal cancer (mCRC) from FIRE-3, TRIBE, and MAVERICC clinical trials.. Journal of Clinical Oncology, 2018, 36, 3575-3575.	0.8	0
121	Abstract 1823: Identification of molecular determinants of vinorelbine resistance in BRAF(V600E) mutated chemorefractory metastatic colorectal cancer patients. , 2018, , .		0
122	Abstract 2579: Loss of chromosome 18q11.2-18q12.1 is predictive for progression-free survival in metastatic colorectal cancer patients treated with bevacizumab. , 2018, , .		0
123	Abstract 206: The Consensus Molecular Classification (CMS) of primary colorectal tumors and their matched liver metastasis: Investigating the concordance. , 2018, , .		0
124	Abstract 2614: Macrophage erythroblast attacher (MAEA) polymorphisms are associated with clinical outcome in TRIBE study mCRC patients treated with 5-fluorouracil/bevacizumab-based therapy. , 2018, , .		0
125	Glycolysis gene expression analysis and selective metabolic advantage in the clinical progression of colorectal cancer. Pharmacogenomics Journal, 2017, 17, 258-264.	0.9	79
126	Variant alleles in factor V, prothrombin, plasminogen activator inhibitor-1, methylenetetrahydrofolate reductase and risk of thromboembolism in metastatic colorectal cancer patients treated with first-line chemotherapy plus bevacizumab. Pharmacogenomics Journal, 2017, 17, 331-336.	0.9	10



#	ARTICLE	IF	CITATIONS
127	Serum LDH predicts benefit from bevacizumab beyond progression in metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2017, 116, 318-323.	2.9	29
128	The role of pharmacogenetics in the new ESMO colorectal cancer guidelines. <i>Pharmacogenomics</i> , 2017, 18, 197-200.	0.6	7
129	Genetic variants of DNA repair-related genes predict efficacy of TAS-102 in patients with refractory metastatic colorectal cancer. <i>Annals of Oncology</i> , 2017, 28, 1015-1022.	0.6	24
130	Single nucleotide polymorphisms in the IGF1R pathway are associated with outcome in mCRC patients enrolled in the FIRE-3 trial. <i>International Journal of Cancer</i> , 2017, 141, 383-392.	2.3	10
131	Immunotherapy for colorectal cancer: where are we heading?. <i>Expert Opinion on Biological Therapy</i> , 2017, 17, 709-721.	1.4	85
132	Autophagy-related polymorphisms predict hypertension in patients with metastatic colorectal cancer treated with FOLFIRI and bevacizumab: Results from TRIBE and FIRE-3 trials. <i>European Journal of Cancer</i> , 2017, 77, 13-20.	1.3	19
133	Homeobox B9 Mediates Resistance to Anti-VEGF Therapy in Colorectal Cancer Patients. <i>Clinical Cancer Research</i> , 2017, 23, 4312-4322.	3.2	41
134	Efficacy of FOLFOXIRI plus bevacizumab in liver-limited metastatic colorectal cancer: A pooled analysis of clinical studies by Gruppo Oncologico del Nord Ovest. <i>European Journal of Cancer</i> , 2017, 73, 74-84.	1.3	54
135	Potential role of polymorphisms in the transporter genes ENT1 and MATE1 / OCT2 in predicting TAS-102 efficacy and toxicity in patients with refractory metastatic colorectal cancer. <i>European Journal of Cancer</i> , 2017, 86, 197-206.	1.3	22
136	QoL is a cool tool. <i>Annals of Oncology</i> , 2017, 28, 2032-2033.	0.6	2
137	BRAF p.V600E-specific immunohistochemical assessment in colorectal cancer endoscopy biopsies is consistent with the mutational profiling. <i>Histopathology</i> , 2017, 71, 1008-1011.	1.6	8
138	Tandem repeat variation near the HIC1 (hypermethylated in cancer 1) promoter predicts outcome of oxaliplatin-based chemotherapy in patients with metastatic colorectal cancer. <i>Cancer</i> , 2017, 123, 4506-4514.	2.0	8
139	Anti-EGFR monoclonal antibody panitumumab for the treatment of patients with metastatic colorectal cancer: an overview of current practice and future perspectives. <i>Expert Opinion on Biological Therapy</i> , 2017, 17, 1297-1308.	1.4	21
140	New perspectives for TAS-102: TASK successful?. <i>Lancet Oncology</i> , The, 2017, 18, 1139-1141.	5.1	1
141	Impact of genetic variations in the MAPK signaling pathway on outcome in metastatic colorectal cancer patients treated with first-line FOLFIRI and bevacizumab: data from FIRE-3 and TRIBE trials. <i>Annals of Oncology</i> , 2017, 28, 2780-2785.	0.6	28
142	Vinorelbine in BRAF V600E mutated metastatic colorectal cancer: a prospective multicentre phase II clinical study. <i>ESMO Open</i> , 2017, 2, e000241.	2.0	10
143	TRIBE-2: a phase III, randomized, open-label, strategy trial in unresectable metastatic colorectal cancer patients by the GONO group. <i>BMC Cancer</i> , 2017, 17, 408.	1.1	28
144	Estimating 12-week death probability in patients with refractory metastatic colorectal cancer: the Colon Life nomogram. <i>Annals of Oncology</i> , 2017, 28, 555-561.	0.6	43

#	ARTICLE	IF	CITATIONS
145	CXCR4 polymorphism predicts progression-free survival in metastatic colorectal cancer patients treated with first-line bevacizumab-based chemotherapy. <i>Pharmacogenomics Journal</i> , 2017, 17, 543-550.	0.9	11
146	Potential contribution of the study nurse to colorectal cancer (CRC) translational research. <i>Annals of Oncology</i> , 2017, 28, vi110.	0.6	1
147	Genetic variants of genes in CCL5/CCR5 pathway to predict regorafenib-induced hand-foot skin reaction in patients with refractory metastatic colorectal cancer: A report of ethnic difference.. <i>Journal of Clinical Oncology</i> , 2017, 35, 615-615.	0.8	1
148	Stereotactic Body Radiotherapy in Patients with Lung Oligometastases from Colorectal Cancer. <i>Anticancer Research</i> , 2017, 37, 315-320.	0.5	21
149	Surrogate Endpoints in Second-Line Trials of Targeted Agents in Metastatic Colorectal Cancer: A Literature-Based Systematic Review and Meta-Analysis. <i>Cancer Research and Treatment</i> , 2017, 49, 834-845.	1.3	12
150	Genetic variations in semaphorin/neuropilin signaling to predict clinical outcome in patients (pts) with metastatic colorectal cancer (mCRC) receiving bevacizumab-based chemotherapy.. <i>Journal of Clinical Oncology</i> , 2017, 35, 11608-11608.	0.8	0
151	Partial splenic embolization in chemotherapy-induced thrombocytopenia: A retrospective analysis with long term follow-up.. <i>Journal of Clinical Oncology</i> , 2017, 35, e21654-e21654.	0.8	0
152	<b>Topoisomerase 1 Promoter Variants and Benefit from Irinotecan in Metastatic Colorectal Cancer Patients.</b> <i>Oncology</i> , 2016, 91, 283-288.	0.9	5
153	FOLFOXIRI plus bevacizumab (bev) followed by maintenance with bev alone or bev plus metronomic chemotherapy (metroCT) in metastatic colorectal cancer (mCRC): The phase II randomized MOMA trial. <i>Annals of Oncology</i> , 2016, 27, vi560.	0.6	7
154	Challenging chemoresistant metastatic colorectal cancer: therapeutic strategies from the clinic and from the laboratory. <i>Annals of Oncology</i> , 2016, 27, 1456-1466.	0.6	51
155	Ramucirumab for the treatment of gastric cancers, colorectal adenocarcinomas, and other gastrointestinal malignancies. <i>Expert Review of Clinical Pharmacology</i> , 2016, 9, 877-885.	1.3	11
156	Location of Primary Tumor and Benefit From Anti-Epidermal Growth Factor Receptor Monoclonal Antibodies in Patients With <i>RAS</i> and <i>BRAF</i> Wild-Type Metastatic Colorectal Cancer. <i>Oncologist</i> , 2016, 21, 988-994.	1.9	94
157	Angiogenesis genotyping and clinical outcome during regorafenib treatment in metastatic colorectal cancer patients. <i>Scientific Reports</i> , 2016, 6, 25195.	1.6	25
158	Oral Chemotherapy and Patient Perspective in Solid Tumors: A National Survey by the Italian Association of Medical Oncology. <i>Tumori</i> , 2016, 102, 108-113.	0.6	4
159	Modified FOLFOXIRI (mFOLFOXIRI) plus cetuximab (cet), followed by cet or bevacizumab (bev) maintenance, in <i>RAS/BRAF</i> wt metastatic colorectal cancer (mCRC): The phase II randomized MACBETH trial by GONO. <i>Annals of Oncology</i> , 2016, 27, vi152.	0.6	3
160	Clinical Significance of <i>TLR1</i> I602S Polymorphism for Patients with Metastatic Colorectal Cancer Treated with FOLFIRI plus Bevacizumab. <i>Molecular Cancer Therapeutics</i> , 2016, 15, 1740-1745.	1.9	9
161	CDX2 as a Prognostic Biomarker in Colon Cancer. <i>New England Journal of Medicine</i> , 2016, 374, 2182-2184.	13.9	23
162	FOLFOXIRI or FOLFOXIRI plus bevacizumab as first-line treatment of metastatic colorectal cancer: a propensity score-adjusted analysis from two randomized clinical trials. <i>Annals of Oncology</i> , 2016, 27, 843-849.	0.6	46

#	ARTICLE	IF	CITATIONS
163	A Review of Clinical Studies and Practical Guide for the Administration of Triplet Chemotherapy Regimens with Bevacizumab in First-line Metastatic Colorectal Cancer. Targeted Oncology, 2016, 11, 293-308.	1.7	18
164	Prognostic Impact of <i>IL6</i> Genetic Variants in Patients with Metastatic Colorectal Cancer Treated with Bevacizumab-Based Chemotherapy. Clinical Cancer Research, 2016, 22, 3218-3226.	3.2	21
165	Clinico-pathological nomogram for predicting BRAF mutational status of metastatic colorectal cancer. British Journal of Cancer, 2016, 114, 30-36.	2.9	56
166	Body Mass Index Is Prognostic in Metastatic Colorectal Cancer: Pooled Analysis of Patients From First-Line Clinical Trials in the ARCAD Database. Journal of Clinical Oncology, 2016, 34, 144-150.	0.8	116
167	Females versus males: Clinical features and outcome differences in large molecularly selected cohort of mCRC patients.. Journal of Clinical Oncology, 2016, 34, 3540-3540.	0.8	1
168	Modified FOLFOXIRI (mFOLFOXIRI) plus cetuximab (cet), followed by cet or bevacizumab (bev) maintenance, in <i>RAS</i> / <i>BRAF</i> wild-type (wt) metastatic colorectal cancer (mCRC): Results of the phase II randomized MACBETH trial by GONO.. Journal of Clinical Oncology, 2016, 34, 3543-3543.	0.8	9
169	Genetic variants of ATM and XRCC3 to predict efficacy of TAS-102 in patients with refractory metastatic colorectal cancer.. Journal of Clinical Oncology, 2016, 34, 3579-3579.	0.8	3
170	Genetic variants of <i>hENT-1</i> to predict efficacy of TAS-102 in patients with refractory metastatic colorectal cancer.. Journal of Clinical Oncology, 2016, 34, 3580-3580.	0.8	2
171	Genetic variations associated with cancer cachexia pathways to predict survival in metastatic colorectal cancer (mCRC): Results from FIRE-3 and TRIBE.. Journal of Clinical Oncology, 2016, 34, 3590-3590.	0.8	1
172	Safety and efficacy of FOLFOXIRI with or without targeted agents as first-line treatment of selected elderly metastatic colorectal cancer patients: A pooled analysis of GONO studies.. Journal of Clinical Oncology, 2016, 34, e15054-e15054.	0.8	0
173	TRIBE-2 by GONO group: A phase III strategy study in the first- and second-line treatment of unresectable metastatic colorectal cancer (mCRC) patients.. Journal of Clinical Oncology, 2016, 34, TPS3629-TPS3629.	0.8	0
174	Lactate dehydrogenase (LDH) levels to predict benefit from the continuation of bevacizumab (bev) beyond progression in metastatic colorectal cancer (mCRC): Subgroup analysis of the randomized BEBYP study.. Journal of Clinical Oncology, 2016, 34, e15127-e15127.	0.8	0
175	Randomized phase II study of first-line FOLFOX plus panitumumab (pan) versus 5FU plus pan in elderly <i>RAS</i> and <i>BRAF</i> wild-type (wt) metastatic colorectal cancer (mCRC) patients (pts): The PANDA study.. Journal of Clinical Oncology, 2016, 34, TPS3627-TPS3627.	0.8	0
176	Genetic variants of <i>Pin1</i> to predict benefit from irinotecan and oxaliplatin based treatment in patients with metastatic colorectal cancer (mCRC).. Journal of Clinical Oncology, 2016, 34, 11589-11589.	0.8	0
177	Identifying predictive SNPs in patients with metastatic colorectal cancer (mCRC) using Random Survival Forests.. Journal of Clinical Oncology, 2016, 34, 3606-3606.	0.8	0
178	NOS2 polymorphisms in the prediction of benefit from FOLFIRI plus bevacizumab in mCRC patients enrolled in TRIBE trial.. Journal of Clinical Oncology, 2016, 34, 11597-11597.	0.8	0
179	Identifying SNPs associated with progression-free survival (PFS) and overall survival (OS) in patients with <i>KRAS</i> wildtype and mutant metastatic colorectal cancer (mCRC) using Random Survival Forests (RSF).. Journal of Clinical Oncology, 2016, 34, 3604-3604.	0.8	0
180	IRS1 and IRS2 polymorphisms and outcome in mCRC patients enrolled in the FIRE-3 trial.. Journal of Clinical Oncology, 2016, 34, 11600-11600.	0.8	0

#	ARTICLE	IF	CITATIONS
181	Abstract 3265: Homeobox B9 (HOXB9) sustains anti-VEGF treatment resistance in gastrointestinal tumors. , 2016, , .		0
182	CDX2 as a Prognostic Biomarker in Colon Cancer. <i>New England Journal of Medicine</i> , 2016, 374, 2183.	13.9	5
183	TRIBE study: are all three cytotoxic drugs crucial? â€œ Authors' reply. <i>Lancet Oncology</i> , The, 2015, 16, e578-e579.	5.1	0
184	Molecular and pathological characterization of the EZH2 rs3757441 single nucleotide polymorphism in colorectal cancer. <i>BMC Cancer</i> , 2015, 15, 874.	1.1	10
185	FOLFOXIRI and Bevacizumab for Metastatic Colorectal Cancer. <i>New England Journal of Medicine</i> , 2015, 372, 290-292.	13.9	11
186	Clonal evolution and resistance to EGFR blockade in the blood of colorectal cancer patients. <i>Nature Medicine</i> , 2015, 21, 795-801.	15.2	809
187	First-line anti-EGFR monoclonal antibodies in panRAS wild-type metastatic colorectal cancer: A systematic review and meta-analysis. <i>Critical Reviews in Oncology/Hematology</i> , 2015, 96, 156-166.	2.0	61
188	TAS-102, a novel antitumor agent: A review of the mechanism of action. <i>Cancer Treatment Reviews</i> , 2015, 41, 777-783.	3.4	115
189	Basal and bevacizumab-based therapy-induced changes of lactate dehydrogenases and fibrinogen levels and clinical outcome of previously untreated metastatic colorectal cancer patients: a multicentric retrospective analysis. <i>Expert Opinion on Biological Therapy</i> , 2015, 15, 155-162.	1.4	27
190	Distinct gene expression profiles of proximal and distal colorectal cancer: implications for cytotoxic and targeted therapy. <i>Pharmacogenomics Journal</i> , 2015, 15, 354-362.	0.9	41
191	Integrin genetic variants and stage-specific tumor recurrence in patients with stage II and III colon cancer. <i>Pharmacogenomics Journal</i> , 2015, 15, 226-234.	0.9	14
192	Continuation or reintroduction of bevacizumab beyond progression to first-line therapy in metastatic colorectal cancer: final results of the randomized BEBYP trial. <i>Annals of Oncology</i> , 2015, 26, 724-730.	0.6	136
193	Primary Tumor Location as a Prognostic Factor in Metastatic Colorectal Cancer. <i>Journal of the National Cancer Institute</i> , 2015, 107, .	3.0	385
194	First-line chemotherapy for mCRCâ€™a review and evidence-based algorithm. <i>Nature Reviews Clinical Oncology</i> , 2015, 12, 607-619.	12.5	138
195	BRAF codons 594 and 596 mutations identify a new molecular subtype of metastatic colorectal cancer at favorable prognosis. <i>Annals of Oncology</i> , 2015, 26, 2092-2097.	0.6	137
196	BRAF and RAS mutations as prognostic factors in metastatic colorectal cancer patients undergoing liver resection. <i>British Journal of Cancer</i> , 2015, 112, 1921-1928.	2.9	146
197	Early tumor shrinkage and depth of response predict long-term outcome in metastatic colorectal cancer patients treated with first-line chemotherapy plus bevacizumab: results from phase III TRIBE trial by the Gruppo Oncologico del Nord Ovest. <i>Annals of Oncology</i> , 2015, 26, 1188-1194.	0.6	153
198	FCGR polymorphisms and cetuximab efficacy in chemorefractory metastatic colorectal cancer: an international consortium study. <i>Gut</i> , 2015, 64, 921-928.	6.1	22

#	ARTICLE	IF	CITATIONS
199	Variations in genes regulating tumor-associated macrophages (TAMs) to predict outcomes of bevacizumab-based treatment in patients with metastatic colorectal cancer: results from TRIBE and FIRE3 trials. <i>Annals of Oncology</i> , 2015, 26, 2450-2456.	0.6	29
200	Polymorphisms in Genes Involved in EGFR Turnover Are Predictive for Cetuximab Efficacy in Colorectal Cancer. <i>Molecular Cancer Therapeutics</i> , 2015, 14, 2374-2381.	1.9	4
201	BRAF-mutated metastatic colorectal cancer between past and future. <i>British Journal of Cancer</i> , 2015, 113, 1634-1635.	2.9	11
202	Single-Agent Panitumumab in Frail Elderly Patients With Advanced <i>RAS</i> and <i>BRAF</i> Wild-Type Colorectal Cancer: Challenging Drug Label to Light Up New Hope. <i>Oncologist</i> , 2015, 20, 1261-1265.	1.9	42
203	TAS-102 for the treatment of metastatic colorectal cancer. <i>Expert Review of Anticancer Therapy</i> , 2015, 15, 1283-1292.	1.1	12
204	Assessment of a HER2 scoring system for colorectal cancer: results from a validation study. <i>Modern Pathology</i> , 2015, 28, 1481-1491.	2.9	226
205	Response. <i>Journal of the National Cancer Institute</i> , 2015, 107, djv205.	3.0	2
206	FOLFOXIRI plus bevacizumab versus FOLFIRI plus bevacizumab as first-line treatment of patients with metastatic colorectal cancer: updated overall survival and molecular subgroup analyses of the open-label, phase 3 TRIBE study. <i>Lancet Oncology</i> , The, 2015, 16, 1306-1315.	5.1	835
207	Phase II study of single-agent cetuximab in KRAS G13D mutant metastatic colorectal cancer. <i>Annals of Oncology</i> , 2015, 26, 2503.	0.6	18
208	Genetic variants of kinase suppressors of Ras (KSR1) to predict survival in patients with ER $\pm$ -positive advanced breast cancer. <i>Pharmacogenomics Journal</i> , 2015, 15, 235-240.	0.9	2
209	Role of <i>NRAS</i> mutations as prognostic and predictive markers in metastatic colorectal cancer. <i>International Journal of Cancer</i> , 2015, 136, 83-90.	2.3	126
210	Genes involved in pericyte-driven tumor maturation predict treatment benefit of first-line FOLFIRI plus bevacizumab in patients with metastatic colorectal cancer. <i>Pharmacogenomics Journal</i> , 2015, 15, 69-76.	0.9	25
211	FOLFOXIRI plus bevacizumab versus FOLFIRI plus bevacizumab as initial treatment for metastatic colorectal cancer (TRIBE study): Updated survival results and final molecular subgroups analyses.. <i>Journal of Clinical Oncology</i> , 2015, 33, 3510-3510.	0.8	8
212	DPYD c.1905+1G>A and c.2846A>T and UGT1A1*28 allelic variants as predictors of toxicity: Pharmacogenetic translational analysis from the phase III TRIBE study in metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2015, 33, 3532-3532.	0.8	2
213	Variations in genes regulating tumor-associated macrophages (TAMs) to predict outcome of bevacizumab (bev)-based treatment in patients with metastatic colorectal cancer (mCRC): Results from TRIBE and FIRE3 trials.. <i>Journal of Clinical Oncology</i> , 2015, 33, 3552-3552.	0.8	1
214	Angiogenesis genotyping and clinical outcome during regorafenib treatment in metastatic colorectal cancer patients.. <i>Journal of Clinical Oncology</i> , 2015, 33, 595-595.	0.8	1
215	FOLFOXIRI plus bevacizumab (bev) versus FOLFIRI plus bev as first-line treatment of metastatic colorectal cancer (mCRC): Updated survival results of the phase III TRIBE trial by the GONO group.. <i>Journal of Clinical Oncology</i> , 2015, 33, 657-657.	0.8	17
216	Prognostic clinical factors in pretreated colorectal cancer patients receiving regorafenib: Implications for clinical management. <i>Oncotarget</i> , 2015, 6, 33982-33992.	0.8	46

#	ARTICLE	IF	CITATIONS
217	Prognostic significance of <i>K-Ras</i> mutation rate in metastatic colorectal cancer patients. <i>Oncotarget</i> , 2015, 6, 31604-31612.	0.8	30
218	Polymorphism of the chemokine CXCR4 to predict treatment benefit of first-line bevacizumab-based chemotherapy in patients with metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2015, 33, 635-635.	0.8	0
219	Prognostic clinical factors in pretreated colorectal cancer patients receiving regorafenib: Implications for clinical management.. <i>Journal of Clinical Oncology</i> , 2015, 33, 591-591.	0.8	0
220	Macrophage polarization related gene variants to predict clinical outcome in metastatic colorectal cancer (mCRC) patients (pts) treated with bevacizumab (bev) in combination with FOLFIRI.. <i>Journal of Clinical Oncology</i> , 2015, 33, 621-621.	0.8	0
221	Effect of genetic variation on overall survival in a clinical trial of metastatic colorectal cancer (mCRC).. <i>Journal of Clinical Oncology</i> , 2015, 33, 3562-3562.	0.8	0
222	Genetic variant of TWEAK to predict clinical outcome in mCRC patients (pts) treated with first line FOLFIRI and Bevacizumab (FOLFIRI/BEV) in FIRE-3 and TRIBE cohorts.. <i>Journal of Clinical Oncology</i> , 2015, 33, 3554-3554.	0.8	0
223	Glucose metabolism enzymes gene expression analysis and selective metabolic advantage in the progression of colorectal cancer (CRC).. <i>Journal of Clinical Oncology</i> , 2015, 33, e14519-e14519.	0.8	0
224	Prognostic significance of KRAS mutation rate in metastatic colorectal cancer (mCRC) patients (pts).. <i>Journal of Clinical Oncology</i> , 2015, 33, e22075-e22075.	0.8	0
225	Genetic variants of kinases suppressors of Ras (KSR) to predict tumor response to first-line cetuximab in patients with mCRC: Prospective analysis in the FIRE 3 trial.. <i>Journal of Clinical Oncology</i> , 2015, 33, 3613-3613.	0.8	0
226	Abstract 616: Blood-based molecular landscapes of resistance to EGFR blockade in colorectal cancer patients. , 2015, , .		0
227	Bevacizumab and first-line chemotherapy for older patients with advanced colorectal cancer: final results of a Community-based Observational Italian Study. <i>Anticancer Research</i> , 2015, 35, 2391-9.	0.5	1
228	KRAS and BRAF genotyping of synchronous colorectal carcinomas. <i>Oncology Letters</i> , 2014, 7, 1532-1536.	0.8	7
229	Gender-specific profiling in SCN1A polymorphisms and time-to-recurrence in patients with stage II/III colorectal cancer treated with adjuvant 5-fluorouracil chemotherapy. <i>Pharmacogenomics Journal</i> , 2014, 14, 135-141.	0.9	11
230	Prospective study of EGFR intron 1 (CA) <sub>n</sub> repeats variants as predictors of benefit from cetuximab and irinotecan in chemo-refractory metastatic colorectal cancer (mCRC) patients. <i>Pharmacogenomics Journal</i> , 2014, 14, 322-327.	0.9	11
231	Biomarkers and Response to Bevacizumab Letter. <i>Clinical Cancer Research</i> , 2014, 20, 1056-1057.	3.2	8
232	EGFR ligands as pharmacodynamic biomarkers in metastatic colorectal cancer patients treated with cetuximab and irinotecan. <i>Targeted Oncology</i> , 2014, 9, 205-214.	1.7	27
233	FOLFOXIRI plus bevacizumab as first-line treatment in BRAF mutant metastatic colorectal cancer. <i>European Journal of Cancer</i> , 2014, 50, 57-63.	1.3	162
234	Initial Therapy with FOLFOXIRI and Bevacizumab for Metastatic Colorectal Cancer. <i>New England Journal of Medicine</i> , 2014, 371, 1609-1618.	13.9	845

#	ARTICLE	IF	CITATIONS
235	Role of immunoglobulin G fragment C receptor polymorphism-mediated antibody-dependant cellular cytotoxicity in colorectal cancer treated with cetuximab therapy. <i>Pharmacogenomics Journal</i> , 2014, 14, 14-19.	0.9	21
236	Association of common gene variants in the WNT/ $\beta$ -catenin pathway with colon cancer recurrence. <i>Pharmacogenomics Journal</i> , 2014, 14, 142-150.	0.9	28
237	Subgroup analyses in RAS mutant, BRAF mutant and all-wt mCRC pts treated with FOLFOXIRI plus bevacizumab (bev) or FOLFIRI plus bev in the TRIBE study.. <i>Journal of Clinical Oncology</i> , 2014, 32, 3519-3519.	0.8	9
238	Phase II study of single-agent cetuximab in KRAS G13D mutant metastatic colorectal cancer (mCRC).. <i>Journal of Clinical Oncology</i> , 2014, 32, 3524-3524.	0.8	6
239	Modified FOLFOXIRI plus cetuximab (cet) as induction treatment in unresectable metastatic colorectal cancer (mCRC) patients (pts): Preliminary results of the phase II randomized Macbeth trial by GONO group.. <i>Journal of Clinical Oncology</i> , 2014, 32, 3596-3596.	0.8	4
240	mRNA expression levels of candidate genes and clinical outcome in mCRC patients treated with FOLFOXIRI plus bevacizumab (bev) or FOLFIRI plus bev in the TRIBE study.. <i>Journal of Clinical Oncology</i> , 2014, 32, 3640-3640.	0.8	1
241	Phase II randomized study of induction FOLFOXIRI plus bevacizumab (bev) followed by maintenance with bev alone or bev plus metronomic chemotherapy (metroCT) in metastatic colorectal cancer (mCRC): The MOMA trial.. <i>Journal of Clinical Oncology</i> , 2014, 32, TPS3664-TPS3664.	0.8	2
242	Early tumor shrinkage (ETS) and deepness of response (DoR) to predict progression-free, postprogression, and overall survival: Results from the phase III TRIBE trial.. <i>Journal of Clinical Oncology</i> , 2014, 32, 521-521.	0.8	2
243	BRAF and KRAS mutations in liver-resected metastatic colorectal cancer (mCRC) patients (pts).. <i>Journal of Clinical Oncology</i> , 2014, 32, 476-476.	0.8	0
244	The Role of Metronomic Chemotherapy in the Treatment of Metastatic Colorectal Cancer Patients. , 2014, , 135-142.		0
245	CCL2 polymorphism as a predictive marker for bevacizumab (Bev) in combination with FOLFIRI as first-line treatment in metastatic colorectal cancer (mCRC) patients (pts).. <i>Journal of Clinical Oncology</i> , 2014, 32, e14556-e14556.	0.8	1
246	Association of the prognostic role of CXCR4/CXCL12 polymorphism with treatment outcomes after bevacizumab-based chemotherapy in metastatic colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2014, 32, e22008-e22008.	0.8	0
247	High-throughput exome array for identification of novel polymorphisms associated with clinical outcome in mCRC patients treated with first-line FOLFOXIRI/BEV versus FOLFIRI/BEV (TRIBE trial); Tj ETQq1 1 0.784314 rgBT (Overloc		
248	Biomarker validation study: Genes involved in ubiquitin proteasome system (UPS) dependent EGFR-degradation for prediction of efficacy in metastatic colorectal cancer patients treated with cetuximab.. <i>Journal of Clinical Oncology</i> , 2014, 32, 3571-3571.	0.8	0
249	Body mass index (BMI) as prognostic in metastatic colorectal cancer (mCRC): A pooled analysis of 21 first-line trials in the ARCAD database.. <i>Journal of Clinical Oncology</i> , 2014, 32, 3537-3537.	0.8	0
250	Common genetic variants in genes involved in the Hippo pathway: Novel biomarkers in metastatic colorectal cancer patients treated with irinotecan plus cetuximab.. <i>Journal of Clinical Oncology</i> , 2014, 32, e14523-e14523.	0.8	0
251	Genetic variants of CBP and SOX9 to predict clinical outcome in metastatic colorectal cancer (mCRC) patients (pts) treated with first-line FOLFIRI and bevacizumab (FOLFIRI/BEV).. <i>Journal of Clinical Oncology</i> , 2014, 32, e14528-e14528.	0.8	0
252	Liquid biopsy: monitoring cancer-genetics in the blood. <i>Nature Reviews Clinical Oncology</i> , 2013, 10, 472-484.	12.5	1,482

#	ARTICLE	IF	CITATIONS
253	Adjuvant Systemic Chemotherapy After Putative Curative Resection of Colorectal Liver and Lung Metastases. <i>Clinical Colorectal Cancer</i> , 2013, 12, 188-194.	1.0	28
254	Histopathologic evaluation of liver metastases from colorectal cancer in patients treated with FOLFOXIRI plus bevacizumab. <i>British Journal of Cancer</i> , 2013, 108, 2549-2556.	2.9	51
255	Dicer and Drosha expression and response to Bevacizumab-based therapy in advanced colorectal cancer patients. <i>European Journal of Cancer</i> , 2013, 49, 1501-1508.	1.3	19
256	Bevacizumab Beyond Progression in Metastatic Colorectal Cancer Patients Receiving a First-Line Treatment Containing Bevacizumab: Update of Bebyp Trial by Gono. <i>Annals of Oncology</i> , 2013, 24, iv22.	0.6	2
257	FOLFOXIRI in combination with panitumumab as first-line treatment in quadruple wild-type (KRAS,) Tj ETQq1 1 0.784314 rgBT /Overlo Nord Ovest (GONO). <i>Annals of Oncology</i> , 2013, 24, 2062-2067.	0.6	86
258	Prognostic Role of Lemur Tyrosine Kinase-3 Germline Polymorphisms in Adjuvant Gastric Cancer in Japan and the United States. <i>Molecular Cancer Therapeutics</i> , 2013, 12, 2261-2272.	1.9	19
259	Caveolin-1 is a novel regulator of K-RAS-dependent migration in colon carcinogenesis. <i>International Journal of Cancer</i> , 2013, 133, 43-57.	2.3	45
260	Reply: Comment on "Histopathologic evaluation of liver metastases from colorectal cancer patients treated with FOLFOXIRI plus bevacizumab". <i>British Journal of Cancer</i> , 2013, 109, 3129-3130.	2.9	3
261	FOLFOXIRI/Bevacizumab Versus FOLFIRI/Bevacizumab as First-Line Treatment in Unresectable Metastatic Colorectal Cancer: Results of Phase III Tribe Trial by Gono Group. <i>Annals of Oncology</i> , 2013, 24, iv21.	0.6	5
262	Long-Survivors with Lung Metastases and Kras Mutations Have an Increased Risk to Develop Brain Metastases From Colorectal Cancer. <i>Annals of Oncology</i> , 2013, 24, iv15.	0.6	1
263	Prospective Analysis of the Early Modulation of Plasma Amphiregulin During Treatment with Cetuximab and Irinotecan in Metastatic Colorectal Cancer Patients. <i>Annals of Oncology</i> , 2013, 24, iv28.	0.6	0
264	Conference Scene: Annual Meeting of the American Society of Clinical Oncology. <i>Colorectal Cancer</i> , 2013, 2, 401-404.	0.8	0
265	Oral multikinase inhibitor regorafenib for the treatment of patients with metastatic colorectal cancer. <i>Colorectal Cancer</i> , 2013, 2, 411-417.	0.8	1
266	Prospective Validation of Candidate SNPs of VEGF/VEGFR Pathway in Metastatic Colorectal Cancer Patients Treated with First-Line FOLFIRI Plus Bevacizumab. <i>PLoS ONE</i> , 2013, 8, e66774.	1.1	64
267	FOLFOXIRI/bevacizumab (bev) versus FOLFIRI/bev as first-line treatment in unresectable metastatic colorectal cancer (mCRC) patients (pts): Results of the phase III TRIBE trial by GONO group.. <i>Journal of Clinical Oncology</i> , 2013, 31, 3505-3505.	0.8	46
268	Gene expression profiles and tumor locations in colorectal cancer (left vs. right vs. rectum).. <i>Journal of Clinical Oncology</i> , 2013, 31, 3527-3527.	0.8	8
269	Analysis of NRAS mutation as poor prognostic indicator and predictor of resistance to anti-EGFR monoclonal antibodies (anti-EGFRs) in metastatic colorectal cancer (mCRC) patients (pts).. <i>Journal of Clinical Oncology</i> , 2013, 31, 3613-3613.	0.8	4
270	Metronomic capecitabine (cape) and cyclophosphamide (CTX) for refractory metastatic colorectal cancer (mCRC): Results of a phase II trial.. <i>Journal of Clinical Oncology</i> , 2013, 31, e14577-e14577.	0.8	2



#	ARTICLE	IF	CITATIONS
271	FOLFOXIRI plus bevacizumab (bev) versus FOLFIRI plus bev as first-line treatment of metastatic colorectal cancer (mCRC): Results of the phase III randomized TRIBE trial.. Journal of Clinical Oncology, 2013, 31, 336-336.	0.8	25
272	Circulating angiogenic factors as predictors of benefit from bevacizumab (bev) beyond progression in metastatic colorectal cancer (mCRC): Translational analyses from the phase III BEBYP trial.. Journal of Clinical Oncology, 2013, 31, 382-382.	0.8	1
273	Gender specific profiling in SCN1A polymorphisms and time to recurrence in patients with stage II/III colorectal cancer.. Journal of Clinical Oncology, 2013, 31, 393-393.	0.8	1
274	Correlation of messenger RNA expression patterns of ERCC1, TS, EGFR, and VEGFR2 with KRAS and BRAF mutational status in advanced colorectal cancer: Implications for targeted therapies.. Journal of Clinical Oncology, 2013, 31, 383-383.	0.8	2
275	“HER Majesty’s a Pretty Nice Girl but She Changes From Day to Day” Journal of Clinical Oncology, 2012, 30, 465-466.	0.8	1
276	High Let-7a MicroRNA Levels in KRAS-Mutated Colorectal Carcinomas May Rescue Anti-EGFR Therapy Effects in Patients with Chemotherapy-Refractory Metastatic Disease. Oncologist, 2012, 17, 823-829.	1.9	74
277	Prognosis of mucinous histology for patients with radically resected stage II and III colon cancer. Annals of Oncology, 2012, 23, 135-141.	0.6	79
278	Influence of Sex on the Survival of Patients With Esophageal Cancer. Journal of Clinical Oncology, 2012, 30, 2265-2272.	0.8	112
279	Analysis of HER-3, insulin growth factor-1, nuclear factor-kB and epidermal growth factor receptor gene copy number in the prediction of clinical outcome for K-RAS wild-type colorectal cancer patients receiving irinotecan and cetuximab. Annals of Oncology, 2012, 23, 1706-1712.	0.6	34
280	Selecting the best targeted agent in first-line treatment of unresectable liver metastases from colorectal cancer: does the bench have the answers?. Journal of Hepato-Biliary-Pancreatic Sciences, 2012, 19, 528-535.	1.4	2
281	The possible role of chemotherapy in antiangiogenic drug resistance. Medical Hypotheses, 2012, 78, 646-648.	0.8	17
282	Outcome of Second-Line Treatment After First-Line Chemotherapy With the GONO FOLFOXIRI Regimen. Clinical Colorectal Cancer, 2012, 11, 71-76.	1.0	17
283	EZH2 polymorphism and benefit from bevacizumab in colorectal cancer: another piece to the puzzle. Annals of Oncology, 2012, 23, 1370-1371.	0.6	7
284	An EZH2 polymorphism is associated with clinical outcome in metastatic colorectal cancer patients. Annals of Oncology, 2012, 23, 1207-1213.	0.6	40
285	Upfront Chemotherapy Regimens in Unresectable Disease: One, Two, or Three Cytotoxics?. Current Colorectal Cancer Reports, 2012, 8, 153-160.	1.0	0
286	Pharmacogenetic Concerns in Metastatic Colorectal Cancer Therapy. Current Colorectal Cancer Reports, 2012, 8, 263-271.	1.0	1
287	PD-0009 Primary Tumor Location is a Major Independent Prognostic Factor for Mrcr Patients. Annals of Oncology, 2012, 23, iv22.	0.6	0
288	P-0262 Prospective Evaluation of Candidate Snps of Vegf/Vegfr Pathway in Metastatic Colorectal Cancer Patients Treated with First-Line Folfiri Plus Bevacizumab (BV). Annals of Oncology, 2012, 23, iv105-iv106.	0.6	0

#	ARTICLE	IF	CITATIONS
289	Cancer Dormancy: A Model of Early Dissemination and Late Cancer Recurrence. <i>Clinical Cancer Research</i> , 2012, 18, 645-653.	3.2	173
290	Circulating endothelial cells and their apoptotic fraction are mutually independent predictive biomarkers in Bevacizumab-based treatment for advanced colorectal cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2012, 138, 1187-1196.	1.2	27
291	Clinical, pharmacokinetic and pharmacodynamic evaluations of metronomic UFT and cyclophosphamide plus celecoxib in patients with advanced refractory gastrointestinal cancers. <i>Angiogenesis</i> , 2012, 15, 275-286.	3.7	61
292	Clinical impact of anti-epidermal growth factor receptor monoclonal antibodies in first-line treatment of metastatic colorectal cancer. <i>Cancer</i> , 2012, 118, 1523-1532.	2.0	34
293	PML as a potential predictive factor of oxaliplatin/fluoropyrimidine-based first line chemotherapy efficacy in colorectal cancer patients. <i>Journal of Cellular Physiology</i> , 2012, 227, 927-933.	2.0	5
294	Prospective evaluation of candidate SNPs of VEGF/VEGFR pathway in metastatic colorectal cancer (mCRC) patients (pts) treated with first-line FOLFIRI plus bevacizumab (BV).. <i>Journal of Clinical Oncology</i> , 2012, 30, 3518-3518.	0.8	1
295	Prospective study of EGFR intron 1 CA tandem repeats to predict factor benefit from cetuximab and irinotecan.. <i>Journal of Clinical Oncology</i> , 2012, 30, 3540-3540.	0.8	6
296	LMTK3 polymorphism in patients with metastatic colon cancer.. <i>Journal of Clinical Oncology</i> , 2012, 30, 471-471.	0.8	6
297	Cytotoxic triplets plus a biologic: state-of-the-art in maximizing the potential of up-front medical treatment of metastatic colorectal cancer. <i>Expert Opinion on Biological Therapy</i> , 2011, 11, 519-531.	1.4	3
298	Retrospective exploratory analysis of VEGF polymorphisms in the prediction of benefit from first-line FOLFIRI plus bevacizumab in metastatic colorectal cancer. <i>BMC Cancer</i> , 2011, 11, 247.	1.1	69
299	Epidermal growth factor receptor (EGFR) gene promoter methylation and cetuximab treatment in colorectal cancer patients. <i>British Journal of Cancer</i> , 2011, 104, 1786-1790.	2.9	65
300	Pharmacodynamic and pharmacogenetic angiogenesis-related markers of first-line FOLFOXIRI plus bevacizumab schedule in metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2011, 104, 1262-1269.	2.9	85
301	Cetuximab plus irinotecan after irinotecan failure in elderly metastatic colorectal cancer patients: Clinical outcome according to KRAS and BRAF mutational status. <i>Critical Reviews in Oncology/Hematology</i> , 2011, 78, 243-251.	2.0	31
302	Bevacizumab pharmacogenetics in tumor treatment: still looking for the right pieces of the puzzle. <i>Pharmacogenomics</i> , 2011, 12, 1077-1080.	0.6	11
303	Randomized Trial of Two Induction Chemotherapy Regimens in Metastatic Colorectal Cancer: An Updated Analysis. <i>Journal of the National Cancer Institute</i> , 2011, 103, 21-30.	3.0	160
304	Early magnesium modifications as a surrogate marker of efficacy of cetuximab-based anticancer treatment in KRAS wild-type advanced colorectal cancer patients. <i>Annals of Oncology</i> , 2011, 22, 1141-1146.	0.6	54
305	Should Oncologists Be Aware in Their Clinical Practice of KRAS Molecular Analysis?. <i>Journal of Clinical Oncology</i> , 2011, 29, e206-e207.	0.8	17
306	Targeting Vascular Endothelial Growth Factor Pathway in First-Line Treatment of Metastatic Colorectal Cancer: State-of-the-Art and Future Perspectives in Clinical and Molecular Selection of Patients. <i>Current Cancer Drug Targets</i> , 2010, 10, 37-45.	0.8	12

#	ARTICLE	IF	CITATIONS
307	Phase II study of sequential cisplatin plus 5-fluorouracil/leucovorin (5-FU/LV) followed by irinotecan plus 5-FU/LV followed by docetaxel plus 5-FU/LV in patients with metastatic gastric or gastro-oesophageal junction adenocarcinoma. <i>Cancer Chemotherapy and Pharmacology</i> , 2010, 66, 559-566.	1.1	5
308	Insulin-like growth factor 1 expression correlates with clinical outcome in KRAS wild type colorectal cancer patients treated with cetuximab and irinotecan. <i>International Journal of Cancer</i> , 2010, 127, 1941-1947.	2.3	67
309	Reply: KRAS status analysis and anti-EGFR therapies: is comprehensiveness a biologist's fancy or a clinical necessity?. <i>British Journal of Cancer</i> , 2010, 102, 1076-1077.	2.9	1
310	Host genetic variants in the IGF binding protein-3 impact on survival of patients with advanced gastric cancer treated with palliative chemotherapy. <i>Pharmacogenomics</i> , 2010, 11, 1247-1256.	0.6	6
311	High concordance of BRAF status between primary colorectal tumours and related metastatic sites: implications for clinical practice. <i>Annals of Oncology</i> , 2010, 21, 1565.	0.6	38
312	Capecitabine after gastrectomy for advanced gastric cancer: have we got the patient right?. <i>Annals of Oncology</i> , 2010, 21, 181.	0.6	5
313	Palliative treatment of unresectable metastatic colorectal cancer. <i>Expert Opinion on Pharmacotherapy</i> , 2010, 11, 63-77.	0.9	18
314	Predictors of Benefit in Colorectal Cancer Treated With Cetuximab: Are We Getting "Lost in Translation"? <i>Journal of Clinical Oncology</i> , 2010, 28, e173-e174.	0.8	4
315	Immunological Effects of Bevacizumab-Based Treatment in Metastatic Colorectal Cancer. <i>Oncology</i> , 2010, 79, 187-196.	0.9	77
316	How useful is adjuvant irinotecan in stage IV CRC?. <i>Nature Reviews Clinical Oncology</i> , 2010, 7, 190-191.	12.5	3
317	Bevacizumab with FOLFOXIRI (irinotecan, oxaliplatin, fluorouracil, and folinate) as first-line treatment for metastatic colorectal cancer: a phase 2 trial. <i>Lancet Oncology</i> , The, 2010, 11, 845-852.	5.1	234
318	Genetic modulation of the Let-7 microRNA binding to KRAS 3'-untranslated region and survival of metastatic colorectal cancer patients treated with salvage cetuximab+irinotecan. <i>Pharmacogenomics Journal</i> , 2010, 10, 458-464.	0.9	109
319	Magnitude of benefit of the addition of bevacizumab to first-line chemotherapy for metastatic colorectal cancer: meta-analysis of randomized clinical trials. <i>Journal of Experimental and Clinical Cancer Research</i> , 2010, 29, 58.	3.5	46
320	Circulating endothelial cells and endothelial progenitors as predictive markers of clinical response to bevacizumab-based first-line treatment in advanced colorectal cancer patients. <i>Annals of Oncology</i> , 2010, 21, 2382-2389.	0.6	94
321	Review: Beyond KRAS: perspectives on new potential markers of intrinsic and acquired resistance to epidermal growth factor receptor inhibitors in metastatic colorectal cancer. <i>Therapeutic Advances in Medical Oncology</i> , 2009, 1, 167-181.	1.4	7
322	Variations in the interleukin-1 receptor antagonist gene impact on survival of patients with advanced colorectal cancer. <i>Pharmacogenomics Journal</i> , 2009, 9, 78-84.	0.9	23
323	Do we need biopsies of metastases for colorectal cancer patients?. <i>British Journal of Cancer</i> , 2009, 101, 374-375.	2.9	2
324	Mucinous histology predicts for poor response rate and overall survival of patients with colorectal cancer and treated with first-line oxaliplatin- and/or irinotecan-based chemotherapy. <i>British Journal of Cancer</i> , 2009, 100, 881-887.	2.9	164

#	ARTICLE	IF	CITATIONS
325	A multicenter phase II study of the combination of oxaliplatin, irinotecan and capecitabine in the first-line treatment of metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2009, 100, 1720-1724.	2.9	30
326	Cigarettes smoking habit may reduce benefit from cetuximab-based treatment in advanced colorectal cancer patients. <i>Expert Opinion on Biological Therapy</i> , 2009, 9, 945-949.	1.4	15
327	Long-Term Outcome of Initially Unresectable Metastatic Colorectal Cancer Patients Treated with 5-Fluorouracil/Leucovorin, Oxaliplatin, and Irinotecan (FOLFOXIRI) Followed by Radical Surgery of Metastases. <i>Annals of Surgery</i> , 2009, 249, 420-425.	2.1	213
328	Epidermal Growth Factor Receptor (EGFR) gene copy number (GCN) correlates with clinical activity of irinotecan-cetuximab in K-RAS wild-type colorectal cancer: a fluorescence in situ (FISH) and chromogenic in situ hybridization (CISH) analysis. <i>BMC Cancer</i> , 2009, 9, 303.	1.1	66
329	VEGF gene polymorphisms and susceptibility to colorectal cancer disease in Italian population. <i>International Journal of Colorectal Disease</i> , 2009, 24, 165-170.	1.0	47
330	A dose finding and pharmacokinetic study of capecitabine in combination with oxaliplatin and irinotecan in metastatic colorectal cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2009, 63, 965-969.	1.1	15
331	Pharmacokinetics, a main actor in a many-sided approach to severe 5-FU toxicity prediction. <i>British Journal of Clinical Pharmacology</i> , 2009, 67, 132-134.	1.1	9
332	Refractory neuroendocrine tumor response to liposomal doxorubicin and capecitabine. <i>Nature Reviews Clinical Oncology</i> , 2009, 6, 670-674.	12.5	5
333	KRAS codon 61, 146 and BRAF mutations predict resistance to cetuximab plus irinotecan in KRAS codon 12 and 13 wild-type metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2009, 101, 715-721.	2.9	509
334	PTEN Expression and KRAS Mutations on Primary Tumors and Metastases in the Prediction of Benefit From Cetuximab Plus Irinotecan for Patients With Metastatic Colorectal Cancer. <i>Journal of Clinical Oncology</i> , 2009, 27, 2622-2629.	0.8	402
335	Long-Term Outcome of Unresectable Metastatic Colorectal Cancer: Does Adjuvant Chemotherapy Play a Role After Resection?. <i>Annals of Surgery</i> , 2009, 250, 655.	2.1	0
336	First-Line Systemic Chemotherapy with Folfoxiri Followed by Radical Surgical Resection of Metastases for the Treatment of Unresectable Metastatic Colorectal Cancer Patients. , 2009, , 285-293.		0
337	Liver-only metastatic colorectal cancer patients and thymidylate synthase polymorphisms for predicting response to 5-fluorouracil-based chemotherapy. <i>British Journal of Cancer</i> , 2008, 99, 716-721.	2.9	29
338	Triplet Combination of Fluoropyrimidines, Oxaliplatin, and Irinotecan in the First-Line Treatment of Metastatic Colorectal Cancer. <i>Clinical Colorectal Cancer</i> , 2008, 7, 7-14.	1.0	15
339	Pharmacogenetic profiling in patients with advanced colorectal cancer treated with first-line FOLFIRI chemotherapy. <i>Pharmacogenomics Journal</i> , 2008, 8, 278-288.	0.9	97
340	Molecular predictive factors of response to anti-EGFR antibodies in colorectal cancer patients. <i>European Journal of Cancer</i> , Supplement, 2008, 6, 86-90.	2.2	2
341	Pharmacogenetic Profiling for Cetuximab Plus Irinotecan Therapy in Patients With Refractory Advanced Colorectal Cancer. <i>Journal of Clinical Oncology</i> , 2008, 26, 1427-1434.	0.8	124
342	Optimal approach to potentially resectable liver metastases from colorectal cancer. <i>Expert Review of Anticancer Therapy</i> , 2008, 8, 1533-1539.	1.1	4

#	ARTICLE	IF	CITATIONS
343	A pharmacokinetic and pharmacodynamic study on metronomic irinotecan in metastatic colorectal cancer patients. <i>British Journal of Cancer</i> , 2008, 98, 1312-1319.	2.9	63
344	High Concordance of <i>KRAS</i> Status Between Primary Colorectal Tumors and Related Metastatic Sites: Implications for Clinical Practice. <i>Oncologist</i> , 2008, 13, 1270-1275.	1.9	218
345	EGF-receptor targeting with monoclonal antibodies in colorectal carcinomas: rationale for a pharmacogenomic approach. <i>Pharmacogenomics</i> , 2008, 9, 55-69.	0.6	12
346	First-line chemotherapy in metastatic colorectal cancer: new approaches and therapeutic algorithms. Always hit hard first?. <i>Current Opinion in Oncology</i> , 2008, 20, 459-465.	1.1	11
347	Nuclear Factor- $\kappa$ B Tumor Expression Predicts Response and Survival in Irinotecan-Refractory Metastatic Colorectal Cancer Treated With Cetuximab-Irinotecan Therapy. <i>Journal of Clinical Oncology</i> , 2007, 25, 3930-3935.	0.8	121
348	Vascular Endothelial Growth Factor Levels in Immunodepleted Plasma of Cancer Patients As a Possible Pharmacodynamic Marker for Bevacizumab Activity. <i>Journal of Clinical Oncology</i> , 2007, 25, 1816-1818.	0.8	56
349	Pharmacogenetic Profiling in Patients With Advanced Colorectal Cancer Treated With First-Line FOLFOX-4 Chemotherapy. <i>Journal of Clinical Oncology</i> , 2007, 25, 1247-1254.	0.8	250
350	Treatment with 5-Fluorouracil/Folinic Acid, Oxaliplatin, and Irinotecan Enables Surgical Resection of Metastases in Patients With Initially Unresectable Metastatic Colorectal Cancer. <i>Annals of Surgical Oncology</i> , 2006, 13, 58-65.	0.7	156
351	First-line 5-fluorouracil/folinic acid, oxaliplatin and irinotecan (FOLFOXIRI) does not impair the feasibility and the activity of second line treatments in metastatic colorectal cancer. <i>Annals of Oncology</i> , 2006, 17, 1249-1254.	0.6	22
352	Are Dose-Finding Studies Still Necessary When Targeted Therapy Is Associated With Chemotherapy?. <i>Journal of Clinical Oncology</i> , 2006, 24, 4668-4669.	0.8	0