

# Pablo Tornero

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2277390/publications.pdf>

Version: 2024-02-01

28  
papers

2,083  
citations

304743

22  
h-index

501196

28  
g-index

28  
all docs

28  
docs citations

28  
times ranked

2623  
citing authors

#	ARTICLE	IF	CITATIONS
1	Gibberellins regulate ovule number through a DELLA-CUC2 complex in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2022, 110, 43-57.	5.7	14
2	On the Way to Ovules: The Hormonal Regulation of Ovule Development. <i>Critical Reviews in Plant Sciences</i> , 2020, 39, 431-456.	5.7	20
3	Gibberellin-mediated RGA-LIKE1 degradation regulates embryo sac development in <i>Arabidopsis</i> . <i>Journal of Experimental Botany</i> , 2020, 71, 7059-7072.	4.8	14
4	Regulation of ovule initiation by gibberellins and brassinosteroids in tomato and <i>Arabidopsis</i> : two plant species, two molecular mechanisms. <i>Plant Journal</i> , 2020, 102, 1026-1041.	5.7	26
5	NPR1 paralogs of <i>Arabidopsis</i> and their role in salicylic acid perception. <i>PLoS ONE</i> , 2018, 13, e0209835.	2.5	44
6	$\beta$ -carbonic anhydrases play a role in salicylic acid perception in <i>Arabidopsis</i> . <i>PLoS ONE</i> , 2017, 12, e0181820.	2.5	44
7	An Allele of <i>Arabidopsis</i> COI1 with Hypo- and Hypermorphic Phenotypes in Plant Growth, Defence and Fertility. <i>PLoS ONE</i> , 2013, 8, e55115.	2.5	1
8	Specific Missense Alleles of the <i>Arabidopsis</i> Jasmonic Acid Co-Receptor COI1 Regulate Innate Immune Receptor Accumulation and Function. <i>PLoS Genetics</i> , 2012, 8, e1003018.	3.5	25
9	Non-Recognition-of-BTH4, an <i>Arabidopsis</i> Mediator Subunit Homolog, Is Necessary for Development and Response to Salicylic Acid. <i>Plant Cell</i> , 2012, 24, 4220-4235.	6.6	99
10	The BLADE-ON-PETIOLE genes of <i>Arabidopsis</i> are essential for resistance induced by methyl jasmonate. <i>BMC Plant Biology</i> , 2012, 12, 199.	3.6	28
11	Quantitative genetic analysis of salicylic acid perception in <i>Arabidopsis</i> . <i>Planta</i> , 2011, 234, 671-684.	3.2	14
12	The <i>Pseudomonas syringae</i> effector protein HopZ1a suppresses effector-triggered immunity. <i>New Phytologist</i> , 2010, 187, 1018-1033.	7.3	52
13	Structure-function analysis of <i>npr1</i> alleles in <i>Arabidopsis</i> reveals a role for its paralogs in the perception of salicylic acid. <i>Plant, Cell and Environment</i> , 2010, 33, 1911-1922.	5.7	67
14	Resistance and biomass in <i>Arabidopsis</i> : a new model for Salicylic Acid perception. <i>Plant Biotechnology Journal</i> , 2010, 8, 126-141.	8.3	47
15	Specific <i>Arabidopsis</i> HSP90.2 alleles recapitulate RAR1 cochaperone function in plant NB-LRR disease resistance protein regulation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 9556-9563.	7.1	95
16	Identification of new type III effectors and analysis of the plant response by competitive index. <i>Molecular Plant Pathology</i> , 2009, 10, 69-80.	4.2	39
17	Molecular analysis of menadione-induced resistance against biotic stress in <i>Arabidopsis</i> . <i>Plant Biotechnology Journal</i> , 2009, 7, 744-762.	8.3	31
18	Cytosolic HSP90 associates with and modulates the <i>Arabidopsis</i> RPM1 disease resistance protein. <i>EMBO Journal</i> , 2003, 22, 5679-5689.	7.8	365

#	ARTICLE	IF	CITATIONS
19	RAR1 and NDR1 Contribute Quantitatively to Disease Resistance in Arabidopsis, and Their Relative Contributions Are Dependent on the R Gene Assayed. <i>Plant Cell</i> , 2002, 14, 1005-1015.	6.6	218
20	Large-Scale Structure and Function Analysis of the Arabidopsis RPM1 Disease Resistance Protein. <i>Plant Cell</i> , 2002, 14, 435-450.	6.6	141
21	NHL25 and NHL3, Two NDR1/HIN1-Like Genes in Arabidopsis thaliana with Potential Role(s) in Plant Defense. <i>Molecular Plant-Microbe Interactions</i> , 2002, 15, 608-616.	2.6	72
22	A high-throughput method for quantifying growth of phytopathogenic bacteria in Arabidopsis thaliana. <i>Plant Journal</i> , 2002, 28, 475-481.	5.7	127
23	A tomato homeobox gene (HD-Zip) is involved in limiting the spread of programmed cell death. <i>Plant Journal</i> , 1999, 20, 591-600.	5.7	49
24	Identification of a New Pathogen-induced Member of the Subtilisin-like Processing Protease Family from Plants. <i>Journal of Biological Chemistry</i> , 1997, 272, 14412-14419.	3.4	127
25	Two PR-1 Genes from Tomato Are Differentially Regulated and Reveal a Novel Mode of Expression for a Pathogenesis-Related Gene During the Hypersensitive Response and Development. <i>Molecular Plant-Microbe Interactions</i> , 1997, 10, 624-634.	2.6	133
26	Characterization of LRP, a leucine-rich repeat (LRR) protein from tomato plants that is processed during pathogenesis. <i>Plant Journal</i> , 1996, 10, 315-330.	5.7	95
27	Phloem-specific expression of a plant homeobox gene during secondary phases of vascular development. <i>Plant Journal</i> , 1996, 9, 639-648.	5.7	45
28	A gene encoding a novel isoform of the PR-1 protein family from tomato is induced upon viroid infection. <i>Molecular Genetics and Genomics</i> , 1994, 243, 47-53.	2.4	51