

Emilio Daniel Valenzuela Espinoza

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2278628/publications.pdf>

Version: 2024-02-01

10
papers

683
citations

1307594

7
h-index

1372567

10
g-index

10
all docs

10
docs citations

10
times ranked

908
citing authors

#	ARTICLE	IF	CITATIONS
1	Hypoxia-related parameters during septic shock resuscitation: Pathophysiological determinants and potential clinical implications. <i>Annals of Translational Medicine</i> , 2020, 8, 784-784.	1.7	5
2	A lactate-targeted resuscitation strategy may be associated with higher mortality in patients with septic shock and normal capillary refill time: a post hoc analysis of the ANDROMEDA-SHOCK study. <i>Annals of Intensive Care</i> , 2020, 10, 114.	4.6	42
3	Effects of capillary refill time-vs. lactate-targeted fluid resuscitation on regional, microcirculatory and hypoxia-related perfusion parameters in septic shock: a randomized controlled trial. <i>Annals of Intensive Care</i> , 2020, 10, 150.	4.6	34
4	Effects of short-term hyperoxia on systemic hemodynamics, oxygen transport, and microcirculation: An observational study in patients with septic shock and healthy volunteers. <i>Journal of Critical Care</i> , 2019, 53, 62-68.	2.2	11
5	Effect of a Resuscitation Strategy Targeting Peripheral Perfusion Status vs Serum Lactate Levels on 28-Day Mortality Among Patients With Septic Shock. <i>JAMA - Journal of the American Medical Association</i> , 2019, 321, 654.	7.4	471
6	Early rise in central venous pressure during a spontaneous breathing trial: A promising test to identify patients at high risk of weaning failure?. <i>PLoS ONE</i> , 2019, 14, e0225181.	2.5	6
7	Norepinephrine, more than a vasopressor. <i>Annals of Translational Medicine</i> , 2019, 7, S25-S25.	1.7	6
8	Relationship of at Admission Lactate, Unmeasured Anions, and Chloride to the Outcome of Critically Ill Patients. <i>Critical Care Medicine</i> , 2017, 45, e1233-e1239.	0.9	35
9	Prognostic value of ventricular diastolic dysfunction in patients with severe sepsis and septic shock. <i>Revista Brasileira De Terapia Intensiva</i> , 2015, 27, 333-9.	0.3	41
10	Lack of agreement between different observers and methods in the measurement of capillary refill time in healthy volunteers: an observational study. <i>Revista Brasileira De Terapia Intensiva</i> , 2014, 26, 269-76.	0.3	32