

Serge Maurice Mbadinga

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2299682/publications.pdf>

Version: 2024-02-01

50
papers

1,789
citations

361413

20
h-index

276875

41
g-index

50
all docs

50
docs citations

50
times ranked

1926
citing authors

#	ARTICLE	IF	CITATIONS
1	Dominant and Active Methanogens in the Production Waters From a High-Temperature Petroleum Reservoir by DNA- and RNA-Based Analysis. <i>Geomicrobiology Journal</i> , 2021, 38, 191-198.	2.0	4
2	Synthesis and mass spectra of rearrangement bio-signature metabolites of anaerobic alkane degradation via fumarate addition. <i>Analytical Biochemistry</i> , 2020, 600, 113746.	2.4	2
3	Genomic and Transcriptomic Evidence Supports Methane Metabolism in <i>Archaeoglobi</i> . <i>MSystems</i> , 2020, 5, .	3.8	33
4	Methanogenic biodegradation of C13 and C14 n-alkanes activated by addition to fumarate. <i>International Biodeterioration and Biodegradation</i> , 2020, 153, 104994.	3.9	6
5	Long-chain n-alkane biodegradation coupling to methane production in an enriched culture from production water of a high-temperature oil reservoir. <i>AMB Express</i> , 2020, 10, 63.	3.0	13
6	Methanogenic biodegradation of C9 to C12n-alkanes initiated by <i>Smithella</i> via fumarate addition mechanism. <i>AMB Express</i> , 2020, 10, 23.	3.0	22
7	The newly proposed TACK and DPANN archaea detected in the production waters from a high-temperature petroleum reservoir. <i>International Biodeterioration and Biodegradation</i> , 2019, 143, 104729.	3.9	11
8	Methanogenic Degradation of Long <i>n</i> -Alkanes Requires Fumarate-Dependent Activation. <i>Applied and Environmental Microbiology</i> , 2019, 85, .	3.1	22
9	High microbial diversity of the nitric oxide dismutation reaction revealed by PCR amplification and analysis of the nod gene. <i>International Biodeterioration and Biodegradation</i> , 2019, 143, 104708.	3.9	10
10	Bioconversion Pathway of CO ₂ in the Presence of Ethanol by Methanogenic Enrichments from Production Water of a High-Temperature Petroleum Reservoir. <i>Energies</i> , 2019, 12, 918.	3.1	2
11	Direct microbial transformation of carbon dioxide to value-added chemicals: A comprehensive analysis and application potentials. <i>Bioresource Technology</i> , 2019, 288, 121401.	9.6	40
12	Simulation of in situ oil reservoir conditions in a laboratory bioreactor testing for methanogenic conversion of crude oil and analysis of the microbial community. <i>International Biodeterioration and Biodegradation</i> , 2019, 136, 24-33.	3.9	14
13	Methanogenic degradation of branched alkanes in enrichment cultures of production water from a high-temperature petroleum reservoir. <i>Applied Microbiology and Biotechnology</i> , 2019, 103, 2391-2401.	3.6	21
14	Accelerated CO ₂ reduction to methane for energy by zero valent iron in oil reservoir production waters. <i>Energy</i> , 2018, 147, 663-671.	8.8	27
15	Characterization of bacterial composition and diversity in a long-term petroleum contaminated soil and isolation of high-efficiency alkane-degrading strains using an improved medium. <i>World Journal of Microbiology and Biotechnology</i> , 2018, 34, 34.	3.6	48
16	Microbial reduction of CO ₂ from injected NaH ₂ CO ₃ with degradation of n-hexadecane in the enrichment culture derived from a petroleum reservoir. <i>International Biodeterioration and Biodegradation</i> , 2018, 127, 192-200.	3.9	12
17	Metabolic capability and in situ activity of microorganisms in an oil reservoir. <i>Microbiome</i> , 2018, 6, 5.	11.1	70
18	Biodiesel production from waste cooking oil using onsite produced purified lipase from <i>Pseudomonas aeruginosa</i> FW_SH-1: Central composite design approach. <i>Renewable Energy</i> , 2017, 109, 93-100.	8.9	60

#	ARTICLE	IF	CITATIONS
19	Microbiota and their affiliation with physiochemical characteristics of different subsurface petroleum reservoirs. <i>International Biodeterioration and Biodegradation</i> , 2017, 120, 170-185.	3.9	63
20	Propionate metabolism and diversity of relevant functional genes by in silico analysis and detection in subsurface petroleum reservoirs. <i>World Journal of Microbiology and Biotechnology</i> , 2017, 33, 182.	3.6	6
21	Iron oxides alter methanogenic pathways of acetate in production water of high-temperature petroleum reservoir. <i>Applied Microbiology and Biotechnology</i> , 2017, 101, 7053-7063.	3.6	16
22	Type II chaperonin gene as a complementary barcode for 16S rRNA gene in study of Archaea diversity of petroleum reservoirs. <i>International Biodeterioration and Biodegradation</i> , 2017, 123, 113-120.	3.9	8
23	Synthesis and Characterization of Anaerobic Degradation Biomarkers of n-Alkanes via Hydroxylation/Carboxylation Pathways. <i>European Journal of Mass Spectrometry</i> , 2016, 22, 31-37.	1.0	7
24	Microbial communities responsible for fixation of CO ₂ revealed by using mcrA, cbbM, cbbL, fthfs, fefe-hydrogenase genes as molecular biomarkers in petroleum reservoirs of different temperatures. <i>International Biodeterioration and Biodegradation</i> , 2016, 114, 164-175.	3.9	14
25	Activation of CO ₂ -reducing methanogens in oil reservoir after addition of nutrient. <i>Journal of Bioscience and Bioengineering</i> , 2016, 122, 740-747.	2.2	12
26	Molecular diversity of bacterial bamA gene involved in anaerobic degradation of aromatic hydrocarbons in mesophilic petroleum reservoirs. <i>International Biodeterioration and Biodegradation</i> , 2016, 114, 122-128.	3.9	36
27	Diversity and abundance of ammonia-oxidizing bacteria (AOB) revealed by PCR amplification of amoA gene in a polyacrylamide transportation system of an oilfield. <i>International Biodeterioration and Biodegradation</i> , 2016, 115, 110-118.	3.9	3
28	Dominance of <i>Desulfotignum</i> in sulfate-reducing community in high sulfate production-water of high temperature and corrosive petroleum reservoirs. <i>International Biodeterioration and Biodegradation</i> , 2016, 114, 45-56.	3.9	59
29	The biofilm property and its correlation with high-molecular-weight polyacrylamide degradation in a water injection pipeline of Daqing oilfield. <i>Journal of Hazardous Materials</i> , 2016, 304, 388-399.	12.4	45
30	Non-destructive characterization using MCT reveals the composition and distribution of impurities in solar carnallite. <i>RSC Advances</i> , 2015, 5, 16230-16233.	3.6	2
31	Efficiently Applicability of Synthetic Cu-TiO ₂ in Tetrachloroethene, Trichloroethene and 1,1,1-Trichloroethane Removal in Aqueous Phase under VUV Irradiation. <i>Asian Journal of Chemistry</i> , 2015, 27, 60-66.	0.3	0
32	Insights into the Anaerobic Biodegradation Pathway of n-Alkanes in Oil Reservoirs by Detection of Signature Metabolites. <i>Scientific Reports</i> , 2015, 5, 9801.	3.3	78
33	Anaerolineaceae and Methanosaeta turned to be the dominant microorganisms in alkanes-dependent methanogenic culture after long-term of incubation. <i>AMB Express</i> , 2015, 5, 117.	3.0	244
34	Synthesis of 2-[2H]-2-(1-methylalkyl)succinic acids. <i>Chinese Chemical Letters</i> , 2015, 26, 619-622.	9.0	3
35	Chemical Structure, Property and Potential Applications of Biosurfactants Produced by <i>Bacillus subtilis</i> in Petroleum Recovery and Spill Mitigation. <i>International Journal of Molecular Sciences</i> , 2015, 16, 4814-4837.	4.1	119
36	Analysis of Bacterial and Archaeal Communities along a High-Molecular-Weight Polyacrylamide Transportation Pipeline System in an Oil Field. <i>International Journal of Molecular Sciences</i> , 2015, 16, 7445-7461.	4.1	15

#	ARTICLE	IF	CITATIONS
37	Acetoclastic methanogenesis is likely the dominant biochemical pathway of palmitate degradation in the presence of sulfate. <i>Applied Microbiology and Biotechnology</i> , 2015, 99, 7757-7769.	3.6	12
38	Role of reactive oxygen species in the dechlorination of trichloroethene and 1.1.1-trichloroethane in aqueous phase in UV/TiO ₂ systems. <i>Chemical Engineering Science</i> , 2015, 123, 367-375.	3.8	21
39	Functional genes (dsr) approach reveals similar sulphidogenic prokaryotes diversity but different structure in saline waters from corroding high temperature petroleum reservoirs. <i>Applied Microbiology and Biotechnology</i> , 2014, 98, 1871-1882.	3.6	45
40	Enhanced Photocatalytic Activity of TiO ₂ Nanosheets by Doping with Cu for Chlorinated Solvent Pollutants Degradation. <i>Industrial & Engineering Chemistry Research</i> , 2014, 53, 1368-1376.	3.7	45
41	Efficiently Synthetic TiO ₂ Nano-sheets for PCE, TCE, and TCA Degradations in Aqueous Phase Under VUV Irradiation. <i>Water, Air, and Soil Pollution</i> , 2014, 225, 1.	2.4	5
42	Efficient dechlorination of chlorinated solvent pollutants under UV irradiation by using the synthesized TiO ₂ nano-sheets in aqueous phase. <i>Journal of Environmental Sciences</i> , 2014, 26, 1188-1194.	6.1	11
43	Synthesis of Anaerobic Degradation Biomarkers Alkyl-, Aryl- and Cycloalkylsuccinic Acids and Their Mass Spectral Characteristics. <i>European Journal of Mass Spectrometry</i> , 2014, 20, 287-297.	1.0	14
44	Optimization of Surfactin Production by <i>Bacillus subtilis</i> HSO121 through Plackett-Burman and Response Surface Method. <i>Protein and Peptide Letters</i> , 2014, 21, 885-893.	0.9	12
45	Evaluation of microbial community composition in thermophilic methane-producing incubation of production water from a high-temperature oil reservoir. <i>Environmental Technology (United Kingdom)</i> 10.1080/09593333.2014.941010	0.784314	10
46	Methanogenic Microbial Community Composition of Oily Sludge and Its Enrichment Amended with Alkanes Incubated for Over 500 Days. <i>Geomicrobiology Journal</i> , 2012, 29, 716-726.	2.0	27
47	Analyses of n-alkanes degrading community dynamics of a high-temperature methanogenic consortium enriched from production water of a petroleum reservoir by a combination of molecular techniques. <i>Ecotoxicology</i> , 2012, 21, 1680-1691.	2.4	67
48	Analysis of alkane-dependent methanogenic community derived from production water of a high-temperature petroleum reservoir. <i>Applied Microbiology and Biotechnology</i> , 2012, 96, 531-542.	3.6	102
49	Microbial communities involved in anaerobic degradation of alkanes. <i>International Biodeterioration and Biodegradation</i> , 2011, 65, 1-13.	3.9	175
50	Characterization of an alkane-degrading methanogenic enrichment culture from production water of an oil reservoir after 274 days of incubation. <i>International Biodeterioration and Biodegradation</i> , 2011, 65, 444-450.	3.9	93