

# Russell B Foxall

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2308645/publications.pdf>

Version: 2024-02-01

28  
papers

1,048  
citations

471509

17  
h-index

552781

26  
g-index

28  
all docs

28  
docs citations

28  
times ranked

1536  
citing authors

#	ARTICLE	IF	CITATIONS
1	Follicular Helper T Cells Are Major Human Immunodeficiency Virus-2 Reservoirs and Support Productive Infection. <i>Journal of Infectious Diseases</i> , 2020, 221, 122-126.	4.0	11
2	Early ART in Acute HIV-1 Infection: Impact on the B-Cell Compartment. <i>Frontiers in Cellular and Infection Microbiology</i> , 2020, 10, 347.	3.9	5
3	Role of Dendritic Cells in HIV-2 Pathogenesis. , 2018, , 1818-1826.		0
4	miRNA profiling of human naive CD4 T cells links miR-34c-5p to cell activation and HIV replication. <i>EMBO Journal</i> , 2017, 36, 346-360.	7.8	32
5	Human naïve regulatory T-cells feature high steady-state turnover and are maintained by IL-7. <i>Oncotarget</i> , 2016, 7, 12163-12175.	1.8	31
6	Thymic HIV-2 Infection Uncovers Posttranscriptional Control of Viral Replication in Human Thymocytes. <i>Journal of Virology</i> , 2015, 89, 2201-2208.	3.4	13
7	Enteric Mucosa Integrity in the Presence of a Preserved Innate Interleukin 22 Compartment in HIV Type 1-Treated Individuals. <i>Journal of Infectious Diseases</i> , 2014, 210, 630-640.	4.0	35
8	Monocyte and Myeloid Dendritic Cell Activation Occurs Throughout HIV Type 2 Infection, an Attenuated Form of HIV Disease. <i>Journal of Infectious Diseases</i> , 2013, 207, 1730-1742.	4.0	22
9	Preserved CD4 T-cell telomere length during long-lasting HIV-2 infection. <i>Aids</i> , 2013, 27, 289-292.	2.2	3
10	Role of Dendritic Cells in HIV-2 Pathogenesis. , 2013, , 1-10.		0
11	Memory B-cell depletion is a feature of HIV-2 infection even in the absence of detectable viremia. <i>Aids</i> , 2012, 26, 1607-1617.	2.2	13
12	PD-1 and its ligand PD-L1 are progressively up-regulated on CD4 and CD8 T-cells in HIV-2 infection irrespective of the presence of viremia. <i>Aids</i> , 2012, 26, 1065-1071.	2.2	20
13	Memory and naive-like regulatory CD4+ T cells expand during HIV-2 infection in direct association with CD4+ T-cell depletion irrespective of viremia. <i>Aids</i> , 2011, 25, 1961-1970.	2.2	19
14	Cell-Associated Viral Burden Provides Evidence of Ongoing Viral Replication in Aviremic HIV-2-Infected Patients. <i>Journal of Virology</i> , 2011, 85, 2429-2438.	3.4	50
15	Dendritic Cell Differentiation and Maturation in the Presence of HIV Type 2 Envelope. <i>AIDS Research and Human Retroviruses</i> , 2009, 25, 425-431.	1.1	9
16	Major Depletion of Plasmacytoid Dendritic Cells in HIV-2 Infection, an Attenuated Form of HIV Disease. <i>PLoS Pathogens</i> , 2009, 5, e1000667.	4.7	35
17	Increased frequency of CD25dimCD4+ T-cells in HIV-2 infection, a naturally occurring attenuated form of HIV-1. <i>Clinical Immunology</i> , 2008, 127, 158-167.	3.2	5
18	Gag-Specific CD4+ T-Cell Frequency Is Inversely Correlated with Proviral Load and Directly Correlated with Immune Activation in Infection with Human Immunodeficiency Virus Type 2 (HIV-2) but Not HIV-1. <i>Journal of Virology</i> , 2008, 82, 9795-9799.	3.4	18

#	ARTICLE	IF	CITATIONS
19	Rate of Increase in Circulating IL-7 and Loss of IL-7R $\alpha$ Expression Differ in HIV-1 and HIV-2 Infections: Two Lymphopenic Diseases with Similar Hyperimmune Activation but Distinct Outcomes. <i>Journal of Immunology</i> , 2007, 178, 3252-3259.	0.8	31
20	Low CD4 T-cell counts despite low levels of circulating HIV: Insights from the comparison of HIV-1 infected patients with a discordant response to antiretroviral therapy to patients with untreated advanced HIV-2 disease. <i>Clinical Immunology</i> , 2007, 125, 67-75.	3.2	9
21	Increased Frequency of Circulating CCR5 + CD4 + T Cells in Human Immunodeficiency Virus Type 2 Infection. <i>Journal of Virology</i> , 2006, 80, 12425-12429.	3.4	30
22	Heterologous cells cooperate to augment stem cell migration, homing, and engraftment. <i>Blood</i> , 2003, 101, 45-51.	1.4	46
23	Thymocyte emigration is mediated by active movement away from stroma-derived factors. <i>Journal of Clinical Investigation</i> , 2002, 109, 1101-1110.	8.2	86
24	Thymocyte emigration is mediated by active movement away from stroma-derived factors. <i>Journal of Clinical Investigation</i> , 2002, 109, 1101-1110.	8.2	43
25	Active movement of T cells away from a chemokine. <i>Nature Medicine</i> , 2000, 6, 543-548.	30.7	283
26	Pulsed electric fields for selection of hematopoietic cells and depletion of tumor cell contaminants. <i>Nature Biotechnology</i> , 2000, 18, 882-887.	17.5	36
27	Efficient generation of human T cells from a tissue-engineered thymic organoid. <i>Nature Biotechnology</i> , 2000, 18, 729-734.	17.5	156
28	The Rabbit Study: Zidovudine and Zalcitabine in Combination in Zidovudine-Experienced and Previously Untreated Patients. <i>AIDS Research and Human Retroviruses</i> , 1999, 15, 1181-1189.	1.1	7