

Michiel Albertus Basson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2315267/publications.pdf>

Version: 2024-02-01

70
papers

7,259
citations

117625

34
h-index

85541

71
g-index

76
all docs

76
docs citations

76
times ranked

11774
citing authors

#	ARTICLE	IF	CITATIONS
1	Distinct, dosage-sensitive requirements for the autism-associated factor CHD8 during cortical development. <i>Molecular Autism</i> , 2021, 12, 16.	4.9	15
2	Cell-type-specific synaptic imbalance and disrupted homeostatic plasticity in cortical circuits of ASD-associated Chd8 haploinsufficient mice. <i>Molecular Psychiatry</i> , 2021, 26, 3614-3624.	7.9	18
3	Inositol treatment inhibits medulloblastoma through suppression of epigenetic-driven metabolic adaptation. <i>Nature Communications</i> , 2021, 12, 2148.	12.8	20
4	Effects of Low-Dose Gestational TCDD Exposure on Behavior and on Hippocampal Neuron Morphology and Gene Expression in Mice. <i>Environmental Health Perspectives</i> , 2021, 129, 57002.	6.0	11
5	<scp><i>ZMYND11</i></scp> variants are a novel cause of centrotemporal and generalised epilepsies with neurodevelopmental disorder. <i>Clinical Genetics</i> , 2021, 100, 412-429.	2.0	5
6	Brain mapping across 16 autism mouse models reveals a spectrum of functional connectivity subtypes. <i>Molecular Psychiatry</i> , 2021, 26, 7610-7620.	7.9	47
7	The chromatin remodelling factor Chd7 protects auditory neurons and sensory hair cells from stress-induced degeneration. <i>Communications Biology</i> , 2021, 4, 1260.	4.4	10
8	A recessive PRDM13 mutation results in congenital hypogonadotropic hypogonadism and cerebellar hypoplasia. <i>Journal of Clinical Investigation</i> , 2021, 131, .	8.2	16
9	Regulation of autism-relevant behaviors by cerebellarâ€“prefrontal cortical circuits. <i>Nature Neuroscience</i> , 2020, 23, 1102-1110.	14.8	149
10	The AHR pathway represses TGFÎ²-SMAD3 signalling and has a potent tumour suppressive role in SHH medulloblastoma. <i>Scientific Reports</i> , 2020, 10, 148.	3.3	22
11	Infraslow State Fluctuations Govern Spontaneous fMRI Network Dynamics. <i>Current Biology</i> , 2019, 29, 2295-2306.e5.	3.9	107
12	Sprouty1 Controls Genitourinary Development via its N-Terminal Tyrosine. <i>Journal of the American Society of Nephrology: JASN</i> , 2019, 30, 1398-1411.	6.1	5
13	Altered Neocortical Gene Expression, Brain Overgrowth and Functional Over-Connectivity in Chd8 Haploinsufficient Mice. <i>Cerebral Cortex</i> , 2018, 28, 2192-2206.	2.9	118
14	Autismâ€“linked <scp>CHD</scp> gene expression patterns during development predict multiâ€“organ disease phenotypes. <i>Journal of Anatomy</i> , 2018, 233, 755-769.	1.5	14
15	Sex bias in autism: new insights from Chd8 mutant mice?. <i>Nature Neuroscience</i> , 2018, 21, 1144-1146.	14.8	7
16	Advanced paternal age effects in neurodevelopmental disordersâ€“review of potential underlying mechanisms. <i>Translational Psychiatry</i> , 2017, 7, e1019-e1019.	4.8	94
17	Engrailed controls epaxial-hypaxial muscle innervation and the establishment of vertebrate three-dimensional mobility. <i>Developmental Biology</i> , 2017, 430, 90-104.	2.0	7
18	Convergence of BMI1 and CHD7 on ERK Signaling in Medulloblastoma. <i>Cell Reports</i> , 2017, 21, 2772-2784.	6.4	31

#	ARTICLE	IF	CITATIONS
19	The neuroanatomy of autism – a developmental perspective. <i>Journal of Anatomy</i> , 2017, 230, 4-15.	1.5	156
20	Distinct cerebellar foliation anomalies in a <i>CHD7</i> haploinsufficient mouse model of CHARGE syndrome. <i>American Journal of Medical Genetics, Part C: Seminars in Medical Genetics</i> , 2017, 175, .	1.6	19
21	Cerebellar Vermis and Midbrain Hypoplasia Upon Conditional Deletion of <i>Chd7</i> from the Embryonic Mid-Hindbrain Region. <i>Frontiers in Neuroanatomy</i> , 2017, 11, 86.	1.7	7
22	The chromatin remodeling factor CHD7 controls cerebellar development by regulating reelin expression. <i>Journal of Clinical Investigation</i> , 2017, 127, 874-887.	8.2	61
23	An <i>FGFR1-SPRY2</i> Signaling Axis Limits Basal Cell Proliferation in the Steady-State Airway Epithelium. <i>Developmental Cell</i> , 2016, 37, 85-97.	7.0	24
24	Functional Insights into Chromatin Remodelling from Studies on CHARGE Syndrome. <i>Trends in Genetics</i> , 2015, 31, 600-611.	6.7	66
25	<i>CHD7</i> Maintains Neural Stem Cell Quiescence and Prevents Premature Stem Cell Depletion in the Adult Hippocampus. <i>Stem Cells</i> , 2015, 33, 196-210.	3.2	74
26	Heparan Sulfotransferases <i>Hs6st1</i> and <i>Hs2st</i> Keep Erk in Check for Mouse Corpus Callosum Development. <i>Journal of Neuroscience</i> , 2014, 34, 2389-2401.	3.6	30
27	Epistatic interactions between <i>Chd7</i> and <i>Fgf8</i> during cerebellar development. <i>Rare Diseases (Austin, Tex)</i> , 2014, 2, e28688.	1.8	11
28	<i>Gli3</i> Controls Corpus Callosum Formation by Positioning Midline Guideposts During Telencephalic Patterning. <i>Cerebral Cortex</i> , 2014, 24, 186-198.	2.9	33
29	Coordinated activity of <i>Spry1</i> and <i>Spry2</i> is required for normal development of the external genitalia. <i>Developmental Biology</i> , 2014, 386, 1-11.	2.0	27
30	Endoderm-specific deletion of <i>Tbx1</i> reveals an FGF-independent role for <i>Tbx1</i> in pharyngeal apparatus morphogenesis. <i>Developmental Dynamics</i> , 2014, 243, 1143-1151.	1.8	24
31	A bi-modal function of Wnt signalling directs an FGF activity gradient to spatially regulate neuronal differentiation in the midbrain. <i>Development (Cambridge)</i> , 2014, 141, 63-72.	2.5	30
32	Congenital hypoplasia of the cerebellum: developmental causes and behavioral consequences. <i>Frontiers in Neuroanatomy</i> , 2013, 7, 29.	1.7	80
33	Deregulated FGF and homeotic gene expression underlies cerebellar vermis hypoplasia in CHARGE syndrome. <i>ELife</i> , 2013, 2, e01305.	6.0	55
34	<i>Sprouty1</i> Haploinsufficiency Prevents Renal Agenesis in a Model of Fraser Syndrome. <i>Journal of the American Society of Nephrology: JASN</i> , 2012, 23, 1790-1796.	6.1	24
35	Localised inhibition of FGF signalling in the third pharyngeal pouch is required for normal thymus and parathyroid organogenesis. <i>Development (Cambridge)</i> , 2012, 139, 3456-3466.	2.5	30
36	<i>Sprouty1</i> is a candidate tumor-suppressor gene in medullary thyroid carcinoma. <i>Oncogene</i> , 2012, 31, 3961-3972.	5.9	31

#	ARTICLE	IF	CITATIONS
37	The aged niche disrupts muscle stem cell quiescence. <i>Nature</i> , 2012, 490, 355-360.	27.8	682
38	Periodic stripe formation by a Turing mechanism operating at growth zones in the mammalian palate. <i>Nature Genetics</i> , 2012, 44, 348-351.	21.4	214
39	Sprouty Is a Negative Regulator of Transforming Growth Factor β^2 -Induced Epithelial-to-Mesenchymal Transition and Cataract. <i>Molecular Medicine</i> , 2012, 18, 861-873.	4.4	49
40	Signaling in Cell Differentiation and Morphogenesis. <i>Cold Spring Harbor Perspectives in Biology</i> , 2012, 4, a008151-a008151.	5.5	121
41	Biallelic expression of <i>Tbx1</i> protects the embryo from developmental defects caused by increased receptor tyrosine kinase signaling. <i>Developmental Dynamics</i> , 2012, 241, 1310-1324.	1.8	9
42	Regulation of CD4+ and CD8+ Effector Responses by Sprouty-1. <i>PLoS ONE</i> , 2012, 7, e49801.	2.5	16
43	Expression of fibroblast growth factors (Fgfs) in murine tooth development. <i>Journal of Anatomy</i> , 2011, 218, 534-543.	1.5	37
44	Sprouty genes are essential for the normal development of epibranchial ganglia in the mouse embryo. <i>Developmental Biology</i> , 2011, 358, 147-155.	2.0	16
45	Sprouty genes prevent excessive FGF signalling in multiple cell types throughout development of the cerebellum. <i>Development (Cambridge)</i> , 2011, 138, 2957-2968.	2.5	53
46	Characterization of a <i>Dchs1</i> mutant mouse reveals requirements for Dchs1-Fat4 signaling during mammalian development. <i>Development (Cambridge)</i> , 2011, 138, 947-957.	2.5	172
47	Sprouty Proteins Inhibit Receptor-mediated Activation of Phosphatidylinositol-specific Phospholipase C. <i>Molecular Biology of the Cell</i> , 2010, 21, 3487-3496.	2.1	45
48	Sprouty1 Regulates Reversible Quiescence of a Self-Renewing Adult Muscle Stem Cell Pool during Regeneration. <i>Cell Stem Cell</i> , 2010, 6, 117-129.	11.1	275
49	FGF ligands emerge as potential specifiers of synaptic identity. <i>Cellscience</i> , 2010, 7, 33-42.	0.3	4
50	Great vessel development requires biallelic expression of Chd7 and Tbx1 in pharyngeal ectoderm in mice. <i>Journal of Clinical Investigation</i> , 2009, 119, 3301-10.	8.2	119
51	Tbx1 controls cardiac neural crest cell migration during arch artery development by regulating <i>Gbx2</i> expression in the pharyngeal ectoderm. <i>Development (Cambridge)</i> , 2009, 136, 3173-3183.	2.5	124
52	Loss of Sprouty1 Rescues Renal Agenesis Caused by Ret Mutation. <i>Journal of the American Society of Nephrology: JASN</i> , 2009, 20, 255-259.	6.1	45
53	Specific regions within the embryonic midbrain and cerebellum require different levels of FGF signaling during development. <i>Development (Cambridge)</i> , 2009, 136, 1962-1962.	2.5	1
54	Fibroblast growth factor (FGF) gene expression in the developing cerebellum suggests multiple roles for FGF signaling during cerebellar morphogenesis and development. <i>Developmental Dynamics</i> , 2009, 238, 2058-2072.	1.8	46

#	ARTICLE	IF	CITATIONS
55	MicroRNA-21 contributes to myocardial disease by stimulating MAP kinase signalling in fibroblasts. <i>Nature</i> , 2008, 456, 980-984.	27.8	2,111
56	An FGF signaling loop sustains the generation of differentiated progeny from stem cells in mouse incisors. <i>Development (Cambridge)</i> , 2008, 135, 377-385.	2.5	150
57	Specific regions within the embryonic midbrain and cerebellum require different levels of FGF signaling during development. <i>Development (Cambridge)</i> , 2008, 135, 889-898.	2.5	124
58	Itch ^Δ and Iti ^Δ T cells independently contribute to autoimmunity in Itchy mice. <i>Blood</i> , 2008, 111, 4273-7282	1.4	42
59	Branching morphogenesis of the ureteric epithelium during kidney development is coordinated by the opposing functions of GDNF and Sprouty1. <i>Developmental Biology</i> , 2006, 299, 466-477.	2.0	141
60	Sprouty proteins: multifaceted negative-feedback regulators of receptor tyrosine kinase signaling. <i>Trends in Cell Biology</i> , 2006, 16, 45-54.	7.9	408
61	Synergistic activity of Sef and Sprouty proteins in regulating the expression of Gbx2 in the mid-hindbrain region. <i>Genesis</i> , 2005, 41, 110-115.	1.6	36
62	Sprouty1 Is a Critical Regulator of GDNF/RET-Mediated Kidney Induction. <i>Developmental Cell</i> , 2005, 8, 229-239.	7.0	327
63	The influence of the src family kinases, Lck and Fyn, on T cell differentiation, survival and activation. <i>Immunological Reviews</i> , 2003, 191, 107-118.	6.0	178
64	Insights into T-Cell Development from Studies Using Transgenic and Knockout Mice. <i>Molecular Biotechnology</i> , 2001, 18, 11-24.	2.4	3
65	The CD4/CD8 lineage decision: integration of signalling pathways. <i>Trends in Immunology</i> , 2000, 21, 509-514.	7.5	44
66	Early Growth Response (Egr)-1 Gene Induction in the Thymus in Response to TCR Ligation During Early Steps in Positive Selection Is Not Required for CD8 Lineage Commitment. <i>Journal of Immunology</i> , 2000, 165, 2444-2450.	0.8	22
67	Greatly reduced efficiency of both positive and negative selection of thymocytes in CD45 tyrosine phosphatase-deficient mice. <i>European Journal of Immunology</i> , 1999, 29, 2923-2933.	2.9	67
68	Greatly reduced efficiency of both positive and negative selection of thymocytes in CD45 tyrosine phosphatase-deficient mice. <i>European Journal of Immunology</i> , 1999, 29, 2923-2933.	2.9	2
69	Molecular requirements for lineage commitment in the thymus - antibody-mediated receptor engagements reveal a central role for Lck in lineage decisions. <i>Immunological Reviews</i> , 1998, 165, 181-194.	6.0	28
70	CD3 Ligation on Immature Thymocytes Generates Antagonist-like Signals Appropriate for CD8 Lineage Commitment, Independently of T Cell Receptor Specificity. <i>Journal of Experimental Medicine</i> , 1998, 187, 1249-1260.	8.5	58