

# Tatjana Stojakovic

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2351361/publications.pdf>

Version: 2024-02-01

95  
papers

3,961  
citations

117625

34  
h-index

133252

59  
g-index

96  
all docs

96  
docs citations

96  
times ranked

8774  
citing authors

#	ARTICLE	IF	CITATIONS
1	Open-label phase III study evaluating safety and efficacy of the non-steroidal farnesoid X receptor agonist PX-104 in non-alcoholic fatty liver disease. <i>Wiener Klinische Wochenschrift</i> , 2021, 133, 441-451.	1.9	27
2	Hypoglycaemia leads to a delayed increase in platelet and coagulation activation markers in people with type 2 diabetes treated with metformin only: Results from a stepwise hypoglycaemic clamp study. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 212-221.	4.4	12
3	E2F-Family Members Engage the PIDDosome to Limit Hepatocyte Ploidy in Liver Development and Regeneration. <i>Developmental Cell</i> , 2020, 52, 335-349.e7.	7.0	40
4	A low-fat spread with added plant sterols and fish omega-3 fatty acids lowers serum triglyceride and LDL-cholesterol concentrations in individuals with modest hypercholesterolaemia and hypertriglyceridaemia. <i>European Journal of Nutrition</i> , 2019, 58, 1615-1624.	3.9	23
5	Comparison of lipoprotein (a) serum concentrations measured by six commercially available immunoassays. <i>Atherosclerosis</i> , 2019, 289, 206-213.	0.8	66
6	HDL subclasses and mortality in acute heart failure patients. <i>Clinica Chimica Acta</i> , 2019, 490, 81-87.	1.1	13
7	LDL triglycerides, hepatic lipase activity, and coronary artery disease: An epidemiologic and Mendelian randomization study. <i>Atherosclerosis</i> , 2019, 282, 37-44.	0.8	38
8	Soluble urokinase plasminogen activation receptor and long-term outcomes in persons undergoing coronary angiography. <i>Scientific Reports</i> , 2019, 9, 475.	3.3	8
9	Serum concentrations of free fatty acids are associated with 3-month mortality in acute heart failure patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2019, 57, 1799-1804.	2.3	6
10	Treatment with PCSK9 inhibitors reduces atherogenic VLDL remnants in a real-world study. <i>Vascular Pharmacology</i> , 2019, 116, 8-15.	2.1	20
11	High Keratin 8/18 Ratio Predicts Aggressive Hepatocellular Cancer Phenotype. <i>Translational Oncology</i> , 2019, 12, 256-268.	3.7	28
12	Hepatocyte-specific lysosomal acid lipase deficiency protects mice from diet-induced obesity but promotes hepatic inflammation. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2019, 1864, 500-511.	2.4	25
13	Large platelet size is associated with poor outcome in patients with metastatic pancreatic cancer. <i>Clinical Chemistry and Laboratory Medicine</i> , 2019, 57, 740-744.	2.3	8
14	The interrelations between PCSK9 metabolism and cholesterol synthesis and absorption. <i>Journal of Lipid Research</i> , 2019, 60, 161-167.	4.2	16
15	Renal function, N-terminal Pro-B-Type natriuretic peptide, propeptide big-endothelin and patients with heart failure and preserved ejection fraction. <i>Peptides</i> , 2019, 111, 112-117.	2.4	8
16	Prospective cohort studies of beta-trace protein and mortality in haemodialysis patients and patients undergoing coronary angiography. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, 1984-1991.	0.7	3
17	Colesevelam attenuates cholestatic liver and bile duct injury in <i>Mdr2<sup>Δ/Δ</sup></i> mice by modulating composition, signalling and excretion of faecal bile acids. <i>Gut</i> , 2018, 67, 1683-1691.	12.1	53
18	Lysyl oxidase-like protein 2 (LOXL2) modulates barrier function in cholangiocytes in cholestasis. <i>Journal of Hepatology</i> , 2018, 69, 368-377.	3.7	27

#	ARTICLE	IF	CITATIONS
19	Vitamin D supplementation and lipoprotein metabolism: A randomized controlled trial. <i>Journal of Clinical Lipidology</i> , 2018, 12, 588-596.e4.	1.5	36
20	Elevated Cardiac Troponin T in Patients With Skeletal Myopathies. <i>Journal of the American College of Cardiology</i> , 2018, 71, 1540-1549.	2.8	150
21	Combined serum free light chain levels are associated with carotid atherosclerosis in type 2 diabetes mellitus. <i>Diabetes and Vascular Disease Research</i> , 2018, 15, 162-164.	2.0	3
22	Hepatocyte specific expression of an oncogenic variant of $\beta$ -catenin results in lethal metabolic dysfunction in mice. <i>Oncotarget</i> , 2018, 9, 11243-11257.	1.8	6
23	Effects of linagliptin on endothelial function and postprandial lipids in coronary artery disease patients with early diabetes: a randomized, placebo-controlled, double-blind trial. <i>Cardiovascular Diabetology</i> , 2018, 17, 71.	6.8	13
24	Elevated amylase in plasma represents an adverse prognostic marker in patients with metastatic pancreatic cancer. <i>Wiener Klinische Wochenschrift</i> , 2018, 130, 569-574.	1.9	4
25	Beta-trace Protein as a new non-invasive immunological Marker for Quinolinic Acid-induced impaired Blood-Brain Barrier Integrity. <i>Scientific Reports</i> , 2017, 7, 43642.	3.3	18
26	Circulating proprotein convertase subtilisin-kexin type 9, all-cause mortality, and cardiovascular mortality: The Ludwigshafen Risk and Cardiovascular Health study. <i>European Journal of Preventive Cardiology</i> , 2017, 24, 1095-1101.	1.8	7
27	Critical evaluation of the potential prognostic value of the pretreatment-derived neutrophil-lymphocyte ratio under consideration of C-reactive protein levels in clear cell renal cell carcinoma. <i>British Journal of Cancer</i> , 2017, 116, 85-90.	6.4	39
28	Prognostic value of free light chains lambda and kappa in early multiple sclerosis. <i>Multiple Sclerosis Journal</i> , 2017, 23, 1496-1505.	3.0	34
29	Relations between lipoprotein(a) concentrations, LPA genetic variants, and the risk of mortality in patients with established coronary heart disease: a molecular and genetic association study. <i>Lancet Diabetes and Endocrinology</i> , 2017, 5, 534-543.	11.4	84
30	Lipidomics by ultrahigh performance liquid chromatography-high resolution mass spectrometry and its application to complex biological samples. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2017, 1053, 72-80.	2.3	87
31	Metabolic preconditioning protects BSEP/ABCB11 mice against cholestatic liver injury. <i>Journal of Hepatology</i> , 2017, 66, 95-101.	3.7	51
32	Impact of Endothelial Lipase on Cholesterol Efflux Capacity of Serum and High-density Lipoprotein. <i>Scientific Reports</i> , 2017, 7, 12485.	3.3	19
33	<sc>FXR</sc> controls <sc>CHOP</sc> expression in steatohepatitis. <i>FEBS Letters</i> , 2017, 591, 3360-3368.	2.8	15
34	ST2 predicts survival in patients undergoing transcatheter aortic valve implantation. <i>International Journal of Cardiology</i> , 2017, 244, 87-92.	1.7	17
35	Propeptide big-endothelin, N-terminal-pro brain natriuretic peptide and mortality. The Ludwigshafen risk and cardiovascular health (LURIC) study. <i>Biomarkers</i> , 2017, 22, 315-320.	1.9	3
36	Bile acid preparation and comprehensive analysis by high performance liquid chromatography-high-resolution mass spectrometry. <i>Clinica Chimica Acta</i> , 2017, 464, 85-92.	1.1	47

#	ARTICLE	IF	CITATIONS
37	Combined Respiratory Chain Deficiency and UQCRC2 Mutations in Neonatal Encephalomyopathy: Defective Supercomplex Assembly in Complex III Deficiencies. <i>Oxidative Medicine and Cellular Longevity</i> , 2017, 2017, 1-11.	4.0	33
38	A New Murine Model of Chronic Kidney Disease-Mineral and Bone Disorder. <i>International Journal of Endocrinology</i> , 2017, 2017, 1-8.	1.5	4
39	Neutrophil gelatinase-associated lipocalin levels are U-shaped in the Ludwigshafen Risk and Cardiovascular Health (LURIC) study—Impact for mortality. <i>PLoS ONE</i> , 2017, 12, e0171574.	2.5	14
40	Effects of Vitamin D Supplementation on Serum 25-Hydroxyvitamin D Concentrations in Cirrhotic Patients: A Randomized Controlled Trial. <i>Nutrients</i> , 2016, 8, 278.	4.1	19
41	CHA2DS2-VASc score and blood biomarkers to identify patients with atrial high-rate episodes and paroxysmal atrial fibrillation. <i>Europace</i> , 2016, 19, euw101.	1.7	18
42	Plasma ceramides predict cardiovascular death in patients with stable coronary artery disease and acute coronary syndromes beyond LDL-cholesterol. <i>European Heart Journal</i> , 2016, 37, 1967-1976.	2.2	433
43	Novel pathomechanisms of cardiomyocyte dysfunction in a model of heart failure with preserved ejection fraction. <i>European Journal of Heart Failure</i> , 2016, 18, 987-997.	7.1	53
44	Prognostic Utility of Galectin-3 for Recurrent Cardiovascular Events During Long-term Follow-up in Patients with Stable Coronary Heart Disease: Results of the KAROLA Study. <i>Clinical Chemistry</i> , 2016, 62, 1372-1379.	3.2	17
45	The elevated preoperative derived neutrophil-to-lymphocyte ratio predicts poor clinical outcome in breast cancer patients. <i>Tumor Biology</i> , 2016, 37, 361-368.	1.8	39
46	Simvastatin Efficiently Lowers Small LDL-IgG Immune Complex Levels: A Therapeutic Quality beyond the Lipid-Lowering Effect. <i>PLoS ONE</i> , 2016, 11, e0148210.	2.5	16
47	Blood-Based Biomarkers Are Associated with Disease Recurrence and Survival in Gastrointestinal Stroma Tumor Patients after Surgical Resection. <i>PLoS ONE</i> , 2016, 11, e0159448.	2.5	30
48	Hepatocyte specific expression of an oncogenic variant of $\beta$ -catenin results in cholestatic liver disease. <i>Oncotarget</i> , 2016, 7, 86985-86998.	1.8	13
49	Acute Physiological Responses to Short- and Long-Stage High-Intensity Interval Exercise in Cardiac Rehabilitation: A Pilot Study. <i>Journal of Sports Science and Medicine</i> , 2016, 15, 80-91.	1.6	13
50	The chronic kidney disease epidemiology collaboration equation combining creatinine and cystatin C accurately assesses renal function in patients with cirrhosis. <i>BMC Nephrology</i> , 2015, 16, 196.	1.8	30
51	<i>Lactobacillus casei</i> Shirota Supplementation Does Not Restore Gut Microbiota Composition and Gut Barrier in Metabolic Syndrome: A Randomized Pilot Study. <i>PLoS ONE</i> , 2015, 10, e0141399.	2.5	45
52	Ratio of Apolipoprotein A-II/B Improves Risk Prediction of Postoperative Survival After Carotid Endarterectomy. <i>Stroke</i> , 2015, 46, 1700-1703.	2.0	6
53	Effect of High-Flux Dialysis on Circulating FGF-23 Levels in End-Stage Renal Disease Patients: Results from a Randomized Trial. <i>PLoS ONE</i> , 2015, 10, e0128079.	2.5	7
54	Dialysis Modalities and HDL Composition and Function. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 2267-2276.	6.1	73

#	ARTICLE	IF	CITATIONS
55	Preoperative serum-gamma-glutamyltransferase (GGT) does not represent an independent prognostic factor in a European cohort of patients with non-metastatic renal cell carcinoma. <i>Journal of Clinical Pathology</i> , 2015, 68, 547-551.	2.0	13
56	Validation of the neutrophil-to-lymphocyte ratio as a prognostic factor in a cohort of European prostate cancer patients. <i>World Journal of Urology</i> , 2015, 33, 1661-1667.	2.2	43
57	Ursodeoxycholic acid exerts farnesoid X receptor-antagonistic effects on bile acid and lipid metabolism in morbid obesity. <i>Journal of Hepatology</i> , 2015, 62, 1398-1404.	3.7	236
58	Cognitive Avoidant Coping Is Associated with Higher Carotid Intima Media Thickness Among Middle-Aged Adults. <i>International Journal of Behavioral Medicine</i> , 2015, 22, 597-604.	1.7	1
59	The derived neutrophil/lymphocyte ratio predicts poor clinical outcome in soft tissue sarcoma patients. <i>American Journal of Surgery</i> , 2015, 210, 111-116.	1.8	34
60	Effect of chronic kidney disease on macrophage cholesterol efflux. <i>Life Sciences</i> , 2015, 136, 1-6.	4.3	19
61	Evaluation of the platelet-to-lymphocyte ratio as a prognostic indicator in a European cohort of patients with prostate cancer treated with radiotherapy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015, 33, 201.e9-201.e16.	1.6	25
62	Pretreatment lymphocyte/monocyte ratio as a potential prognostic factor in a cohort of patients with upper tract urothelial carcinoma. <i>Journal of Clinical Pathology</i> , 2015, 68, 351-355.	2.0	38
63	Oxidized LDL Is Strictly Limited to Hyperthyroidism Irrespective of Fat Feeding in Female Sprague Dawley Rats. <i>International Journal of Molecular Sciences</i> , 2015, 16, 11689-11698.	4.1	2
64	The Preoperative AST/ALT (De Ritis) Ratio Represents a Poor Prognostic Factor in a Cohort of Patients with Nonmetastatic Renal Cell Carcinoma. <i>Journal of Urology</i> , 2015, 194, 30-35.	0.4	105
65	The elevated C-reactive protein level is associated with poor prognosis in prostate cancer patients treated with radiotherapy. <i>European Journal of Cancer</i> , 2015, 51, 610-619.	2.8	49
66	Profound Changes in Sex Hormone Levels during Cross-Sex Hormone Therapy of Transsexuals do not Alter Serum Cholesterol Acceptor Capacity. <i>Journal of Sexual Medicine</i> , 2015, 12, 1436-1439.	0.6	4
67	Soluble klotho and mortality: The Ludwigshafen Risk and Cardiovascular Health Study. <i>Atherosclerosis</i> , 2015, 242, 483-489.	0.8	38
68	An elevated preoperative plasma fibrinogen level is associated with poor disease-specific and overall survival in breast cancer patients. <i>Breast</i> , 2015, 24, 667-672.	2.2	31
69	The association of an elevated plasma fibrinogen level with cancer-specific and overall survival in prostate cancer patients. <i>World Journal of Urology</i> , 2015, 33, 1467-1473.	2.2	27
70	Mid-regional proadrenomedullin levels predict recurrence of atrial fibrillation after catheter ablation. <i>International Journal of Cardiology</i> , 2015, 180, 129-133.	1.7	8
71	HDL Cholesterol, Apolipoproteins, and Cardiovascular Risk in Hemodialysis Patients. <i>Journal of the American Society of Nephrology: JASN</i> , 2015, 26, 484-492.	6.1	37
72	Prognostic relevance of ischemia-modified albumin and NT-proBNP in patients with peripheral arterial occlusive disease. <i>Clinica Chimica Acta</i> , 2015, 438, 255-260.	1.1	12

#	ARTICLE	IF	CITATIONS
73	Serum Bile Acids in Repaired Tetralogy of Fallot: A Marker for Liver and Heart?. PLoS ONE, 2015, 10, e0144745.	2.5	5
74	Preoperative Leucocytosis, Thrombocytosis and Anemia as Potential Prognostic Factors in Non-metastatic Renal Cell Carcinoma. Anticancer Research, 2015, 35, 3463-9.	1.1	17
75	Evaluation of Uric Acid as a Prognostic Blood-Based Marker in a Large Cohort of Pancreatic Cancer Patients. PLoS ONE, 2014, 9, e104730.	2.5	39
76	Pre-Treatment Anemia Is a Poor Prognostic Factor in Soft Tissue Sarcoma Patients. PLoS ONE, 2014, 9, e107297.	2.5	31
77	Editorial for "Randomized controlled trial assessing the effect of simvastatin in primary biliary cirrhosis". Liver International, 2014, 34, 328-329.	3.9	1
78	The Arachidonic Acid Metabolome Serves as a Conserved Regulator of Cholesterol Metabolism. Cell Metabolism, 2014, 20, 787-798.	16.2	92
79	Effect of High-Dose Vitamin D <sub>3</sub> on Hospital Length of Stay in Critically Ill Patients With Vitamin D Deficiency. JAMA - Journal of the American Medical Association, 2014, 312, 1520.	7.4	368
80	The elevated preoperative platelet to lymphocyte ratio predicts decreased time to recurrence in colon cancer patients. American Journal of Surgery, 2014, 208, 210-214.	1.8	97
81	Cardiovascular Function and Predictors of Exercise Capacity in Patients With Colorectal Cancer. Journal of the American College of Cardiology, 2014, 64, 1310-1319.	2.8	93
82	Gestational diabetes mellitus modulates neonatal high-density lipoprotein composition and its functional heterogeneity. Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids, 2014, 1841, 1619-1627.	2.4	35
83	Low preoperative lymphocyte-monocyte ratio (LMR) represents a potentially poor prognostic factor in nonmetastatic clear cell renal cell carcinoma. Urologic Oncology: Seminars and Original Investigations, 2014, 32, 1041-1048.	1.6	82
84	Adipocyte cell size, free fatty acids and apolipoproteins are associated with non-alcoholic liver injury progression in severely obese patients. Metabolism: Clinical and Experimental, 2014, 63, 1542-1552.	3.4	88
85	Fibroblast growth factor 23 (FGF23) and mortality: The Ludwigshafen Risk and Cardiovascular Health Study. Atherosclerosis, 2014, 237, 53-59.	0.8	79
86	Validation of the Preoperative Plasma Fibrinogen Level as a Prognostic Factor in a European Cohort of Patients with Localized Upper Tract Urothelial Carcinoma. Journal of Urology, 2014, 191, 920-925.	0.4	41
87	High preoperative monocyte-lymphocyte ratio (MLR) as a prognostic factor in nonmetastatic clear cell renal cell carcinoma.. Journal of Clinical Oncology, 2014, 32, 534-534.	1.6	2
88	Ethanol Causes Protein Precipitation"New Safety Issues for Catheter Locking Techniques. PLoS ONE, 2013, 8, e84869.	2.5	40
89	Validation of the pretreatment neutrophil-lymphocyte ratio as prognostic factor regarding cancer-specific, metastasis-free, and overall survival in a European cohort of patients with renal cell carcinoma.. Journal of Clinical Oncology, 2013, 31, 410-410.	1.6	2
90	Research update for articles published in EJCI in 2010. European Journal of Clinical Investigation, 2012, 42, 1149-1164.	3.4	1

#	ARTICLE	IF	CITATIONS
91	Low-dose atorvastatin improves dyslipidemia and vascular function in patients with primary biliary cirrhosis after one year of treatment. <i>Atherosclerosis</i> , 2010, 209, 178-183.	0.8	50
92	Serum gamma-glutamyl transferase and mortality in persons undergoing coronary angiographyâ€”The Ludwigshafen Risk and Cardiovascular Health Study. <i>Atherosclerosis</i> , 2010, 208, 564-571.	0.8	42
93	Low density lipoprotein triglycerides and lipoprotein(a) are risk factors for retinal vascular occlusion. <i>Clinica Chimica Acta</i> , 2007, 382, 77-81.	1.1	30
94	Atorvastatin in patients with primary biliary cirrhosis and incomplete biochemical response to ursodeoxycholic acid. <i>Hepatology</i> , 2007, 46, 776-784.	7.3	77
95	ApoE: Crossroads between Alzheimer's Disease and Atherosclerosis. <i>Seminars in Vascular Medicine</i> , 2004, 4, 279-285.	2.1	10