Jing Wang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2385636/publications.pdf

Version: 2024-02-01

		1163117	1125743
16	176	8	13
papers	citations	h-index	g-index
16	16	16	296
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Comparison of Antifungal Prophylaxis Drugs in Patients With Hematological Disease or Undergoing Hematopoietic Stem Cell Transplantation. JAMA Network Open, 2020, 3, e2017652.	5.9	30
2	MYC protein expression is associated with poor prognosis in diffuse large B cell lymphoma patients treated with RCHOP chemotherapy. Tumor Biology, 2014, 35, 6757-6762.	1.8	19
3	Pretreatment C-reactive protein was an independent prognostic factor for patients with diffuse large B-cell lymphoma treated with RCHOP. Clinica Chimica Acta, 2016, 459, 150-154.	1.1	18
4	Prognostic role of pretreatment neutrophil-lymphocyte ratio in patients with diffuse large B-cell lymphoma treated with RCHOP. Medicine (United States), 2016, 95, e4893.	1.0	17
5	Chest CT Findings and Differential Diagnosis of Mycoplasma pneumoniae Pneumonia and Mycoplasma pneumoniae Combined with Streptococcal Pneumonia in Children. Journal of Healthcare Engineering, 2021, 2021, 1-10.	1.9	17
6	Prophylaxis with itraconazole is more effective than prophylaxis with fluconazole in neutropenic patients with hematological malignancies: a meta-analysis of randomized-controlled trials. Medical Oncology, 2010, 27, 1082-1088.	2.5	14
7	Primary central nervous system lymphoma in the United States, 1975–2017. Therapeutic Advances in Hematology, 2022, 13, 204062072110661.	2.5	14
8	Combination of Bcl-2 and MYC protein expression improves high-risk stratification in diffuse large B-cell lymphoma. OncoTargets and Therapy, 2015, 8, 2645.	2.0	10
9	Nomogram for Predicting the Overall Survival of Adult Patients With Primary Gastrointestinal Diffuse Large B Cell Lymphoma: A SEER- Based Study. Frontiers in Oncology, 2020, 10, 1093.	2.8	9
10	Prognostic value of expression of nuclear factor kappa-B/p65 in non-GCB DLBCL patients. Oncotarget, $2017, 8, 9708-9716$.	1.8	9
11	MYC and BCL-2 adjusted-International Prognostic Index (A-IPI) is a better predictor of outcome than the standard IPI for patients with diffuse large B-cell lymphoma treated with R-CHOP. Histology and Histopathology, 2016, 31, 285-92.	0.7	7
12	Methylenetetrahydrofolate reductase gene polymorphisms association with the risk of diffuse large B cell lymphoma: a meta-analysis. Tumor Biology, 2013, 34, 3587-3591.	1.8	3
13	Methylenetetrahydrofolate reductase gene polymorphisms association with the risk of follicular lymphoma: a meta-analysis. Tumor Biology, 2013, 34, 1467-1471.	1.8	3
14	Clonal evolution in a chronic neutrophilic leukemia patient. Hematology, 2019, 24, 455-458.	1.5	3
15	Autologous Cytokine-Induced Killer Cell Immunotherapy for Patients with High-Risk Diffuse Large B Cell Lymphoma After the First Complete Remission. OncoTargets and Therapy, 2020, Volume 13, 5879-5885.	2.0	3
16	A novel mutation in the JAK2 gene (R588S) in a Chinese patient with erythrocytosis. Annals of Hematology, 2020, 100, 2643-2645.	1.8	0