

Tae-You Kim

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/239047/publications.pdf>

Version: 2024-02-01

135
papers

11,204
citations

109264

35
h-index

31818

101
g-index

136
all docs

136
docs citations

136
times ranked

15534
citing authors

#	ARTICLE	IF	CITATIONS
1	Nivolumab in patients with advanced hepatocellular carcinoma (CheckMate 040): an open-label, non-comparative, phase 1/2 dose escalation and expansion trial. <i>Lancet, The</i> , 2017, 389, 2492-2502.	6.3	3,224
2	Ramucirumab plus paclitaxel versus placebo plus paclitaxel in patients with previously treated advanced gastric or gastro-oesophageal junction adenocarcinoma (RAINBOW): a double-blind, randomised phase 3 trial. <i>Lancet Oncology, The</i> , 2014, 15, 1224-1235.	5.1	1,932
3	Efficacy and Safety of Nivolumab Plus Ipilimumab in Patients With Advanced Hepatocellular Carcinoma Previously Treated With Sorafenib. <i>JAMA Oncology</i> , 2020, 6, e204564.	3.4	746
4	Open versus laparoscopic surgery for mid-rectal or low-rectal cancer after neoadjuvant chemoradiotherapy (COREAN trial): survival outcomes of an open-label, non-inferiority, randomised controlled trial. <i>Lancet Oncology, The</i> , 2014, 15, 767-774.	5.1	713
5	Phase 1 study of MRX34, a liposomal miR-34a mimic, in patients with advanced solid tumours. <i>British Journal of Cancer</i> , 2020, 122, 1630-1637.	2.9	472
6	Lineage-dependent gene expression programs influence the immune landscape of colorectal cancer. <i>Nature Genetics</i> , 2020, 52, 594-603.	9.4	380
7	Oxaliplatin, fluorouracil, and leucovorin versus fluorouracil and leucovorin as adjuvant chemotherapy for locally advanced rectal cancer after preoperative chemoradiotherapy (ADORE): an open-label, multicentre, phase 2, randomised controlled trial. <i>Lancet Oncology, The</i> , 2014, 15, 1245-1253.	5.1	336
8	Safety, Efficacy, and Pharmacodynamics of Tremelimumab Plus Durvalumab for Patients With Unresectable Hepatocellular Carcinoma: Randomized Expansion of a Phase I/II Study. <i>Journal of Clinical Oncology</i> , 2021, 39, 2991-3001.	0.8	257
9	Skeletal Muscle Depletion Predicts the Prognosis of Patients with Advanced Pancreatic Cancer Undergoing Palliative Chemotherapy, Independent of Body Mass Index. <i>PLoS ONE</i> , 2015, 10, e0139749.	1.1	183
10	Histone deacetylase inhibitor, suberoylanilide hydroxamic acid (SAHA), enhances anti-tumor effects of the poly (ADP-ribose) polymerase (PARP) inhibitor olaparib in triple-negative breast cancer cells. <i>Breast Cancer Research</i> , 2015, 17, 33.	2.2	138
11	Phase I/II study of durvalumab and tremelimumab in patients with unresectable hepatocellular carcinoma (HCC): Phase I safety and efficacy analyses.. <i>Journal of Clinical Oncology</i> , 2017, 35, 4073-4073.	0.8	133
12	STAT3 inhibits the degradation of HIF-1 α by pVHL-mediated ubiquitination. <i>Experimental and Molecular Medicine</i> , 2008, 40, 479.	3.2	103
13	Oxaliplatin-Based Adjuvant Chemotherapy for Rectal Cancer After Preoperative Chemoradiotherapy (ADORE): Long-Term Results of a Randomized Controlled Trial. <i>Journal of Clinical Oncology</i> , 2019, 37, 3111-3123.	0.8	100
14	Loss of CDX2 expression is associated with poor prognosis in colorectal cancer patients. <i>World Journal of Gastroenterology</i> , 2015, 21, 1457.	1.4	98
15	Tumor Mutation Burden and Prognosis in Patients with Colorectal Cancer Treated with Adjuvant Fluoropyrimidine and Oxaliplatin. <i>Clinical Cancer Research</i> , 2019, 25, 6141-6147.	3.2	98
16	Soluble programmed death-ligand 1 (sPDL1) and neutrophil-to-lymphocyte ratio (NLR) predicts survival in advanced biliary tract cancer patients treated with palliative chemotherapy. <i>Oncotarget</i> , 2016, 7, 76604-76612.	0.8	93
17	Phase I Study of OPB-31121, an Oral STAT3 Inhibitor, in Patients with Advanced Solid Tumors. <i>Cancer Research and Treatment</i> , 2015, 47, 607-615.	1.3	93
18	Randomized phase II trial of nimotuzumab plus irinotecan versus irinotecan alone as second-line therapy for patients with advanced gastric cancer. <i>Gastric Cancer</i> , 2015, 18, 824-832.	2.7	91

#	ARTICLE	IF	CITATIONS
19	Nivolumab (nivo) in sorafenib (sor)-naive and -experienced pts with advanced hepatocellular carcinoma (HCC): CheckMate 040 study.. <i>Journal of Clinical Oncology</i> , 2017, 35, 4013-4013.	0.8	76
20	Association Between <i>Fusobacterium nucleatum</i> , Pathway Mutation, and Patient Prognosis in Colorectal Cancer. <i>Annals of Surgical Oncology</i> , 2018, 25, 3389-3395.	0.7	69
21	Therapeutic implication of HER2 in advanced biliary tract cancer. <i>Oncotarget</i> , 2016, 7, 58007-58021.	0.8	63
22	Histone Deacetylase Inhibitors for Cancer Therapy. <i>Epigenetics</i> , 2006, 1, 15-24.	1.3	61
23	Optimal Patient Selection for Trastuzumab Treatment in HER2-Positive Advanced Gastric Cancer. <i>Clinical Cancer Research</i> , 2015, 21, 2520-2529.	3.2	59
24	The Impact of Diabetes Mellitus and Metformin Treatment on Survival of Patients with Advanced Pancreatic Cancer Undergoing Chemotherapy. <i>Cancer Research and Treatment</i> , 2016, 48, 171-179.	1.3	56
25	Prognostic implication of antitumor immunity measured by the neutrophil-lymphocyte ratio and serum cytokines and angiogenic factors in gastric cancer. <i>Gastric Cancer</i> , 2017, 20, 254-262.	2.7	51
26	Association between mutations of critical pathway genes and survival outcomes according to the tumor location in colorectal cancer. <i>Cancer</i> , 2017, 123, 3513-3523.	2.0	50
27	Neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, and their dynamic changes during chemotherapy is useful to predict a more accurate prognosis of advanced biliary tract cancer. <i>Oncotarget</i> , 2017, 8, 2329-2341.	0.8	50
28	Open versus laparoscopic surgery for mid or low rectal cancer after neoadjuvant chemoradiotherapy (COREAN trial): 10-year follow-up of an open-label, non-inferiority, randomised controlled trial. <i>The Lancet Gastroenterology and Hepatology</i> , 2021, 6, 569-577.	3.7	50
29	Methylation and microsatellite status and recurrence following adjuvant FOLFOX in colorectal cancer. <i>International Journal of Cancer</i> , 2013, 132, 2209-2216.	2.3	49
30	Clinical Implications of VEGF, TGF-beta1, and IL-1beta in Patients with Advanced Non-small Cell Lung Cancer. <i>Cancer Research and Treatment</i> , 2013, 45, 325-333.	1.3	49
31	Activation of WNT/catenin signaling results in resistance to a dual PI3K/mTOR inhibitor in colorectal cancer cells harboring PIK3CA mutations. <i>International Journal of Cancer</i> , 2019, 144, 389-401.	2.3	48
32	A Phase III Randomized Trial of Combined Chemoradiotherapy Versus Radiotherapy Alone in Locally Advanced Non-Small-Cell Lung Cancer. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 2002, 25, 238-243.	0.6	45
33	Adverse prognostic impact of the CpG island methylator phenotype in metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2016, 115, 164-171.	2.9	43
34	A Phase II Study of Avelumab Monotherapy in Patients with Mismatch Repair-Deficient/Microsatellite Instability-High or POLE-Mutated Metastatic or Unresectable Colorectal Cancer. <i>Cancer Research and Treatment</i> , 2020, 52, 1135-1144.	1.3	43
35	Distinct clinical outcomes of two CIMP-positive colorectal cancer subtypes based on a revised CIMP classification system. <i>British Journal of Cancer</i> , 2017, 116, 1012-1020.	2.9	40
36	Phase I Dose-Finding Study of OPB-111077, a Novel STAT3 Inhibitor, in Patients with Advanced Hepatocellular Carcinoma. <i>Cancer Research and Treatment</i> , 2019, 51, 510-518.	1.3	39

#	ARTICLE	IF	CITATIONS
37	Whole-Slide Image Analysis Reveals Quantitative Landscape of Tumor Immune Microenvironment in Colorectal Cancers. <i>Clinical Cancer Research</i> , 2020, 26, 870-881.	3.2	37
38	Phase I/II study of first-line combination therapy with sorafenib plus resminostat, an oral HDAC inhibitor, versus sorafenib monotherapy for advanced hepatocellular carcinoma in east Asian patients. <i>Investigational New Drugs</i> , 2018, 36, 1072-1084.	1.2	32
39	p53 expression status is associated with cancer-specific survival in stage III and high-risk stage II colorectal cancer patients treated with oxaliplatin-based adjuvant chemotherapy. <i>British Journal of Cancer</i> , 2019, 120, 797-805.	2.9	32
40	Effectiveness of nivolumab versus regorafenib in hepatocellular carcinoma patients who failed sorafenib treatment. <i>Clinical and Molecular Hepatology</i> , 2020, 26, 328-339.	4.5	32
41	Weight loss at the first month of palliative chemotherapy predicts survival outcomes in patients with advanced gastric cancer. <i>Gastric Cancer</i> , 2016, 19, 597-606.	2.7	31
42	Safety and antitumor activity of nivolumab (nivo) in patients (pts) with advanced hepatocellular carcinoma (HCC): Interim analysis of dose-expansion cohorts from the phase 1/2 CheckMate-040 study.. <i>Journal of Clinical Oncology</i> , 2016, 34, 4078-4078.	0.8	30
43	Total lesion glycolysis (TLG) as an imaging biomarker in metastatic colorectal cancer patients treated with regorafenib. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2017, 44, 757-764.	3.3	27
44	Phase II Study of Avelumab in Patients with Advanced Hepatocellular Carcinoma Previously Treated with Sorafenib. <i>Clinical Cancer Research</i> , 2021, 27, 713-718.	3.2	27
45	A Phase III Study to Compare the Efficacy and Safety of Paclitaxel Versus Irinotecan in Patients with Metastatic or Recurrent Gastric Cancer Who Failed in First-line Therapy (KCSG ST10-01). <i>Oncologist</i> , 2019, 24, 18-e24.	1.9	25
46	Annexin A10 expression in colorectal cancers with emphasis on the serrated neoplasia pathway. <i>World Journal of Gastroenterology</i> , 2015, 21, 9749.	1.4	24
47	A Phase I/Randomized Phase II Study to Evaluate the Safety, Pharmacokinetics, and Efficacy of Nintedanib versus Sorafenib in Asian Patients with Advanced Hepatocellular Carcinoma. <i>Liver Cancer</i> , 2018, 7, 165-178.	4.2	23
48	Randomised Phase 1b/2 trial of tepotinib vs sorafenib in Asian patients with advanced hepatocellular carcinoma with MET overexpression. <i>British Journal of Cancer</i> , 2021, 125, 200-208.	2.9	22
49	Dynamic cohesin-mediated chromatin architecture controls epithelial-mesenchymal plasticity in cancer. <i>EMBO Reports</i> , 2016, 17, 1343-1359.	2.0	21
50	MRX34, a liposomal miR-34 mimic, in patients with advanced solid tumors: Final dose-escalation results from a first-in-human phase I trial of microRNA therapy.. <i>Journal of Clinical Oncology</i> , 2016, 34, 2508-2508.	0.8	21
51	Clinical Implication of Anti-Angiogenic Effect of Regorafenib in Metastatic Colorectal Cancer. <i>PLoS ONE</i> , 2015, 10, e0145004.	1.1	20
52	Phase I Study of CKD-516, a Novel Vascular Disrupting Agent, in Patients with Advanced Solid Tumors. <i>Cancer Research and Treatment</i> , 2016, 48, 28-36.	1.3	20
53	Clinical and pathological significance of ROS1 expression in intrahepatic cholangiocarcinoma. <i>BMC Cancer</i> , 2015, 15, 721.	1.1	19
54	TTP as a surrogate endpoint in advanced hepatocellular carcinoma treated with molecular targeted therapy: meta-analysis of randomised controlled trials. <i>British Journal of Cancer</i> , 2016, 115, 1201-1205.	2.9	19

#	ARTICLE	IF	CITATIONS
55	Korean Cancer Patients's Awareness of Clinical Trials, Perceptions on the Benefit and Willingness to Participate. <i>Cancer Research and Treatment</i> , 2017, 49, 1033-1043.	1.3	19
56	Liquid biopsy-based tumor profiling for metastatic colorectal cancer patients with ultra-deep targeted sequencing. <i>PLoS ONE</i> , 2020, 15, e0232754.	1.1	19
57	Nivolumab dose escalation and expansion in patients with advanced hepatocellular carcinoma (HCC): The CheckMate 040 study.. <i>Journal of Clinical Oncology</i> , 2017, 35, 226-226.	0.8	19
58	Reduced cohesin destabilizes high-level gene amplification by disrupting pre-replication complex bindings in human cancers with chromosomal instability. <i>Nucleic Acids Research</i> , 2016, 44, 558-572.	6.5	18
59	The Impact of Body Mass Index Dynamics on Survival of Patients With Advanced Pancreatic Cancer Receiving Chemotherapy. <i>Journal of Pain and Symptom Management</i> , 2014, 48, 13-25.	0.6	17
60	Different prognostic effect of CpG island methylation according to sex in colorectal cancer patients treated with adjuvant FOLFOX. <i>Clinical Epigenetics</i> , 2015, 7, 63.	1.8	17
61	Patient-derived organoids as a preclinical platform for precision medicine in colorectal cancer. <i>Molecular Oncology</i> , 2022, 16, 2396-2412.	2.1	17
62	Transcriptional induction of DLC-1 gene through Sp1 sites by histone deacetylase inhibitors in gastric cancer cells. <i>Experimental and Molecular Medicine</i> , 2008, 40, 639.	3.2	15
63	Metabolic Characteristics of Advanced Biliary Tract Cancer Using 18F-Fluorodeoxyglucose Positron Emission Tomography and Their Clinical Implications. <i>Oncologist</i> , 2015, 20, 926-933.	1.9	15
64	Development of a Nomogram to Predict the Recurrence Score of 21-Gene Prediction Assay in Hormone Receptor-Positive Early Breast Cancer. <i>Clinical Breast Cancer</i> , 2020, 20, 98-107.e1.	1.1	15
65	Survival Outcomes According to Adjuvant Treatment and Prognostic Factors Including Host Immune Markers in Patients with Curatively Resected Ampulla of Vater Cancer. <i>PLoS ONE</i> , 2016, 11, e0151406.	1.1	15
66	Splenomegaly and Its Associations with Genetic Polymorphisms and Treatment Outcome in Colorectal Cancer Patients Treated with Adjuvant FOLFOX. <i>Cancer Research and Treatment</i> , 2016, 48, 990-997.	1.3	15
67	Circulating tumor DNA sequencing in colorectal cancer patients treated with first-line chemotherapy with anti-EGFR. <i>Scientific Reports</i> , 2021, 11, 16333.	1.6	14
68	More Accurate Prediction of Metastatic Pancreatic Cancer Patients' Survival with Prognostic Model Using Both Host Immunity and Tumor Metabolic Activity. <i>PLoS ONE</i> , 2016, 11, e0145692.	1.1	14
69	Signature of cytokines and angiogenic factors (CAFs) defines a clinically distinct subgroup of gastric cancer. <i>Gastric Cancer</i> , 2017, 20, 164-174.	2.7	13
70	Cyclophosphamide, Methotrexate, and 5-Fluorouracil as Palliative Treatment for Heavily Pretreated Patients with Metastatic Breast Cancer: A Multicenter Retrospective Analysis. <i>Journal of Breast Cancer</i> , 2017, 20, 347.	0.8	13
71	TGF- β 2 Suppresses COX-2 Expression by Tristetraprolin-Mediated RNA Destabilization in A549 Human Lung Cancer Cells. <i>Cancer Research and Treatment</i> , 1970, 47, 101-109.	1.3	13
72	Treatment Patterns and Changes in Quality of Life during First-Line Palliative Chemotherapy in Korean Patients with Advanced Gastric Cancer. <i>Cancer Research and Treatment</i> , 2019, 51, 223-239.	1.3	13

#	ARTICLE	IF	CITATIONS
73	NFATC3-PLA2G15 Fusion Transcript Identified by RNA Sequencing Promotes Tumor Invasion and Proliferation in Colorectal Cancer Cell Lines. <i>Cancer Research and Treatment</i> , 2019, 51, 391-401.	1.3	13
74	EMT-mediated regulation of CXCL1/5 for resistance to anti-EGFR therapy in colorectal cancer. <i>Oncogene</i> , 2022, 41, 2026-2038.	2.6	13
75	Hypermethylation of PDX1, EN2, and MSX1 predicts the prognosis of colorectal cancer. <i>Experimental and Molecular Medicine</i> , 2022, 54, 156-168.	3.2	13
76	Chasing targets for EGFR tyrosine kinase inhibitors in non-small-cell lung cancer: Asian perspectives. <i>Expert Review of Molecular Diagnostics</i> , 2007, 7, 821-836.	1.5	12
77	Landscape of Actionable Genetic Alterations Profiled from 1,071 Tumor Samples in Korean Cancer Patients. <i>Cancer Research and Treatment</i> , 2019, 51, 211-222.	1.3	12
78	Prognostic impact of AJCC response criteria for neoadjuvant chemotherapy in stage II/III breast cancer patients: breast cancer subtype analyses. <i>BMC Cancer</i> , 2016, 16, 515.	1.1	11
79	Metabolic landscape of advanced gastric cancer according to HER2 and its prognostic implications. <i>Gastric Cancer</i> , 2016, 19, 421-430.	2.7	11
80	Comparison of Quantitative Methods on FDG PET/CT for Treatment Response Evaluation of Metastatic Colorectal Cancer. <i>Nuclear Medicine and Molecular Imaging</i> , 2017, 51, 147-153.	0.6	11
81	Blood-Based Detection of Colorectal Cancer Using Cancer-Specific DNA Methylation Markers. <i>Diagnostics</i> , 2021, 11, 51.	1.3	11
82	Regorafenib as Salvage Treatment in Korean Patients with Refractory Metastatic Colorectal Cancer. <i>Cancer Research and Treatment</i> , 2015, 47, 790-795.	1.3	10
83	Src as a Therapeutic Target in Biliary Tract Cancer. <i>Molecular Cancer Therapeutics</i> , 2016, 15, 1515-1524.	1.9	10
84	Macrophage migration inhibitory factor promotes resistance to MEK blockade in KRAS mutant colorectal cancer cells. <i>Molecular Oncology</i> , 2018, 12, 1398-1409.	2.1	10
85	Interpretation of EBV infection in pan-cancer genome considering viral life cycle: LiEB (Life cycle of) Tj ETQq1 1 0.784314 rgBT /Overlo	1.6	10
86	Body mass index and body weight change during adjuvant chemotherapy in colon cancer patients: results from the AVANT trial. <i>Scientific Reports</i> , 2020, 10, 19467.	1.6	10
87	Tumor microenvironment-adjusted prognostic implications of the KRAS mutation subtype in patients with stage III colorectal cancer treated with adjuvant FOLFOX. <i>Scientific Reports</i> , 2021, 11, 14609.	1.6	10
88	Concurrent Chemoradiotherapy Versus Chemotherapy Alone for Unresectable Locally Advanced Pancreatic Cancer: A Retrospective Cohort Study. <i>Cancer Research and Treatment</i> , 2016, 48, 1045-1055.	1.3	10
89	Prognostic influence of body mass index and body weight gain during adjuvant FOLFOX chemotherapy in Korean colorectal cancer patients. <i>BMC Cancer</i> , 2015, 15, 690.	1.1	9
90	Perceived needs for the information communication technology (ICT)-based personalized health management program, and its association with information provision, health-related quality of life (HRQOL), and decisional conflict in cancer patients. <i>Psycho-Oncology</i> , 2017, 26, 1810-1817.	1.0	9

#	ARTICLE	IF	CITATIONS
91	Phase II Trial of Postoperative Adjuvant Gemcitabine and Cisplatin Chemotherapy Followed by Chemoradiotherapy with Gemcitabine in Patients with Resected Pancreatic Cancer. <i>Cancer Research and Treatment</i> , 2021, 53, 1096-1103.	1.3	9
92	Prognostic implication of serum hepatocyte growth factor in stage II/III breast cancer patients who received neoadjuvant chemotherapy. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 707-714.	1.2	8
93	A phase II trial of S-1 and oxaliplatin in patients with advanced hepatocellular carcinoma. <i>BMC Cancer</i> , 2018, 18, 252.	1.1	7
94	RAINBOW: A global, phase III, randomized, double-blind study of ramucirumab plus paclitaxel versus placebo plus paclitaxel patients with previously treated gastric or gastroesophageal junction (GEJ) adenocarcinoma: Quality-of-life (QoL) results.. <i>Journal of Clinical Oncology</i> , 2014, 32, 4058-4058.	0.8	7
95	Efficacy and safety of nintedanib versus sorafenib in Asian patients with advanced hepatocellular carcinoma (HCC): A randomized phase II trial.. <i>Journal of Clinical Oncology</i> , 2015, 33, 339-339.	0.8	7
96	Complete responses (CR) in patients receiving atezolizumab (atezo) + bevacizumab (bev) versus sorafenib (sor) in IMbrave150: A phase III clinical trial for unresectable hepatocellular carcinoma (HCC).. <i>Journal of Clinical Oncology</i> , 2020, 38, 4596-4596.	0.8	7
97	Immunohistochemical features associated with sensitivity to lapatinib-plus-capecitabine and resistance to trastuzumab in HER2-positive breast cancer. <i>Anticancer Research</i> , 2014, 34, 4275-80.	0.5	7
98	Evaluation of Lapatinib Powder-Entrapped Biodegradable Polymeric Microstructures Fabricated by X-Ray Lithography for a Targeted and Sustained Drug Delivery System. <i>Materials</i> , 2015, 8, 519-534.	1.3	6
99	Identification of Long-Range Epigenetic Silencing on Chromosome 15q25 and Its Clinical Implication in Gastric Cancer. <i>American Journal of Pathology</i> , 2015, 185, 666-678.	1.9	6
100	Targeted next-generation sequencing-based detection of microsatellite instability in colorectal carcinomas. <i>PLoS ONE</i> , 2021, 16, e0246356.	1.1	6
101	A phase I study of HM781-36B, a novel pan-HER inhibitor, in patients (pts) with advanced solid tumors.. <i>Journal of Clinical Oncology</i> , 2012, 30, 3076-3076.	0.8	6
102	Phase I study of PF-03446962 (anti-ALK-1 mAb) in hepatocellular carcinoma (HCC).. <i>Journal of Clinical Oncology</i> , 2013, 31, 4121-4121.	0.8	6
103	The distinct signatures of VEGF and soluble VEGFR2 increase prognostic implication in gastric cancer. <i>American Journal of Cancer Research</i> , 2015, 5, 3376-88.	1.4	6
104	Ramucirumab in patients with previously treated advanced hepatocellular carcinoma: Impact of liver disease aetiology. <i>Liver International</i> , 2021, 41, 2759-2767.	1.9	5
105	BioPATH: A Biomarker Study in Asian Patients with HER2+ Advanced Breast Cancer Treated with Lapatinib and Other Anti-HER2 Therapy. <i>Cancer Research and Treatment</i> , 2019, 51, 1527-1539.	1.3	5
106	Radiation Response Prediction Model Based on Integrated Clinical and Genomic Data Analysis. <i>Cancer Research and Treatment</i> , 2022, 54, 383-395.	1.3	4
107	First-in-human trial of microRNA cancer therapy with MRX34, a liposomal miR-34 mimic: Phase Ia expansion in patients with advanced solid tumors.. <i>Journal of Clinical Oncology</i> , 2016, 34, TPS2597-TPS2597.	0.8	4
108	KEYNOTE-164: Phase 2 study of pembrolizumab for patients with previously treated, microsatellite instability-high advanced colorectal carcinoma.. <i>Journal of Clinical Oncology</i> , 2016, 34, TPS3631-TPS3631.	0.8	4

#	ARTICLE	IF	CITATIONS
109	Resminostat and sorafenib combination therapy for advanced hepatocellular carcinoma in patients previously untreated with systemic chemotherapy.. Journal of Clinical Oncology, 2017, 35, 252-252.	0.8	4
110	Phenotype-based single cell sequencing identifies diverse genetic subclones in CD133 positive cancer stem cells. Biochemical and Biophysical Research Communications, 2021, 558, 209-215.	1.0	3
111	Phase I dose-finding study of OPB-111077, a novel STAT3 inhibitor, in patients with advanced hepatocellular carcinoma.. Journal of Clinical Oncology, 2018, 36, 4078-4078.	0.8	3
112	The impact of diabetes mellitus and metformin on survival of patients with advanced pancreatic cancer receiving chemotherapy.. Journal of Clinical Oncology, 2013, 31, 4044-4044.	0.8	3
113	Association of pathway mutation with survival after recurrence in colorectal cancer patients treated with adjuvant fluoropyrimidine and oxaliplatin chemotherapy. BMC Cancer, 2019, 19, 421.	1.1	2
114	Efficacy of infusional 5-fluorouracil, doxorubicin, and mitomycin-C (iFAM) chemotherapy and analysis of prognostic factors in previously treated advanced hepatocellular carcinoma.. Journal of Clinical Oncology, 2012, 30, 269-269.	0.8	2
115	Prognostic Impact of Extramural Lymphatic, Vascular, and Perineural Invasion in Stage II Colon Cancer: A Comparison With Intramural Invasion. Diseases of the Colon and Rectum, 2022, Publish Ahead of Print, .	0.7	2
116	A phase 1 dose-escalation and dose-expansion study to assess the safety and efficacy of CKD-516, a novel vascular disrupting agent, in combination with Irinotecan in patients with previously treated metastatic colorectal cancer. Investigational New Drugs, 2021, 39, 1335-1347.	1.2	1
117	Efficacy of Letrozole as First-Line Treatment of Postmenopausal Women with Hormone Receptor-Positive Metastatic Breast Cancer in Korea. Cancer Research and Treatment, 2017, 49, 454-463.	1.3	1
118	Usefulness of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, and their dynamic changes during chemotherapy to predict prognosis of advanced biliary tract cancer.. Journal of Clinical Oncology, 2017, 35, 416-416.	0.8	1
119	Longitudinal monitoring of circulating tumor DNA (ctDNA) during disease course of metastatic colorectal cancer (mCRC).. Journal of Clinical Oncology, 2022, 40, 189-189.	0.8	1
120	Role of Dedicated Subspecialized Radiologists in Multidisciplinary Team Discussions on Lower Gastrointestinal Tract Cancers. Korean Journal of Radiology, 2022, 23, .	1.5	1
121	Fabrication of biodegradable polymeric micro chambers encapsulated with pulverized drug for bacteria-based microrobots. , 2014, , .		0
122	Prognostic Role of Body Mass Index in Advanced Small Bowel Adenocarcinoma Patients Receiving Palliative Chemotherapy. Nutrition and Cancer, 2016, 68, 750-755.	0.9	0
123	Reply to "Comment on "Distinct clinical outcomes of two CIMP-positive colorectal cancer subtypes based on a revised CIMP classification system" British Journal of Cancer, 2018, 118, e4-e4.	2.9	0
124	A Case of Extranodal NK/T Cell Lymphoma, Nasal Type Involving Anus. The Korean Journal of Hematology, 2005, 40, 192.	0.7	0
125	Genetic polymorphisms and ethnic difference in outcome of adjuvant FOLFOX chemotherapy in Korean patients with colon cancer.. Journal of Clinical Oncology, 2012, 30, 623-623.	0.8	0
126	Methylations of NEUROG1, p16, and MLH1 and recurrence following adjuvant FOLFOX in colorectal cancer.. Journal of Clinical Oncology, 2012, 30, 3624-3624.	0.8	0

#	ARTICLE	IF	CITATIONS
127	The efficacy and toxicity of 3-weekly TS-1 containing chemotherapy in patients with unresectable advanced gastric cancer.. Journal of Clinical Oncology, 2012, 30, e14580-e14580.	0.8	0
128	Breast-conserving surgery after neoadjuvant chemotherapy for stage III breast cancer patients.. Journal of Clinical Oncology, 2012, 30, e11532-e11532.	0.8	0
129	Phosphorylated Akt expression as a favorable prognostic factor for patients undergoing curative resection and adjuvant chemoradiotherapy for proximal extrahepatic bile duct cancer.. Journal of Clinical Oncology, 2013, 31, 182-182.	0.8	0
130	The impact of body mass index dynamics on survival of patients with advanced pancreatic cancer receiving chemotherapy.. Journal of Clinical Oncology, 2013, 31, e15066-e15066.	0.8	0
131	ABCB1 polymorphism as a prognostic factor in breast cancer patients with neoadjuvant chemotherapy.. Journal of Clinical Oncology, 2014, 32, 1038-1038.	0.8	0
132	TTP or PFS as a surrogate endpoint in advanced hepatocellular carcinoma treated with systemic therapy.. Journal of Clinical Oncology, 2016, 34, 4075-4075.	0.8	0
133	Korean Cancer Patients's Awareness of Clinical Trials: Perceptions on the benefit and willingness to participate.. Journal of Clinical Oncology, 2016, 34, 10067-10067.	0.8	0
134	Skeletal muscle depletion to predict survival of patients with advanced biliary tract cancer undergoing palliative chemotherapy.. Journal of Clinical Oncology, 2017, 35, 460-460.	0.8	0
135	Surrogate endpoint in advanced hepatocellular carcinoma treated with molecular targeted therapy: Meta-analysis of randomized controlled trials.. Journal of Clinical Oncology, 2017, 35, 454-454.	0.8	0