

Ladislav Senolt

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/239363/publications.pdf>

Version: 2024-02-01

144
papers

13,486
citations

101384

36
h-index

26548

107
g-index

189
all docs

189
docs citations

189
times ranked

18989
citing authors

#	ARTICLE	IF	CITATIONS
1	Plasma Hsp90 levels in patients with systemic sclerosis and relation to lung and skin involvement: a cross-sectional and longitudinal study. <i>Scientific Reports</i> , 2021, 11, 1.	1.6	9,439
2	The role of resistin as a regulator of inflammation: Implications for various human pathologies. <i>Clinical Immunology</i> , 2009, 133, 157-170.	1.4	345
3	Activation of STAT3 integrates common profibrotic pathways to promote fibroblast activation and tissue fibrosis. <i>Nature Communications</i> , 2017, 8, 1130.	5.8	245
4	Resistin in rheumatoid arthritis synovial tissue, synovial fluid and serum. <i>Annals of the Rheumatic Diseases</i> , 2006, 66, 458-463.	0.5	226
5	Association of circulating miR-223 and miR-16 with disease activity in patients with early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 1898-1904.	0.5	165
6	Increased adiponectin is negatively linked to the local inflammatory process in patients with rheumatoid arthritis. <i>Cytokine</i> , 2006, 35, 247-252.	1.4	141
7	Increased level of cytokines and matrix metalloproteinases in osteoarthritic subchondral bone. <i>Cytokine</i> , 2007, 38, 151-156.	1.4	140
8	Prospective new biological therapies for rheumatoid arthritis. <i>Autoimmunity Reviews</i> , 2009, 9, 102-107.	2.5	119
9	Increased serum adiponectin levels in female patients with erosive compared with non-erosive osteoarthritis: Figure 1. <i>Annals of the Rheumatic Diseases</i> , 2009, 68, 295-296.	0.5	112
10	Vaspin and omentin: new adipokines differentially regulated at the site of inflammation in rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, 1410-1411.	0.5	94
11	Up-regulation of metastasis-promoting S100A4 (Mts-1) in rheumatoid arthritis: Putative involvement in the pathogenesis of rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 2007, 56, 779-789.	6.7	91
12	Hyaluronic acid levels may have predictive value for the progression of knee osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2004, 12, 277-283.	0.6	89
13	Heat shock protein 90 (Hsp90) inhibition targets canonical TGF- β 2 signalling to prevent fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 1215-1222.	0.5	78
14	S100A4 is expressed at site of invasion in rheumatoid arthritis synovium and modulates production of matrix metalloproteinases. <i>Annals of the Rheumatic Diseases</i> , 2006, 65, 1645-1648.	0.5	73
15	The metastasis-associated protein S100A4 promotes the inflammatory response of mononuclear cells via the TLR4 signalling pathway in rheumatoid arthritis. <i>Rheumatology</i> , 2014, 53, 1520-1526.	0.9	72
16	Decreases in serum levels of S100A8/9 (calprotectin) correlate with improvements in total swollen joint count in patients with recent-onset rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2011, 13, R122.	1.6	69
17	Demonstration of extracellular peptidylarginine deiminase (PAD) activity in synovial fluid of patients with rheumatoid arthritis using a novel assay for citrullination of fibrinogen. <i>Arthritis Research and Therapy</i> , 2014, 16, 498.	1.6	68
18	Increased pentosidine, an advanced glycation end product, in serum and synovial fluid from patients with knee osteoarthritis and its relation with cartilage oligomeric matrix protein. <i>Annals of the Rheumatic Diseases</i> , 2005, 64, 886-890.	0.5	67

#	ARTICLE	IF	CITATIONS
19	Serum calprotectin (S100A8/9): an independent predictor of ultrasound synovitis in patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2015, 17, 252.	1.6	60
20	Increasing the infliximab dose in rheumatoid arthritis patients: a randomised, double blind study failed to confirm its efficacy. <i>Annals of the Rheumatic Diseases</i> , 2009, 68, 1285-1289.	0.5	59
21	An update on biomarkers in axial spondyloarthritis. <i>Autoimmunity Reviews</i> , 2016, 15, 501-509.	2.5	58
22	The metastasis associated protein S100A4: a potential novel link to inflammation and consequent aggressive behaviour of rheumatoid arthritis synovial fibroblasts. <i>Annals of the Rheumatic Diseases</i> , 2008, 67, 1499-1504.	0.5	57
23	The pre-clinical phase of rheumatoid arthritis: From risk factors to prevention of arthritis. <i>Autoimmunity Reviews</i> , 2021, 20, 102797.	2.5	56
24	Low circulating Dickkopf-1 and its link with severity of spinal involvement in diffuse idiopathic skeletal hyperostosis. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, 71-74.	0.5	55
25	The transcription factor GLI2 as a downstream mediator of transforming growth factor- β -induced fibroblast activation in SSc. <i>Annals of the Rheumatic Diseases</i> , 2017, 76, 756-764.	0.5	53
26	S100A4 amplifies TGF- β -induced fibroblast activation in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 1748-1755.	0.5	52
27	The IL-20 Cytokine Family in Rheumatoid Arthritis and Spondyloarthritis. <i>Frontiers in Immunology</i> , 2018, 9, 2226.	2.2	51
28	MicroRNA-125b: association with disease activity and the treatment response of patients with early rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2016, 18, 124.	1.6	48
29	Pro-inflammatory effects of interleukin-35 in rheumatoid arthritis. <i>Cytokine</i> , 2015, 73, 36-43.	1.4	44
30	Increased levels of peptidylarginine deiminase 2 in synovial fluid from anti-CCP-positive rheumatoid arthritis patients: Association with disease activity and inflammatory markers. <i>Rheumatology</i> , 2016, 55, 918-927.	0.9	42
31	Emerging therapies in rheumatoid arthritis: focus on monoclonal antibodies. <i>F1000Research</i> , 2019, 8, 1549.	0.8	41
32	Metastasis-inducing S100A4 protein is associated with the disease activity of rheumatoid arthritis. <i>Rheumatology</i> , 2009, 48, 1590-1594.	0.9	40
33	Clinical improvement and reduction in serum calprotectin levels after an intensive exercise programme for patients with ankylosing spondylitis and non-radiographic axial spondyloarthritis. <i>Arthritis Research and Therapy</i> , 2016, 18, 275.	1.6	40
34	Monoclonal antibodies to human cartilage oligomeric matrix protein: epitope mapping and characterization of sandwich ELISA. <i>Clinica Chimica Acta</i> , 2003, 328, 59-69.	0.5	39
35	Serum hyaluronic acid as a potential marker with a predictive value for further radiographic progression of hand osteoarthritis. <i>Osteoarthritis and Cartilage</i> , 2009, 17, 1615-1619.	0.6	39
36	Efficacy and Safety of Anti-Interleukin-20 Monoclonal Antibody in Patients With Rheumatoid Arthritis: A Randomized Phase IIa Trial. <i>Arthritis and Rheumatology</i> , 2015, 67, 1438-1448.	2.9	39

#	ARTICLE	IF	CITATIONS
37	Tribbles homologue 3 stimulates canonical TGF- β 2 signalling to regulate fibroblast activation and tissue fibrosis. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 609-616.	0.5	38
38	Increased serum concentration of immune cell derived microparticles in polymyositis/dermatomyositis. <i>Immunology Letters</i> , 2010, 128, 124-130.	1.1	35
39	The metastasis promoting protein S100A4 is increased in idiopathic inflammatory myopathies. <i>Rheumatology</i> , 2011, 50, 1766-1772.	0.9	35
40	Calgizzarin (S100A11): a novel inflammatory mediator associated with disease activity of rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2017, 19, 79.	1.6	35
41	Progranulin Is Associated with Disease Activity in Patients with Rheumatoid Arthritis. <i>Mediators of Inflammation</i> , 2015, 2015, 1-6.	1.4	34
42	Relationship between serum calprotectin (S100A8/9) and clinical, laboratory and ultrasound parameters of disease activity in rheumatoid arthritis: A large cohort study. <i>PLoS ONE</i> , 2017, 12, e0183420.	1.1	32
43	The level of serum visfatin (PBEF) is associated with total number of B cells in patients with rheumatoid arthritis and decreases following B cell depletion therapy. <i>Cytokine</i> , 2011, 55, 116-121.	1.4	31
44	S100A4 (Mts1): Is there any relation to the pathogenesis of rheumatoid arthritis?. <i>Autoimmunity Reviews</i> , 2006, 5, 129-131.	2.5	30
45	Association between circulating miRNAs and spinal involvement in patients with axial spondyloarthritis. <i>PLoS ONE</i> , 2017, 12, e0185323.	1.1	30
46	Adiponectin relation to skin changes and dyslipidemia in systemic sclerosis. <i>Cytokine</i> , 2012, 58, 165-168.	1.4	29
47	Interleukin 35 Synovial Fluid Levels Are Associated with Disease Activity of Rheumatoid Arthritis. <i>PLoS ONE</i> , 2015, 10, e0132674.	1.1	28
48	Laboratory biomarkers or imaging in the diagnostics of rheumatoid arthritis?. <i>BMC Medicine</i> , 2014, 12, 49.	2.3	27
49	Relative efficiencies of peptidylarginine deiminase 2 and 4 in generating target sites for anti-citrullinated protein antibodies in fibrinogen, alpha-enolase and histone H3. <i>PLoS ONE</i> , 2018, 13, e0203214.	1.1	27
50	Decreased Circulating Visfatin Is Associated with Improved Disease Activity in Early Rheumatoid Arthritis: Data from the PERAC Cohort. <i>PLoS ONE</i> , 2014, 9, e103495.	1.1	25
51	Serum calprotectin may reflect inflammatory activity in patients with active rheumatoid arthritis despite normal to low C-reactive protein. <i>Clinical Rheumatology</i> , 2018, 37, 2055-2062.	1.0	23
52	Circulating S100 proteins effectively discriminate SLE patients from healthy controls: a cross-sectional study. <i>Rheumatology International</i> , 2019, 39, 469-478.	1.5	23
53	Resistin in idiopathic inflammatory myopathies. <i>Arthritis Research and Therapy</i> , 2012, 14, R111.	1.6	21
54	MicroRNA-331 and microRNA-151-3p as biomarkers in patients with ST-segment elevation myocardial infarction. <i>Scientific Reports</i> , 2020, 10, 5845.	1.6	21

#	ARTICLE	IF	CITATIONS
55	Lower serum clusterin levels in patients with erosive hand osteoarthritis are associated with more pain. <i>BMC Musculoskeletal Disorders</i> , 2018, 19, 264.	0.8	20
56	The level of fatty acid-binding protein 4, a novel adipokine, is increased in rheumatoid arthritis and correlates with serum cholesterol levels. <i>Cytokine</i> , 2013, 64, 441-447.	1.4	19
57	No effect of physiotherapy on the serum levels of adipocytokines in patients with ankylosing spondylitis. <i>Clinical Rheumatology</i> , 2012, 31, 67-71.	1.0	18
58	Identification of Novel Native Autoantigens in Rheumatoid Arthritis. <i>Biomedicines</i> , 2020, 8, 141.	1.4	18
59	Interleukin-35 is upregulated in systemic sclerosis and its serum levels are associated with early disease. <i>Rheumatology</i> , 2015, 54, kev260.	0.9	17
60	Pro-inflammatory S100A11 is elevated in inflammatory myopathies and reflects disease activity and extramuscular manifestations in myositis. <i>Cytokine</i> , 2019, 116, 13-20.	1.4	17
61	S100A11 (calgizzarin) is released via NETosis in rheumatoid arthritis (RA) and stimulates IL-6 and TNF secretion by neutrophils. <i>Scientific Reports</i> , 2021, 11, 6063.	1.6	17
62	Chronic repetitive nonprogressive epilepsy partialis continua due to rheumatoid meningitis. <i>Seizure: the Journal of the British Epilepsy Association</i> , 2013, 22, 80-82.	0.9	16
63	Expanding the citrullinome of synovial fibrinogen from rheumatoid arthritis patients. <i>Journal of Proteomics</i> , 2019, 208, 103484.	1.2	16
64	Cytokines and inflammatory mediators as promising markers of polymyositis/dermatomyositis. <i>Current Opinion in Rheumatology</i> , 2020, 32, 534-541.	2.0	16
65	Applicability of Small-Molecule Inhibitors in the Study of Peptidyl Arginine Deiminase 2 (PAD2) and PAD4. <i>Frontiers in Immunology</i> , 2021, 12, 716250.	2.2	16
66	The Role of Resistin in Inflammatory Myopathies. <i>Current Rheumatology Reports</i> , 2013, 15, 336.	2.1	15
67	Serum tenascin-C discriminates patients with active SLE from inactive patients and healthy controls and predicts the need to escalate immunosuppressive therapy: a cohort study. <i>Arthritis Research and Therapy</i> , 2015, 17, 341.	1.6	14
68	Biomarkers of hand osteoarthritis. <i>Rheumatology International</i> , 2018, 38, 725-735.	1.5	14
69	Serum Calprotectin Discriminates Subclinical Disease Activity from Ultrasound-Defined Remission in Patients with Rheumatoid Arthritis in Clinical Remission. <i>PLoS ONE</i> , 2016, 11, e0165498.	1.1	14
70	The metastasis promoting protein S100A4 levels associate with disease activity rather than cancer development in patients with idiopathic inflammatory myopathies. <i>Arthritis Research and Therapy</i> , 2014, 16, 468.	1.6	13
71	High levels of metastasis-inducing S100A4 protein and treatment outcome in early rheumatoid arthritis: data from the PERAC cohort. <i>Biomarkers</i> , 2015, 20, 47-51.	0.9	12
72	Adipokines: role in local and systemic inflammation of rheumatic diseases. <i>Expert Review of Clinical Immunology</i> , 2017, 13, 1-3.	1.3	11

#	ARTICLE	IF	CITATIONS
73	Response to secukinumab on synovitis using Power Doppler ultrasound in psoriatic arthritis: 12-week results from a phase III study, ULTIMATE. <i>Rheumatology</i> , 2022, 61, 1867-1876.	0.9	11
74	IL-40: A New B Cell-Associated Cytokine Up-Regulated in Rheumatoid Arthritis Decreases Following the Rituximab Therapy and Correlates With Disease Activity, Autoantibodies, and NETosis. <i>Frontiers in Immunology</i> , 2021, 12, 745523.	2.2	11
75	Muscle and Fat Metabolism in Obesity After Kidney Transplantation: No Effect of Peritoneal Dialysis or Hemodialysis. <i>Journal of Intensive Care Medicine</i> , 2012, 22, 166-170.		10
76	Interleukin-20 is triggered by TLR ligands and associates with disease activity in patients with rheumatoid arthritis. <i>Cytokine</i> , 2017, 97, 187-192.	1.4	10
77	Clusterin serum levels are elevated in patients with early rheumatoid arthritis and predict disease activity and treatment response. <i>Scientific Reports</i> , 2021, 11, 11525.	1.6	10
78	Adipokine profile is modulated in subcutaneous adipose tissue by TNF α inhibitors in patients with rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, 2054-2056.	0.5	9
79	Cross-sectional study of patients with axial spondyloarthritis fulfilling imaging arm of ASAS classification criteria: baseline clinical characteristics and subset differences in a single-centre cohort. <i>BMJ Open</i> , 2019, 9, e024713.	0.8	9
80	Early changes in serum visfatin after abdominal surgery: a new pro-inflammatory marker in diagnosis?. <i>Biomedical Papers of the Medical Faculty of the University Palacky&#x0301;, Olomouc, Czechoslovakia</i> , 2015, 159, 489-496.	0.2	9
81	The effect of a 24-week training focused on activities of daily living, muscle strengthening, and stability in idiopathic inflammatory myopathies: a monocentric controlled study with follow-up. <i>Arthritis Research and Therapy</i> , 2021, 23, 173.	1.6	8
82	Real world use of secukinumab for treatment of axial spondyloarthritis and psoriatic arthritis: nationwide results from the ATTRA registry. <i>Clinical and Experimental Rheumatology</i> , 2019, 37, 342-343.	0.4	8
83	High miR-451 expression in peripheral blood mononuclear cells from subjects at risk of developing rheumatoid arthritis. <i>Scientific Reports</i> , 2021, 11, 4719.	1.6	7
84	Serum visfatin levels in patients with axial spondyloarthritis and their relationship to disease activity and spinal radiographic damage: a cross-sectional study. <i>Rheumatology International</i> , 2019, 39, 1037-1043.	1.5	6
85	Sexual function in patients with idiopathic inflammatory myopathies: a cross-sectional study. <i>Rheumatology</i> , 2021, 60, 5060-5072.	0.9	6
86	Pentosidine, an Advanced Glycation End-Product, May Reflect Clinical and Morphological Features of Hand Osteoarthritis. <i>Open Rheumatology Journal</i> , 2012, 6, 64-69.	0.1	6
87	Inhibition of Hsp90 Counteracts the Established Experimental Dermal Fibrosis Induced by Bleomycin. <i>Biomedicines</i> , 2021, 9, 650.	1.4	5
88	Identification of potential autoantigens in anti-CCP-positive and anti-CCP-negative rheumatoid arthritis using citrulline-specific protein arrays. <i>Scientific Reports</i> , 2021, 11, 17300.	1.6	5
89	Chemokine and Cytokine Profiles in Patients with Hand Osteoarthritis. <i>Biomolecules</i> , 2021, 11, 4.	1.8	5
90	Cardiovascular Risk in Myositis Patients Compared to the General Population: Preliminary Data From a Single-Center Cross-Sectional Study. <i>Frontiers in Medicine</i> , 2022, 9, .	1.2	5

#	ARTICLE	IF	CITATIONS
91	Interleukin-35 in idiopathic inflammatory myopathies. <i>Cytokine</i> , 2021, 137, 155350.	1.4	3
92	P187â€¦Secukinumab significantly decreased joint synovitis measured by Power Doppler ultrasonography in biologic-naïve patients with active psoriatic arthritis: primary (12week) results from a randomised, placebo-controlled Phase 3 study. <i>Rheumatology</i> , 2021, 60, .	0.9	3
93	Plasma heat shock protein 90 levels in patients with spondyloarthritis and their relation to structural changes: a cross-sectional study. <i>Biomarkers in Medicine</i> , 2021, 15, 5-13.	0.6	3
94	Female Sexual Dysfunction and Pelvic Floor Muscle Function Associated with Systemic Sclerosis: A Cross-Sectional Study. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 612.	1.2	3
95	Hsp90 Levels in Idiopathic Inflammatory Myopathies and Their Association With Muscle Involvement and Disease Activity: A Cross-Sectional and Longitudinal Study. <i>Frontiers in Immunology</i> , 2022, 13, 811045.	2.2	2
96	Increased visfatin levels are associated with higher disease activity in anti-Jo-1-positive myositis patients. <i>Clinical and Experimental Rheumatology</i> , 2016, 34, 222-9.	0.4	2
97	The link between metastasis-associated protein S100A4 and rheumatoid arthritis. <i>Joint Bone Spine</i> , 2008, 75, 242-243.	0.8	1
98	MiRs in RA: possible biomarkers and therapeutic targets. <i>Arthritis Research and Therapy</i> , 2012, 14, .	1.6	1
99	Calprotectin (a S100 protein) as a sensitive biomarker for rheumatoid arthritis: new perspectives for an old finding. <i>International Journal of Clinical Rheumatology</i> , 2012, 7, 127-129.	0.3	1
100	THU0528â€¦Decrease in Circulating Visfatin Levels is Associated with Disease Activity Improvement in Early Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 366.1-366.	0.5	1
101	A1.48â€¦Enhanced expression of CD11c on non-classical CD16+peripheral blood monocytes in early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, A20.2-A21.	0.5	1
102	A3.16â€¦Serum S100A4 correlates with skin fibrosis, lung involvement and disease activity in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, A48.1-A48.	0.5	1
103	AB0203â€¦S100A4 Serum Levels Correlate with Disease Activity, Skin Fibrosis and Lung Involvement in Systemic Sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 871.1-871.	0.5	1
104	A6.26â€¦Analysis of circulating mirnas in patients with axial spondyloarthritis with different spinal involvement. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, A66.1-A66.	0.5	1
105	A3.10â€¦Plasma levels of heat shock protein 90 correlate with disease activity, lung involvement and skin fibrosis in systemic sclerosis. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, A35.1-A35.	0.5	1
106	S100A4 is elevated in axial spondyloarthritis: a potential link to disease severity. <i>BMC Rheumatology</i> , 2020, 4, 13.	0.6	1
107	S100A11 (calgizzarin) is released by circulating mononuclear cells and its elevated plasma levels distinguish systemic lupus erythematosus patients from healthy individuals. <i>Clinical and Experimental Rheumatology</i> , 2019, 37, 338-339.	0.4	1
108	Clusterin is upregulated in serum and muscle tissue in idiopathic inflammatory myopathies and associates with clinical disease activity and cytokine profile. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 1021-1032.	0.4	1

#	ARTICLE	IF	CITATIONS
109	The effect of a 24-week physiotherapy and occupational therapy program in systemic sclerosis: a monocentric controlled study with follow-up. <i>Clinical and Experimental Rheumatology</i> , 0, , .	0.4	1
110	Title is missing!. <i>Arthritis Research</i> , 2003, 5, 167.	2.0	0
111	Metastasis-Promoting protein S100A4 (MTS1) upregulates production of proinflammatory cytokines via TLR-4. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, A38-A39.	0.5	0
112	Immune cell-derived microparticles in polymyositis/dermatomyositis. <i>Annals of the Rheumatic Diseases</i> , 2010, 69, A55-A56.	0.5	0
113	Modulation of subcutaneous adipose tissue adipokines by TNF- α blockade therapy in patients with inflammatory arthritides. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, A85-A85.	0.5	0
114	Potential role of S100A4 in an immune response of rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, A85-A85.	0.5	0
115	Metastasis-inducing S100A4 protein is associated with treatment response in patients with recent-onset rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, A15.2-A15.	0.5	0
116	A8.3â€¦Deficit of S100A4 Prevents Joint Destruction and Systemic Bone Loss in hTNFtg Mouse Model. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A58.1-A58.	0.5	0
117	A7.3â€¦Association of Circulating miR-223 and miR-16 with Disease Activity in Patients with Early Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A48.3-A49.	0.5	0
118	THU0168â€¦Monotherapy with Biological Disease-Modifying Anti-Rheumatic Drugs-Data from the Czech Registry Attra:. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 238.2-238.	0.5	0
119	OP0240â€¦Signal Transducer and Activator of Transcription 3 Regulates Fibroblast Activation as A Downstream Mediator of Transforming Growth Factor-Beta. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 153.1-153.	0.5	0
120	AB0114â€¦Comparative Analysis of the Expression of Protein Tyrosine Phosphatase Non-Receptor Type 2 (PTPN2) in Autoimmune Disease. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 841.3-842.	0.5	0
121	SAT0205â€¦Potential of US 7 Score in Evaluating of Disease Activity of Patients with Rheumatoid Arthritis in the State of Remission. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 664.1-664.	0.5	0
122	AB0163â€¦Comprehensive Analysis of Circulating Mirnas in Patients with Axial Spondyloarthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 856.4-857.	0.5	0
123	AB0818â€¦Bone Mineral Density and Metabolic Risk Factors in Patients with Non-Erosive and Erosive Hand Osteoarthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 1073.3-1074.	0.5	0
124	SAT0565â€¦The Expression and Regulation of the Argonaute Protein Family Member PIWIL4 in RA. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 795.1-795.	0.5	0
125	SAT0188â€¦Serum Calprotectin (S100A8/9) Correlates with Clinical and Ultrasound Outcomes in Patients with Early Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 658.1-658.	0.5	0
126	A3.10â€¦Serum calprotectinis elevated in patients with early rheumatoid arthritis but not in patients at risk of developing rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, A36.1-A36.	0.5	0

#	ARTICLE	IF	CITATIONS
127	SAT0466â€¦Progression of Pain and Ultrasound Detected Synovitis in Patients with Erosive Hand Osteoarthritis over One Year. Annals of the Rheumatic Diseases, 2016, 75, 839.3-840.	0.5	0
128	AB0952â€¦Serum Visfatin and Resistin, but Not Adiponectin or Leptin, Are Associated with Ultrasound Synovitis in Patients with Rheumatoid Arthritis. Annals of the Rheumatic Diseases, 2016, 75, 1227.1-1227.	0.5	0
129	A9.02â€¦Heat shock protein 90 plasma levels correlate with disease activity, skeletal muscle, lung and heart involvement in idiopathic inflammatory myopathies. Annals of the Rheumatic Diseases, 2016, 75, A70.2-A71.	0.5	0
130	A10.17â€¦Interleukin-20 serum levels correlate with disease activity in rheumatoid arthritis patients and decrease in response to rituximab treatment. Annals of the Rheumatic Diseases, 2016, 75, A79.2-A79.	0.5	0
131	THU0353â€¦Serum Tweak Levels in Patients with Neuropsychiatric Systemic Lupus Erythematosus. Annals of the Rheumatic Diseases, 2016, 75, 315.1-315.	0.5	0
132	SAT0068â€¦Circulating MIR-126 Is A Marker of Disease Activity in Patients with Early RA but Does Not Differentiate Patients at Risk of Developing RA. Annals of the Rheumatic Diseases, 2016, 75, 689.2-689.	0.5	0
133	AB0696â€¦Clinical Characteristics of Spondyloarthritis in The Czech Republic: A Single Centre Cohort. Annals of the Rheumatic Diseases, 2016, 75, 1142.2-1143.	0.5	0
134	A8.06â€¦MicroRNA-125b expression in PBMCS is inversely associated with disease activity in patients with early rheumatoid arthritis. Annals of the Rheumatic Diseases, 2016, 75, A67.1-A67.	0.5	0
135	AB0599â€¦Interleukin-35 in Idiopathic Inflammatory Myopathies. Annals of the Rheumatic Diseases, 2016, 75, 1109.2-1109.	0.5	0
136	08.48â€¦Increased body fat but decreased lean body mass and bone mineral density in myositis patients are associated with disease duration, inflammatory status, skeletal muscle involvement and physical activity. , 2017, , .		0
137	08.49â€¦Efficacy of an intensive 24-week physiotherapy programme in scleroderma patients â€“ preliminary data from a single-centre controlled study. , 2017, , .		0
138	08.47â€¦Decreased body fat, lean body mass and bone mineral density in scleroderma patients are associated with disease activity and physical activity. , 2017, , .		0
139	02.30â€¦High levels of mir-451a differentiate patients with clinically suspect arthralgia from healthy controls. , 2017, , .		0
140	AB0687â€¦EFFECTIVENESS OF SPECIALIZED AND INTENSIVE ADL TRAINING IN PATIENTS WITH IDIOPATHIC INFLAMMATORY MYOPATHIES â€“ PRELIMINARY RESULTS OF A ONE-YEAR CONTROLLED STUDY. , 2019, , .		0
141	FRI0107â€¦THE EFFECT OF CO-MEDICATION WITH METHOTREXATE OR OTHER CONVENTIONAL SYNTHETIC DISEASE MODIFYING ANTI-RHEUMATIC DRUGS ON FIRST TUMOR NECROSIS INHIBITOR DRUG SURVIVAL IN PATIENTS WITH RHEUMATOID ARTHRITIS: DATA FROM THE CZECH ATTRA REGISTRY. , 2019, , .		0
142	OP0066â€¦EFFECTIVENESS OF SPECIALIZED HAND/FACE PHYSICAL-OCCUPATIONAL THERAPY IN PATIENTS WITH SYSTEMIC SCLEROSIS â€“ PRELIMINARY RESULTS OF A ONE-YEAR CONTROLLED STUDY. , 2019, , .		0
143	THU0062â€¦THE DYSREGULATION OF NK CELLS AND NON-CLASSICAL AND CLASSICAL MONOCYTE SUBPOPULATIONS IN INDIVIDUALS AT RISK OF DEVELOPING RHEUMATOID ARTHRITIS. , 2019, , .		0
144	The effect of a 24-week physiotherapy and occupational therapy program in systemic sclerosis: a monocentric controlled study with follow-up.. Clinical and Experimental Rheumatology, 2022, , .	0.4	0