

Hartmut R StÃ¼tzel

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2401863/publications.pdf>

Version: 2024-02-01

91
papers

2,372
citations

201674

27
h-index

243625

44
g-index

94
all docs

94
docs citations

94
times ranked

2758
citing authors

#	ARTICLE	IF	CITATIONS
1	Biomass partitioning, specific leaf area, and water use efficiency of vegetable amaranth (<i>Amaranthus</i>) Tj ETQq1 1 0,784314 rgBT /Ove	3.6	225
2	Breeding improves wheat productivity under contrasting agrochemical input levels. <i>Nature Plants</i> , 2019, 5, 706-714.	9.3	194
3	Root growth and dry matter partitioning of cauliflower under drought stress conditions: measurement and simulation. <i>European Journal of Agronomy</i> , 2004, 20, 379-394.	4.1	121
4	Ontogenetic Changes of 2-Propenyl and 3-Indolylmethyl Glucosinolates in <i>Brassica carinata</i> Leaves as Affected by Water Supply. <i>Journal of Agricultural and Food Chemistry</i> , 2009, 57, 7259-7263.	5.2	85
5	Glucosinolate Concentration in Turnip (<i>Brassica rapa</i> ssp. <i>rapifera</i> L.) Roots as Affected by Nitrogen and Sulfur Supply. <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 8452-8457.	5.2	81
6	Leaf water relations of vegetable amaranth (<i>Amaranthus</i> spp.) in response to soil drying. <i>European Journal of Agronomy</i> , 2002, 16, 137-150.	4.1	66
7	Water supply and growing season influence glucosinolate concentration and composition in turnip root (<i>Brassica rapa</i> ssp. <i>rapifera</i> L.). <i>Journal of Plant Nutrition and Soil Science</i> , 2008, 171, 255-265.	1.9	66
8	Leaf Expansion, Stomatal Conductance, and Transpiration of Vegetable Amaranth (<i>Amaranthus</i> sp.) in Response to Soil Drying. <i>Journal of the American Society for Horticultural Science</i> , 2002, 127, 878-883.	1.0	63
9	Determining the stable Fe isotope signature of plant-available iron in soils. <i>Chemical Geology</i> , 2010, 277, 269-280.	3.3	60
10	Disentangling the contributions of osmotic and ionic effects of salinity on stomatal, mesophyll, biochemical and light limitations to photosynthesis. <i>Plant, Cell and Environment</i> , 2015, 38, 1528-1542.	5.7	51
11	Modelling photo-modulated internode elongation in growing glasshouse cucumber canopies. <i>New Phytologist</i> , 2011, 190, 697-708.	7.3	49
12	Crop model based QTL analysis across environments and QTL based estimation of time to floral induction and flowering in <i>Brassica oleracea</i> . <i>Molecular Breeding</i> , 2008, 21, 205-216.	2.1	45
13	What is the most prominent factor limiting photosynthesis in different layers of a greenhouse cucumber canopy?. <i>Annals of Botany</i> , 2014, 114, 677-688.	2.9	45
14	Simulation of faba bean (<i>Vicia faba</i> L.) root system development under Mediterranean conditions. <i>European Journal of Agronomy</i> , 1998, 9, 259-272.	4.1	44
15	A new method to estimate photosynthetic parameters through net assimilation rate~intercellular space CO_2 concentration (A/C_i ~ C_i) curve and chlorophyll fluorescence measurements. <i>New Phytologist</i> , 2017, 213, 1543-1554.	7.3	44
16	Quantification of the effects of architectural traits on dry mass production and light interception of tomato canopy under different temperature regimes using a dynamic functional~structural plant model. <i>Journal of Experimental Botany</i> , 2014, 65, 6399-6410.	4.8	42
17	A new method for assessing the sustainability of land-use systems (II): Evaluating impact indicators. <i>Ecological Economics</i> , 2009, 68, 1288-1300.	5.7	41
18	Title is missing!. <i>Plant and Soil</i> , 2000, 223, 133-147.	3.7	40

#	ARTICLE	IF	CITATIONS
19	Modelling leaf phototropism in a cucumber canopy. <i>Functional Plant Biology</i> , 2008, 35, 876.	2.1	39
20	Simulation of faba bean (<i>Vicia faba</i> L.) growth and development under Mediterranean conditions: Model adaptation and evaluation. <i>European Journal of Agronomy</i> , 1998, 9, 273-293.	4.1	35
21	Differences in the enzymatic hydrolysis of glucosinolates increase the defense metabolite diversity in 19 <i>Arabidopsis thaliana</i> accessions. <i>Plant Physiology and Biochemistry</i> , 2018, 124, 126-135.	5.8	35
22	Decoupling of impact factors reveals the response of German winter wheat yields to climatic changes. <i>Global Change Biology</i> , 2020, 26, 3601-3626.	9.5	35
23	A new method for assessing the sustainability of land-use systems (I): Identifying the relevant issues. <i>Ecological Economics</i> , 2009, 68, 1275-1287.	5.7	33
24	Genetic dissection of temperature-dependent sorghum growth during juvenile development. <i>Theoretical and Applied Genetics</i> , 2014, 127, 1935-1948.	3.6	32
25	Prediction of flowering time in <i>Brassica oleracea</i> using a quantitative trait loci-based phenology model. <i>Plant Biology</i> , 2012, 14, 179-189.	3.8	31
26	Title is missing!. <i>Plant and Soil</i> , 2002, 246, 201-209.	3.7	30
27	Evaluation of a radiosity based light model for greenhouse cucumber canopies. <i>Agricultural and Forest Meteorology</i> , 2011, 151, 906-915.	4.8	30
28	Biodiversity in European agricultural landscapes: transformative societal changes needed. <i>Trends in Ecology and Evolution</i> , 2021, 36, 1067-1070.	8.7	29
29	A simple empirical model for predicting development and dry matter partitioning in cauliflower (<i>Brassica oleracea</i> L. botrytis). <i>Scientia Horticulturae</i> , 1999, 80, 19-38.	3.6	28
30	A Three-Dimensional Approach to Modeling Light Interception in Heterogeneous Canopies. <i>Agronomy Journal</i> , 1999, 91, 1024-1032.	1.8	28
31	Genetic dissection of the temperature dependent emergence processes in sorghum using a cumulative emergence model and stability parameters. <i>Theoretical and Applied Genetics</i> , 2012, 125, 1647-1661.	3.6	25
32	Quantitative aspects of <i>Orobanche crenata</i> infestation in faba beans as affected by abiotic factors and parasite soil seedbank. <i>Weed Research</i> , 2001, 41, 311-324.	1.7	24
33	High light aggravates functional limitations of cucumber canopy photosynthesis under salinity. <i>Annals of Botany</i> , 2018, 121, 797-807.	2.9	23
34	Co-Evolution of Sink and Source in the Recent Breeding History of Winter Wheat in Germany. <i>Frontiers in Plant Science</i> , 2019, 10, 1771.	3.6	23
35	Quantitative trait loci controlling leaf appearance and curd initiation of cauliflower in relation to temperature. <i>Theoretical and Applied Genetics</i> , 2016, 129, 1273-1288.	3.6	21
36	Determining Ion Toxicity in Cucumber under Salinity Stress. <i>Agronomy</i> , 2020, 10, 677.	3.0	21

#	ARTICLE	IF	CITATIONS
37	Dry matter partitioning models for the simulation of individual fruit growth in greenhouse cucumber canopies. <i>Annals of Botany</i> , 2011, 108, 1075-1084.	2.9	20
38	A model for light competition between vegetable crops and weeds. <i>European Journal of Agronomy</i> , 2001, 14, 13-29.	4.1	19
39	A Model of the Partitioning of New Above-ground Dry Matter. <i>Annals of Botany</i> , 1988, 61, 481-487.	2.9	18
40	Modelling Nitrogen Content and Distribution in Cauliflower (<i>Brassica oleracea</i> L. botrytis). <i>Annals of Botany</i> , 2000, 86, 963-973.	2.9	18
41	Predicting dry matter production of cauliflower (<i>Brassica oleracea</i> L. botrytis) under unstressed conditions. <i>Scientia Horticulturae</i> , 2001, 87, 171-190.	3.6	18
42	Plant growth, water relations and transpiration of two species of African nightshade (<i>Solanum</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 54 water-limited conditions. <i>Scientia Horticulturae</i> , 2006, 110, 7-15.	3.6	18
43	Environmental triggers for photosynthetic protein turnover determine the optimal nitrogen distribution and partitioning in the canopy. <i>Journal of Experimental Botany</i> , 2019, 70, 2419-2434.	4.8	18
44	High temperature and vapor pressure deficit aggravate architectural effects but ameliorate non-architectural effects of salinity on dry mass production of tomato. <i>Frontiers in Plant Science</i> , 2015, 6, 887.	3.6	17
45	Optimal Nitrogen Content and Photosynthesis in Cauliflower (<i>Brassica oleracea</i> L. botrytis). Scaling up from a Leaf to the Whole Plant. <i>Annals of Botany</i> , 2000, 85, 779-787.	2.9	16
46	Modelling the effects of soil water limitations on transpiration and stomatal regulation of cauliflower. <i>European Journal of Agronomy</i> , 2007, 26, 375-383.	4.1	16
47	Plant Growth, Water Relations, and Transpiration of Spiderplant [<i>Gynandropsis gynandra</i> (L.) Briq.] under Water-limited Conditions. <i>Journal of the American Society for Horticultural Science</i> , 2005, 130, 469-477.	1.0	16
48	Estimation of Geometric Attributes and Masses of Individual Cucumber Organs Using Three-dimensional Digitizing and Allometric Relationships. <i>Journal of the American Society for Horticultural Science</i> , 2007, 132, 439-446.	1.0	16
49	Simplification of a light-based model for estimating final internode length in greenhouse cucumber canopies. <i>Annals of Botany</i> , 2011, 108, 1055-1063.	2.9	14
50	The Future of Field Trials in Europe: Establishing a Network Beyond Boundaries. <i>Trends in Plant Science</i> , 2016, 21, 92-95.	8.8	14
51	The physiological causes of mixing effects in cultivar mixtures: A general hypothesis. <i>Agricultural Systems</i> , 1990, 32, 41-53.	6.1	13
52	Prediction of time to harvest and its variability in broccoli (<i>Brassica oleracea</i> var. <i>italica</i>) Part I. Plant developmental variation and forecast of time to head induction. <i>Scientia Horticulturae</i> , 2016, 198, 424-433.	3.6	13
53	Dry Matter Partitioning in a Determinate and an Indeterminate Cultivar of <i>Vicia faba</i> L. Under Contrasting Plant Distributions and Densities. <i>Annals of Botany</i> , 1991, 67, 487-495.	2.9	12
54	Dry matter production and partitioning of <i>Chenopodium album</i> in contrasting competitive environments. <i>Weed Research</i> , 2001, 41, 129-142.	1.7	12

#	ARTICLE	IF	CITATIONS
55	Aspects of nitrogen use efficiency of cauliflower II. Productivity and nitrogen partitioning as influenced by N supply. <i>Journal of Agricultural Science</i> , 2003, 141, 17-29.	1.3	12
56	Nitrogen use efficiency of organically fertilized white cabbage and residual effects on subsequent beetroot. <i>Plant and Soil</i> , 2014, 382, 237-251.	3.7	12
57	Irrigation Scheduling of Kohlrabi (<i>Brassica oleracea</i> var. <i>gongylodes</i>) Using Crop Water Stress Index. <i>Hortscience: A Publication of the American Society for Horticultural Science</i> , 2004, 39, 276-279.	1.0	12
58	Canopy development of a determinate and an indeterminate cultivar of <i>Vicia faba</i> L. under contrasting plant distributions and densities. <i>Annals of Applied Biology</i> , 1991, 118, 185-199.	2.5	11
59	A Modeling Approach to Quantify the Effects of Stomatal Behavior and Mesophyll Conductance on Leaf Water Use Efficiency. <i>Frontiers in Plant Science</i> , 2016, 7, 875.	3.6	11
60	A mechanistic view of the reduction in photosynthetic protein abundance under diurnal light fluctuation. <i>Journal of Experimental Botany</i> , 2019, 70, 3705-3708.	4.8	10
61	Yield determinants of Guar (<i>Cyamopsis tetragonoloba</i>): 1. Grain yield and pod number. <i>Field Crops Research</i> , 1989, 21, 29-37.	5.1	9
62	Prediction of time to harvest and its variability of broccoli (<i>Brassica oleracea</i> var. <i>italica</i>) part II. Growth model description, parameterisation and field evaluation. <i>Scientia Horticulturae</i> , 2016, 200, 151-160.	3.6	9
63	Modeling the Effects of Drought Stress on Leaf Development in a <i>Brassica oleracea</i> Doubled Haploid Population Using Two-phase Linear Functions. <i>Journal of the American Society for Horticultural Science</i> , 2009, 134, 543-552.	1.0	9
64	Prediction of winter wheat cultivar performance in Germany: At national, regional and location scale. <i>European Journal of Agronomy</i> , 2014, 52, 210-217.	4.1	8
65	Physiological and morphological responses of different spring barley genotypes to water deficit and associated QTLs. <i>PLoS ONE</i> , 2020, 15, e0237834.	2.5	8
66	Changes in plant growth, leaf relative water content and physiological traits in response to salt stress in peanut (<i>Arachis hypogaea</i> L.) varieties. <i>Notulae Botanicae Horti Agrobotanici Cluj-Napoca</i> , 2021, 49, 12049.	1.1	8
67	Physiological and Morphological Responses of Cassava Genotypes to Fertilization Regimes in Chromi-Haplic Acrisols Soils. <i>Agronomy</i> , 2021, 11, 1757.	3.0	8
68	Unraveling the genetic complexity underlying sorghum response to water availability. <i>PLoS ONE</i> , 2019, 14, e0215515.	2.5	7
69	Seasonal Efficiency of Supplemental LED Lighting on Growth and Photomorphogenesis of Sweet Basil. <i>Frontiers in Plant Science</i> , 2021, 12, 609975.	3.6	7
70	Effects of winter barley cultivar mixtures on lodging. <i>Journal of Agricultural Science</i> , 1989, 112, 47-55.	1.3	6
71	Canopy development of <i>Chenopodium album</i> in pure and mixed stands. <i>Weed Research</i> , 2001, 41, 111-128.	1.7	6
72	Nitrogen efficiency of Brussels sprouts under different organic N fertilization rates. <i>Scientia Horticulturae</i> , 2012, 134, 7-12.	3.6	6

#	ARTICLE	IF	CITATIONS
73	Editorial: Virtual Plants: Modeling Plant Architecture in Changing Environments. <i>Frontiers in Plant Science</i> , 2016, 7, 1734.	3.6	6
74	How does structure matter? Comparison of canopy photosynthesis using one- and three-dimensional light models: a case study using greenhouse cucumber canopies. <i>In Silico Plants</i> , 2021, 3, .	1.9	6
75	Sorten-Mischungseffekte in Wintergerstenbeständen in Abhängigkeit von Standort und Produktionsintensität. <i>Journal of Agronomy and Crop Science</i> , 1989, 162, 180-191.	3.5	4
76	Decomposition of lupine seeds and seedlings as N fertilizer in organic vegetable production. <i>Plant and Soil</i> , 2012, 357, 59-71.	3.7	3
77	Narrow-leaved lupine as an N source alternative to grass-clover swards in organic vegetable rotations. <i>Biological Agriculture and Horticulture</i> , 2017, 33, 125-142.	1.0	3
78	Impact of the source of organic manure on persistence of <i>E. coli</i> O157:H7 gfp + in rocket (<i>Diplotaxis</i>) Tj ETQq0 0 0 gBT /Overlock 10 T	3.5	3
79	Experiments for in silico evaluation of Optimality of Photosynthetic Nitrogen Distribution and Partitioning in the Canopy: an Example Using Greenhouse Cucumber Plants. <i>Bio-protocol</i> , 2020, 10, e3556.	0.4	3
80	Effekte der Sorten- (Weizen) und der Arten- (Weizen, Roggen) Mischung auf die Ertragsleistung krankheitsfreier Bestände. <i>Journal of Agronomy and Crop Science</i> , 1989, 163, 319-329.	3.5	2
81	Nitrogen Status and Light Environment Influence Dry Matter Partitioning in Cauliflower. <i>Journal of the American Society for Horticultural Science</i> , 2001, 126, 750-756.	1.0	2
82	Performance of Cassava under Lime, Fertilizer, and Legume Intercropping on Exhausted Land in Northern Zambia. <i>International Journal of Agronomy</i> , 2022, 2022, 1-17.	1.2	2
83	Grain Yield of Intercropped Sorghum and Pearl Millet as Influenced by Sorghum Genotype and Cropping Pattern. <i>Journal of Agronomy and Crop Science</i> , 1988, 160, 191-197.	3.5	1
84	Yield determinants of guar (<i>Cyamopsis tetragonoloba</i>): 2. Nitrogen accumulation and growth at high plant density. <i>Field Crops Research</i> , 1989, 21, 39-47.	5.1	1
85	Lichtaufnahme und Stoffproduktion eines konventionellen und eines epigonalen Genotyps der Weißen Lupine (<i>Lupinus albus</i>). <i>Journal of Agronomy and Crop Science</i> , 1993, 171, 1-12.	3.5	1
86	Predicting dry-matter partitioning between individual cauliflower leaves using a source limitation/sink hierarchy model. <i>Journal of Horticultural Science and Biotechnology</i> , 2003, 78, 537-548.	1.9	1
87	Modeling temperature-modulated stem growth of cucumber plants (<i>Cucumis sativus</i> L.), 2012, , .		1
88	Determining photosynthetic limitations under saturated and non-saturated light conditions. , 2012, , .		1
89	Interspecific variation in leaf traits, photosynthetic light response, and whole-plant productivity in amaranths (<i>Amaranthus</i> spp. L.). <i>PLoS ONE</i> , 2022, 17, e0270674.	2.5	1
90	Die Ertragsbildung und Ertragsstruktur eines konventionellen und eines epigonalen Genotyps der Weißen Lupine (<i>Lupinus albus</i>). <i>Journal of Agronomy and Crop Science</i> , 1993, 170, 177-186.	3.5	0

#	ARTICLE	IF	CITATIONS
91	First Form, Then Function: 3D Reconstruction of Cucumber Plants (<i>Cucumis sativus</i> L.) Allows Early Detection of Stress Effects through Leaf Dimensions. <i>Remote Sensing</i> , 2022, 14, 1094.	4.0	0