

Lars Henrik Jensen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2404099/publications.pdf>

Version: 2024-02-01

94
papers

3,906
citations

361413

20
h-index

138484

58
g-index

97
all docs

97
docs citations

97
times ranked

4868
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Pembrolizumab in Microsatellite-Instability-High Advanced Colorectal Cancer. <i>New England Journal of Medicine</i> , 2020, 383, 2207-2218. | 27.0 | 1,513 |
| 2 | High-dose chemoradiotherapy and watchful waiting for distal rectal cancer: a prospective observational study. <i>Lancet Oncology</i> , The, 2015, 16, 919-927. | 10.7 | 435 |
| 3 | Pembrolizumab versus chemotherapy for microsatellite instability-high or mismatch repair-deficient metastatic colorectal cancer (KEYNOTE-177): final analysis of a randomised, open-label, phase 3 study. <i>Lancet Oncology</i> , The, 2022, 23, 659-670. | 10.7 | 282 |
| 4 | Adjuvant chemotherapy with gemcitabine and cisplatin compared to observation after curative intent resection of cholangiocarcinoma and muscle invasive gallbladder carcinoma (ACTICCA-1 trial) - a randomized, multidisciplinary, multinational phase III trial. <i>BMC Cancer</i> , 2015, 15, 564. | 2.6 | 182 |
| 5 | Pembrolizumab versus chemotherapy for microsatellite instability-high/mismatch repair deficient metastatic colorectal cancer: The phase 3 KEYNOTE-177 Study.. <i>Journal of Clinical Oncology</i> , 2020, 38, LBA4-LBA4. | 1.6 | 150 |
| 6 | Health-related quality of life in patients with microsatellite instability-high or mismatch repair deficient metastatic colorectal cancer treated with first-line pembrolizumab versus chemotherapy (KEYNOTE-177): an open-label, randomised, phase 3 trial. <i>Lancet Oncology</i> , The, 2021, 22, 665-677. | 10.7 | 110 |
| 7 | Phase II marker-driven trial of panitumumab and chemotherapy in KRAS wild-type biliary tract cancer. <i>Annals of Oncology</i> , 2012, 23, 2341-2346. | 1.2 | 89 |
| 8 | Prognostic factors for progression-free and overall survival in advanced biliary tract cancer. <i>Annals of Oncology</i> , 2016, 27, 134-140. | 1.2 | 88 |
| 9 | Neoadjuvant chemotherapy in locally advanced colon cancer. A phase II trial. <i>Acta OncolÃ³gica</i> , 2015, 54, 1747-1753. | 1.8 | 84 |
| 10 | Final overall survival for the phase III KN177 study: Pembrolizumab versus chemotherapy in microsatellite instability-high/mismatch repair deficient (MSI-H/dMMR) metastatic colorectal cancer (mCRC).. <i>Journal of Clinical Oncology</i> , 2021, 39, 3500-3500. | 1.6 | 51 |
| 11 | Randomized cross-over study of patient preference for oral or intravenous vinorelbine in combination with carboplatin in the treatment of advanced NSCLC. <i>Lung Cancer</i> , 2008, 62, 85-91. | 2.0 | 50 |
| 12 | Strategy in clinical practice for classification of unselected colorectal tumours based on mismatch repair deficiency. <i>Colorectal Disease</i> , 2008, 10, 490-497. | 1.4 | 39 |
| 13 | KEYNOTE-177: Phase III randomized study of pembrolizumab versus chemotherapy for microsatellite instability-high advanced colorectal cancer.. <i>Journal of Clinical Oncology</i> , 2021, 39, 6-6. | 1.6 | 39 |
| 14 | An Update on Immune Checkpoint Therapy for the Treatment of Lynch Syndrome. <i>Clinical and Experimental Gastroenterology</i> , 2021, Volume 14, 181-197. | 2.3 | 36 |
| 15 | Tumor-stroma ratio predicts recurrence in patients with colon cancer treated with neoadjuvant chemotherapy. <i>Acta OncolÃ³gica</i> , 2018, 57, 528-533. | 1.8 | 36 |
| 16 | Long-Term Patient-Reported Outcomes After High-Dose Chemoradiation Therapy for Nonsurgical Management of Distal Rectal Cancer. <i>International Journal of Radiation Oncology Biology Physics</i> , 2020, 106, 556-563. | 0.8 | 32 |
| 17 | EGF61A>G polymorphism as predictive marker of clinical outcome to first-line capecitabine and oxaliplatin in metastatic colorectal cancer. <i>Annals of Oncology</i> , 2010, 21, 535-539. | 1.2 | 30 |
| 18 | Prognostic impact of CDX2 in stage II colon cancer: results from two nationwide cohorts. <i>British Journal of Cancer</i> , 2018, 119, 1367-1373. | 6.4 | 30 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Early identification of treatment benefit by methylated circulating tumor DNA in metastatic colorectal cancer. <i>Therapeutic Advances in Medical Oncology</i> , 2020, 12, 175883592091847. | 3.2 | 26 |
| 20 | Landmark survival analysis and impact of anatomic site of origin in prospective clinical trials of biliary tract cancer. <i>Journal of Hepatology</i> , 2020, 73, 1109-1117. | 3.7 | 25 |
| 21 | A phase II study of retifanlimab (INCMGA00012) in patients with squamous carcinoma of the anal canal who have progressed following platinum-based chemotherapy (POD1UM-202). <i>ESMO Open</i> , 2022, 7, 100529. | 4.5 | 23 |
| 22 | Local staging of sigmoid colon cancer using MRI. <i>Acta Radiologica Open</i> , 2017, 6, 205846011772095. | 0.6 | 22 |
| 23 | Serum IL6 as a Prognostic Biomarker and IL6R as a Therapeutic Target in Biliary Tract Cancers. <i>Clinical Cancer Research</i> , 2020, 26, 5655-5667. | 7.0 | 21 |
| 24 | A new model of early, integrated palliative care: palliative rehabilitation for newly diagnosed patients with non-resectable cancer. <i>Supportive Care in Cancer</i> , 2019, 27, 3291-3300. | 2.2 | 20 |
| 25 | Early ctDNA response to chemotherapy. A potential surrogate marker for overall survival. <i>European Journal of Cancer</i> , 2021, 149, 128-133. | 2.8 | 20 |
| 26 | Decline in CA19-9 during chemotherapy predicts survival in four independent cohorts of patients with inoperable bile duct cancer. <i>European Journal of Cancer</i> , 2015, 51, 1381-1388. | 2.8 | 19 |
| 27 | Early, integrated palliative rehabilitation improves quality of life of patients with newly diagnosed advanced cancer: The Pal-Rehab randomized controlled trial. <i>Palliative Medicine</i> , 2021, 35, 1344-1355. | 3.1 | 19 |
| 28 | Regulation of MLH1 mRNA and protein expression by promoter methylation in primary colorectal cancer: a descriptive and prognostic cancer marker study. <i>Cellular Oncology (Dordrecht)</i> , 2013, 36, 411-419. | 4.4 | 18 |
| 29 | A Phase Iâ€”II dose escalation study of fixed-dose rate gemcitabine, oxaliplatin and capecitabine every two weeks in advanced cholangiocarcinomas. <i>Acta OncolÃ³gica</i> , 2011, 50, 448-454. | 1.8 | 17 |
| 30 | A parallel-group randomized clinical trial of individually tailored, multidisciplinary, palliative rehabilitation for patients with newly diagnosed advanced cancer: the Pal-Rehab study protocol. <i>BMC Cancer</i> , 2017, 17, 560. | 2.6 | 17 |
| 31 | NPY Gene Methylation as a Universal, Longitudinal Plasma Marker for Evaluating the Clinical Benefit from Last-Line Treatment with Regorafenib in Metastatic Colorectal Cancer. <i>Cancers</i> , 2019, 11, 1649. | 3.7 | 17 |
| 32 | Treatment of Patients with Advanced Biliary Tract Cancer with Either Oxaliplatin, Gemcitabine, and Capecitabine or Cisplatin and Gemcitabineâ€”A Randomized Phase II Trial. <i>Cancers</i> , 2020, 12, 1975. | 3.7 | 17 |
| 33 | Drug Repositioning Based on the Reversal of Gene Expression Signatures Identifies TOP2A as a Therapeutic Target for Rectal Cancer. <i>Cancers</i> , 2021, 13, 5492. | 3.7 | 17 |
| 34 | Systemic therapy in younger and elderly patients with advanced biliary cancer: sub-analysis of ABC-02 and twelve other prospective trials. <i>BMC Cancer</i> , 2017, 17, 262. | 2.6 | 16 |
| 35 | How participatory action research changed our view of the challenges of shared decision-making training. <i>Patient Education and Counseling</i> , 2018, 101, 639-646. | 2.2 | 16 |
| 36 | Molecular biology from bench-to-bedside â€” Which colorectal cancer patients should be referred for genetic counselling and risk assessment. <i>European Journal of Cancer</i> , 2010, 46, 1823-1828. | 2.8 | 15 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | The prognostic value of simultaneous tumor and serum <sc>RAS</sc>/<sc>RAF</sc> mutations in localized colon cancer. <i>Cancer Medicine</i> , 2017, 6, 928-936. | 2.8 | 15 |
| 38 | Correlation Between Natural Killer Cell Activity and Treatment Effect in Patients with Disseminated Cancer. <i>Translational Oncology</i> , 2019, 12, 968-972. | 3.7 | 15 |
| 39 | Radiotherapy for metastatic spinal cord compression with increased radiation doses (RAMSES-01): a prospective multicenter study. <i>BMC Cancer</i> , 2019, 19, 1163. | 2.6 | 14 |
| 40 | Predictive Value of MSH2 Gene Expression in Colorectal Cancer Treated with Capecitabine. <i>Clinical Colorectal Cancer</i> , 2007, 6, 433-435. | 2.3 | 13 |
| 41 | Open dialogue about complementary and alternative medicine (CAM) integrated in conventional oncology care, characteristics and impact. A systematic review. <i>Patient Education and Counseling</i> , 2020, 103, 2224-2234. | 2.2 | 13 |
| 42 | Study protocol designed to investigate tumour response to calcium electroporation in cancers affecting the skin: a non-randomised phase II clinical trial. <i>BMJ Open</i> , 2021, 11, e046779. | 1.9 | 13 |
| 43 | Microsatellite Instability and the Association with Plasma Homocysteine and Thymidylate Synthase in Colorectal Cancer. <i>Cancer Investigation</i> , 2008, 26, 583-589. | 1.3 | 12 |
| 44 | Clinical outcome in 520 consecutive Danish rectal cancer patients treated with short course preoperative radiotherapy. <i>European Journal of Surgical Oncology</i> , 2010, 36, 237-243. | 1.0 | 12 |
| 45 | Laser microdissection and microsatellite analysis of colorectal adenocarcinomas. <i>Anticancer Research</i> , 2006, 26, 2069-74. | 1.1 | 12 |
| 46 | Combining biological agents and chemotherapy in the treatment of cholangiocarcinoma. <i>Expert Review of Anticancer Therapy</i> , 2011, 11, 589-600. | 2.4 | 10 |
| 47 | Lynch syndrome-associated epithelial ovarian cancer and its immunological profile. <i>Gynecologic Oncology</i> , 2021, 162, 686-693. | 1.4 | 10 |
| 48 | Improving continuity by bringing the cancer patient, general practitioner and oncologist together in a shared video-based consultation " protocol for a randomised controlled trial. <i>BMC Family Practice</i> , 2019, 20, 86. | 2.9 | 9 |
| 49 | Plasma Dynamics of RAS/RAF Mutations in Patients With Metastatic Colorectal Cancer Receiving Chemotherapy and Anti-EGFR Treatment. <i>Clinical Colorectal Cancer</i> , 2019, 18, 28-33.e3. | 2.3 | 9 |
| 50 | Cross-sectoral video consultations in cancer care: perspectives of cancer patients, oncologists and general practitioners. <i>Supportive Care in Cancer</i> , 2021, 29, 107-116. | 2.2 | 9 |
| 51 | The relationship between serum vascular endothelial growth factor A and microsatellite instability in colorectal cancer. <i>Colorectal Disease</i> , 2011, 13, 984-988. | 1.4 | 8 |
| 52 | Randomized <sc>Phase II</sc> trial of combination chemotherapy with panitumumab or bevacizumab for patients with inoperable biliary tract cancer without <sc>KRAS</sc> exon 2 mutations. <i>International Journal of Cancer</i> , 2021, 149, 119-126. | 5.1 | 8 |
| 53 | Molecular Screening for Lynch Syndrome: From Bench to Bedside. <i>Journal of Clinical Oncology</i> , 2009, 27, e224-e224. | 1.6 | 7 |
| 54 | Prognostic significance of circulating intact and cleaved forms of urokinase plasminogen activator receptor in inoperable chemotherapy treated cholangiocarcinoma patients. <i>Clinical Biochemistry</i> , 2014, 47, 599-604. | 1.9 | 7 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 55 | Correlation Between Tumor-Specific Mutated and Methylated DNA in Colorectal Cancer. <i>JCO Precision Oncology</i> , 2019, 3, 1-8. | 3.0 | 7 |
| 56 | Randomized phase II crossover trial exploring the clinical benefit from targeting EGFR or VEGF with combination chemotherapy in patients with non-resectable biliary tract cancer.. <i>Journal of Clinical Oncology</i> , 2015, 33, 4071-4071. | 1.6 | 7 |
| 57 | Investigating whether shared video-based consultations with patients, oncologists, and GPs can benefit patient-centred cancer care: a qualitative study. <i>BJGP Open</i> , 2020, 4, bjgpopen20X101023. | 1.8 | 7 |
| 58 | Gene expression of the mismatch repair gene MSH2 in primary colorectal cancer. <i>Tumor Biology</i> , 2011, 32, 977-983. | 1.8 | 6 |
| 59 | Gemcitabine, capecitabine and oxaliplatin in advanced biliary tract carcinoma. <i>Acta Oncologica</i> , 2014, 53, 1448-1450. | 1.8 | 6 |
| 60 | Study protocol: a randomized controlled trial comparing the efficacy of therapist guided internet-delivered cognitive therapy (TG-iConquerFear) with augmented treatment as usual in reducing fear of cancer recurrence in Danish colorectal cancer survivors. <i>BMC Cancer</i> , 2020, 20, 223. | 2.6 | 6 |
| 61 | Biliary-tract cancer: improving therapy by adding molecularly targeted agents. <i>Lancet Oncology</i> , The, 2012, 13, 118-119. | 10.7 | 5 |
| 62 | The Soluble Urokinase-Type Plasminogen Activator Receptor as a Biomarker for Survival and Early Treatment Effect in Metastatic Colorectal Cancer. <i>Cancers</i> , 2021, 13, 5100. | 3.7 | 5 |
| 63 | Clinical aspects and perspectives of erlotinib in the treatment of patients with biliary tract cancer. <i>Expert Opinion on Investigational Drugs</i> , 2016, 25, 359-365. | 4.1 | 4 |
| 64 | A single-center randomized clinical trial of palliative rehabilitation versus standard care alone in patients with newly diagnosed non-resectable cancer.. <i>Journal of Clinical Oncology</i> , 2018, 36, 75-75. | 1.6 | 4 |
| 65 | Impact of Open Dialogue about Complementary Alternative Medicine”A Phase II Randomized Controlled Trial. <i>Cancers</i> , 2022, 14, 952. | 3.7 | 4 |
| 66 | Pseudoprogression during treatment with pembrolizumab followed by rechallenge with chemotherapy in metastatic colorectal cancer: A case report. <i>Clinical Case Reports (discontinued)</i> , 2019, 7, 1445-1449. | 0.5 | 3 |
| 67 | Phase II study of gemcitabine, oxaliplatin and capecitabine in patients with KRAS exon 2 mutated biliary tract cancers. <i>Acta Oncologica</i> , 2020, 59, 298-301. | 1.8 | 3 |
| 68 | The impact of mismatch repair status to the preoperative staging of colon cancer: implications for clinical management. <i>Colorectal Cancer</i> , 2020, 9, CRC20. | 0.8 | 3 |
| 69 | The “œImmunescore” in rectal cancer: could we search quality beyond quantity of life?. <i>Oncotarget</i> , 2022, 13, 18-31. | 1.8 | 3 |
| 70 | <i>BRAF</i> refines clinical interpretation of mismatch repair deficiency in colorectal cancer. <i>Colorectal Cancer</i> , 2014, 3, 1-4. | 0.8 | 2 |
| 71 | Dynamic contrast-enhanced computed tomography as a potential biomarker in patients with metastatic colorectal cancer treated with regorafenib. <i>Acta Radiologica</i> , 2019, 60, 836-845. | 1.1 | 2 |
| 72 | Cross-sectoral communication by bringing together patient with cancer, general practitioner and oncologist in a video-based consultation: a qualitative study of oncologists” and nurse specialists” perspectives. <i>BMJ Open</i> , 2021, 11, e043038. | 1.9 | 2 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Re-exposure to immunotherapy in metastatic colon cancer: A case report. <i>Clinical Case Reports</i> (discontinued), 2021, 9, e04349. | 0.5 | 2 |
| 74 | Cross-sectoral video consultation in cancer care: GPS™ evaluation of a randomised controlled trial. <i>BJGP Open</i> , 2021, 5, BJGPO.2020.0114. | 1.8 | 2 |
| 75 | Treosulfan in platinum-resistant ovarian cancer. <i>International Journal of Gynecological Cancer</i> , 2021, 31, ijgc-2021-002395. | 2.5 | 1 |
| 76 | Functional precision medicine in colorectal cancer based on patient-derived tumoroids and in-vitro sensitivity drug testing. <i>Journal of Clinical Oncology</i> , 2021, 39, e15567-e15567. | 1.6 | 1 |
| 77 | Adjuvant chemotherapy with gemcitabine and cisplatin compared to observation after curative intent resection of cholangiocarcinoma and muscle invasive gallbladder carcinoma (ACTICCA-1): A randomized, multidisciplinary, multinational phase III trial. <i>Journal of Clinical Oncology</i> , 2015, 33, TPS4140-TPS4140. | 1.6 | 1 |
| 78 | A marker-driven phase II trial of neoadjuvant chemotherapy in locally advanced colon cancer. <i>Journal of Clinical Oncology</i> , 2014, 32, 3621-3621. | 1.6 | 1 |
| 79 | Natural killer cell activity: A test for immune reactivity with clinical perspectives. <i>Journal of Clinical Oncology</i> , 2018, 36, 87-87. | 1.6 | 1 |
| 80 | The prognostic importance of thymidylate gene polymorphism in colon cancer stage II. <i>International Journal of Colorectal Disease</i> , 2008, 23, 1267-1267. | 2.2 | 0 |
| 81 | The prognostic impact of RAS and RAF serum mutations in localized colon cancer. <i>Annals of Oncology</i> , 2016, 27, vi27. | 1.2 | 0 |
| 82 | Extended RAS and BRAF mutation analysis of circulating tumor DNA in patients with biliary tract cancer. <i>Annals of Oncology</i> , 2016, 27, vi235. | 1.2 | 0 |
| 83 | The Clinical Impact of MicroRNA-21 in Low Rectal Cancer Treated with High-Dose Chemoradiotherapy in the Organ Preserving Setting. <i>Gastrointestinal Disorders</i> , 2020, 2, 378-384. | 0.8 | 0 |
| 84 | PS1-2 Pembrolizumab vs chemotherapy for MSI-high/dMMR metastatic colorectal cancer: Asia subgroup of phase 3 KEYNOTE-177. <i>Annals of Oncology</i> , 2021, 32, S284. | 1.2 | 0 |
| 85 | Effectiveness study of gemcitabine, oxaliplatin, and capecitabine as first-line treatment for nonresectable biliary tract cancer. <i>Journal of Clinical Oncology</i> , 2014, 32, 334-334. | 1.6 | 0 |
| 86 | Feasibility of molecular patient selection in rare cancers: Phase II study of gemcitabine, oxaliplatin and capecitabine in KRAS mutated biliary tract cancer. <i>Journal of Clinical Oncology</i> , 2016, 34, e15620-e15620. | 1.6 | 0 |
| 87 | Monitoring the effect of first-line treatment in RAS/RAF mutated metastatic colorectal cancer by serial analysis of tumor specific DNA in plasma. <i>Journal of Clinical Oncology</i> , 2017, 35, 3593-3593. | 1.6 | 0 |
| 88 | Postponement of death weighed against duration of treatment and toxicity as key components in shared decision making about last line oncologic treatment. <i>Journal of Clinical Oncology</i> , 2017, 35, e21555-e21555. | 1.6 | 0 |
| 89 | Tumor specific methylation of NPY compared to RAS mutation in plasma DNA in the monitoring of colorectal cancer patients treated with last-line regorafenib. <i>Journal of Clinical Oncology</i> , 2018, 36, e15541-e15541. | 1.6 | 0 |
| 90 | Correlation between natural killer cell activity and treatment effect in patients with disseminated cancer. <i>Journal of Clinical Oncology</i> , 2018, 36, 12029-12029. | 1.6 | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 91 | Prognostic impact of CDX2 in stage II colon cancer: Results from two nationwide cohorts.. Journal of Clinical Oncology, 2018, 36, 3610-3610. | 1.6 | 0 |
| 92 | MicroRNA-126 and epidermal growth factor-like domain 7 predict recurrence in patients with colon cancer treated with neoadjuvant chemotherapy. , 2019, 2, 885-896. | | 0 |
| 93 | Prognostic impact of SOX9 in stage II colon cancer: Results from a large nationwide cohort.. Journal of Clinical Oncology, 2019, 37, e15165-e15165. | 1.6 | 0 |
| 94 | Efficacy of open dialogue about complementary and alternative medicine compared with standard care in improving quality of life in patients undergoing conventional oncology treatment (CAMONCO 2): protocol for a randomised controlled trial. BMJ Open, 2022, 12, e059960. | 1.9 | 0 |