
Rut Fernandez-Torres

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2425696/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Bioaccumulation and biochemical responses in the peppery furrow shell Scrobicularia plana exposed to a pharmaceutical cocktail at sub-lethal concentrations. Ecotoxicology and Environmental Safety, 2022, 242, 113845.	6.0	4
2	Uptake study in Juncus sp. and Salicornia europaea of six pharmaceuticals by liquid chromatography quadrupole time-of-flight mass spectrometry. Chemosphere, 2021, 266, 128995.	8.2	5
3	Urine and saliva biomonitoring by HF-LPME-LC/MS to assess dinitrophenols exposure. Microchemical Journal, 2021, 166, 106193.	4.5	8
4	Assessment of pharmaceutical mixture (ibuprofen, ciprofloxacin and flumequine) effects to the crayfish Procambarus clarkii: A multilevel analysis (biochemical, transcriptional and proteomic) Tj ETQq0 0 0 rgB	T/Otværloc	k 1 2 5Tf 50 611
5	A microfluidic liquid phase microextraction method for drugs and parabens monitoring in human urine. Microchemical Journal, 2021, 169, 106577.	4.5	8
6	Monitoring of pharmaceuticals in aquatic biota (Procambarus clarkii) of the Doñana National Park (Spain). Journal of Environmental Management, 2021, 297, 113314.	7.8	7
7	Comparison of three electromembrane-based extraction systems for NSAIDs analysis in human urine samples. Analytical and Bioanalytical Chemistry, 2020, 412, 6811-6822.	3.7	11
8	A Method for the Determination of Veterinary Drugs from Different Therapeutic Classes in Animal Urine. Journal of Chromatographic Science, 2020, 58, 127-135.	1.4	2
9	Simultaneous determination of metformin and glimepiride in human serum by ultra high performance liquid chromatography quadrupole time of flight mass spectrometry detection. Journal of Pharmaceutical and Biomedical Analysis, 2019, 165, 276-283.	2.8	12
10	Liquid chromatography quadrupole time-of-flight mass spectrometry determination of six pharmaceuticals in vegetal biota. Uptake study in Lavandula dentata. Science of the Total Environment, 2018, 622-623, 655-663.	8.0	7
11	Multiresidue determination of 21 pharmaceuticals in crayfish (Procambarus clarkii) using enzymatic microwave-assisted liquid extraction and ultrahigh-performance liquid chromatography-triple quadrupole mass spectrometry analysis. Journal of Pharmaceutical and Biomedical Analysis, 2018, 160, 144-151.	2.8	20
12	New nanostructured support for carrier-mediated electromembrane extraction of high polar compounds. Talanta, 2017, 162, 32-37.	5.5	40
13	New developments in microextraction techniques in bioanalysis. AÂreview. Analytica Chimica Acta, 2016, 905, 8-23.	5.4	169
14	Electromembrane extraction for the determination of parabens in water samples. Analytical and Bioanalytical Chemistry, 2016, 408, 1615-1621.	3.7	28
15	New developments in the extraction and determination of parabens in cosmetics and environmental samples. A review. Analytica Chimica Acta, 2015, 858, 1-15.	5.4	102
16	Hollow-fiber liquid-phase microextraction for the direct determination of flumequine in urban wastewaters by flow-injection analysis with terbium-sensitized chemiluminescence. Journal of Separation Science, 2014, 37, 2738-2744.	2.5	8
17	Application of chemiluminescence in the analysis of wastewaters – A review. Talanta, 2014, 122, 214-222.	5.5	54
18	A novel approach for electromembrane extraction based on the use of silver nanometallic-decorated hollow fibers. Analytica Chimica Acta, 2014, 849, 7-11.	5.4	47

Rut Fernandez-Torres

#	Article	IF	CITATIONS
19	Hollow fiber liquidâ€phase microextraction and determination of nonsteroidal antiâ€inflammatories by capillary electrophoresis and sulfonamides by HPLC in human urine. Biomedical Chromatography, 2013, 27, 246-253.	1.7	9
20	A novel application of three phase hollow fiber based liquid phase microextraction (HF-LPME) for the HPLC determination of two endocrine disrupting compounds (EDCs), n-octylphenol and n-nonylphenol, in environmental waters. Science of the Total Environment, 2013, 443, 1-6.	8.0	52
21	Electromembrane extraction (EME)—an easy, novel and rapid extraction procedure for the HPLC determination of fluoroquinolones in wastewater samples. Analytical and Bioanalytical Chemistry, 2013, 405, 2575-2584.	3.7	47
22	Identification of the specified impurities of silver sulfadiazine using a screening of degradation products in different stress physico-chemical media. Talanta, 2013, 116, 653-662.	5.5	9
23	Application of three phase hollow fiber based liquid phase microextraction (HF-LPME) for the simultaneous HPLC determination of phenol substituting compounds (alkyl-, chloro- and) Tj ETQq1 1 0.78431	4 rg &ī. \$Ovei	ിodങ10 Tf 50
24	Analytical Applications of Hollow Fiber Liquid Phase Microextraction (HF-LPME): A Review. Analytical Letters, 2012, 45, 804-830.	1.8	115
25	Simultaneous Determination of Selected Veterinary Antibiotics and their Main Metabolites in Fish and Mussel Samples by High-Performance Liquid Chromatography with Diode Array-Fluorescence (HPLC-DAD-FLD) Detection. Analytical Letters, 2011, 44, 2357-2372.	1.8	28
26	Electromembrane extraction (EME) and HPLC determination of non-steroidal anti-inflammatory drugs (NSAIDs) in wastewater samples. Talanta, 2011, 85, 394-399.	5.5	119
27	Hollow fiber-based liquid phase microextraction (HF-LPME) for a highly sensitive HPLC determination of sulfonamides and their main metabolites. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2011, 879, 197-204.	2.3	81
28	Capillary electrophoresis determination of nonsteroidal antiâ€inflammatory drugs in wastewater using hollow fiber liquidâ€phase microextraction. Electrophoresis, 2011, 32, 2107-2113.	2.4	31
29	Trace-metal distribution of cigarette ashes as marker of tobacco brands. Forensic Science International, 2011, 204, 119-125.	2.2	25
30	Enzymatic-microwave assisted extraction and high-performance liquid chromatography–mass spectrometry for the determination of selected veterinary antibiotics in fish and mussel samples. Journal of Pharmaceutical and Biomedical Analysis, 2011, 54, 1146-1156.	2.8	58
31	Hollow fiber-based liquid phase microextraction (HF-LPME) as a new approach for the HPLC determination of fluoroquinolones in biological and environmental matrices. Journal of Pharmaceutical and Biomedical Analysis, 2011, 55, 332-341.	2.8	76
32	Temporal Evolution of Linear Alkylbenzene Sulfonates and Heavy Metals in Sludge from Wastewater Treatment Plant. Water Environment Research, 2011, 83, 411-417.	2.7	1
33	Application of enzymatic probe sonication extraction for the determination of selected veterinary antibiotics and their main metabolites in fish and mussel samples. Analytica Chimica Acta, 2010, 675, 156-164.	5.4	26
34	Simultaneous determination of 11 antibiotics and their main metabolites from four different groups by reversed-phase high-performance liquid chromatography–diode array–fluorescence (HPLC–DAD–FLD) in human urine samples. Talanta, 2010, 81, 871-880.	5.5	67
35	Application of hollow fiber-based liquid-phase microextraction (HF-LPME) for the determination of acidic pharmaceuticals in wastewaters. Talanta, 2010, 82, 854-858.	5.5	110
36	Rapid flow injection method for the determination of sulfite in wine using the permanganate–luminol luminescence system. Talanta, 2010, 82, 2003-2006.	5.5	31

#	Article	IF	CITATIONS
37	Rapid Flow-Injection Method for the Determination of Colistin by Sensitized Chemiluminescence Using the Acidic Permanganate–Sulfite System. Analytical Letters, 2009, 42, 1471-1478.	1.8	4
38	HPLC determination of ibuprofen, diclofenac and salicylic acid using hollow fiber-based liquid phase microextraction (HF-LPME). Analytica Chimica Acta, 2009, 653, 184-190.	5.4	129
39	Hollow fiber-based liquid-phase microextraction (HF-LPME) of ibuprofen followed by FIA-chemiluminescence determination using the acidic permanganate–sulfite system. Talanta, 2009, 79, 911-915.	5.5	32
40	Determination of imipenem and rifampicin in mouse plasma by high performance liquid chromatography–diode array detection. Analytica Chimica Acta, 2008, 608, 204-210.	5.4	13
41	Application of probe sonication extraction for the determination of linear alkylbenzene sulfonates from sewage sludge. Comparison with other extraction methods. Microchemical Journal, 2008, 90, 164-170.	4.5	11
42	Urea as new stabilizing agent for imipenem determination. Talanta, 2008, 77, 241-248.	5.5	12
43	Mineral content and botanical origin of Spanish honeys. Talanta, 2005, 65, 686-691.	5.5	198
44	Spectrofluorimetric determination of acrivastine in spiked human urine and pharmaceuticals. Talanta, 2002, 56, 571-576.	5.5	4
45	Electrochemical behaviour and determination of acrivastine in pharmaceuticals and human urine. Journal of Pharmaceutical and Biomedical Analysis, 2002, 30, 1215-1222.	2.8	16
46	Electrochemical oxidation of cisatracurium on carbon paste electrode and its analytical applications. Talanta, 2001, 53, 1179-1185.	5.5	15
47	Spectrofluorimetric determination of cisatracurium and mivacurium in spiked human serum and pharmaceuticals. Talanta, 1999, 49, 881-887.	5.5	7
48	New Methodologies for Assessing the Presence and Ecological Effects of Pesticides in DonÌfana National Park (SW Spain). , 0, , .		2