

Xiang-Shan Wang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2431599/publications.pdf>

Version: 2024-02-01

197
papers

2,846
citations

201674

27
h-index

254184

43
g-index

231
all docs

231
docs citations

231
times ranked

2053
citing authors

#	ARTICLE	IF	CITATIONS
1	An efficient synthesis of 6-arylpyrazolo[4,5-pyrimido[2,1-a]isoquinolin-8(9H)-one derivatives catalyzed by AgOTf. <i>Journal of Heterocyclic Chemistry</i> , 2022, 59, 890-898.	2.6	1
2	Copper-assisted Wittig-type olefination of aldehydes with <i>p</i> -toluenesulfonylmethyl isocyanide. <i>Organic Chemistry Frontiers</i> , 2022, 9, 4158-4163.	4.5	4
3	Silver-Catalyzed Controlled Intermolecular Cross-Coupling of Silyl Enol Ethers: Scalable Access to 1,4-Diketones. <i>Organic Letters</i> , 2022, 24, 4513-4518.	4.6	18
4	CuI-catalyzed synthesis of Benzoimidazo[1,4]diazepinoindoles/indazoles via double Ullmann cross-coupling reaction. <i>Tetrahedron</i> , 2022, 121, 132835.	1.9	3
5	Synthesis of Sulfonylated Heterocycles via Copper-Catalyzed Heteroaromatization/Sulfonyl Transfer of Propargylic Alcohols. <i>Chemistry - an Asian Journal</i> , 2021, 16, 30-33.	3.3	9
6	Silver-Catalyzed [3+1+1] Annulation of Nitrones with Isocynoacetates as an Approach to 1,4,5-Trisubstituted Imidazoles. <i>European Journal of Organic Chemistry</i> , 2021, 2021, 964-968.	2.4	7
7	An efficient synthesis of diimidazo[1,2-a:1,2-c]quinazolines via a copper-catalyzed double Ullmann cross-coupling reaction. <i>Tetrahedron</i> , 2021, 81, 131918.	1.9	7
8	Silver-Promoted (4 + 1) Annulation of Isocynoacetates with Alkylpyridinium Salts: Divergent Regioselective Synthesis of 1,2-Disubstituted Indolizines. <i>Organic Letters</i> , 2021, 23, 7555-7560.	4.6	14
9	Pd(II)-Catalyzed Arylation/Oxidation of Benzylic C-H of 8-Methylquinolines: Access to 8-Benzoylquinolines. <i>Journal of Organic Chemistry</i> , 2021, 86, 15423-15432.	3.2	3
10	Synthesis of 15-Arylisoquinolino[2,1-a:1,2]imidazo[4,5-f][1,10]phenanthrolines catalyzed by Copper(I)/ <i>o</i> -Phen. <i>Research on Chemical Intermediates</i> , 2021, 47, 2063-2074.	2.7	1
11	Modular synthesis of 3-substituted isocoumarins via silver-catalyzed aerobic oxidation/6-endo heterocyclization of ortho-alkynylbenzaldehydes. <i>Organic and Biomolecular Chemistry</i> , 2021, 19, 6657-6664.	2.8	8
12	A rearrangement of saccharin-derived cyclic ketimines with 3-chlorooxindoles leading to spiro-1,3-benzothiazine oxindoles. <i>Chemical Communications</i> , 2021, 57, 11322-11325.	4.1	5
13	Synthesis of Benzo[4,5]imidazo[1,2-a]naphthyridine and Benzo[4,5]imidazo[2,1-a]isoquinoline Derivatives Catalyzed by CuI/L-Proline. <i>Polycyclic Aromatic Compounds</i> , 2020, 40, 465-474.	2.6	1
14	Cascade C-N and C-O bond constructions for the synthesis of dibenzoimidazo[1,4]oxazepines catalyzed by CuI/ <i>o</i> -phen. <i>Journal of Heterocyclic Chemistry</i> , 2020, 57, 851-858.	2.6	5
15	CuI catalyzed synthesis of Dibenzo[b,f]imidazo[1,2-d][1,4]thiazepines via C-N and C-S bond Ullmann cross-coupling reaction. <i>Tetrahedron</i> , 2020, 76, 130915.	1.9	7
16	Copper(I)-catalyzed synthesis of isoindolo[1,2-b]quinazoline derivatives via an $\hat{\pi}$ -arylation under Pd and ligand free conditions. <i>Tetrahedron Letters</i> , 2020, 61, 152508.	1.4	1
17	An efficient synthesis of 6-benzyl-2-arylthieno[2,3-d]pyrimidin-4(3H)-ones catalyzed by HCl involving a Friedel-Crafts alkylation reaction. <i>Journal of Heterocyclic Chemistry</i> , 2020, 57, 3970-3979.	2.6	0
18	CuBr-Catalyzed $\hat{\pi}$ -Arylation and Aerobic Oxidative Dehydrogenative C-N Coupling for the Synthesis of Spiro[cyclohexane-1,12-isoindolo[1,2-b]quinazolin]-10-one Derivatives. <i>Organic Letters</i> , 2020, 22, 2887-2891.	4.6	9

#	ARTICLE	IF	CITATIONS
19	A Cascade synthesis of 11 <i>bH</i> -imidazo[1,2- <i>c</i>]isoquinolino[2,1- <i>a</i>]quinazoline derivatives catalyzed by AgOTf. <i>Journal of Heterocyclic Chemistry</i> , 2020, 57, 2203-2212.	2.6	4
20	Silver-Assisted [3 + 2] Annulation of Nitrones with Isocyanides: Synthesis of 2,3,4-Trisubstituted 1,2,4-Oxadiazolidin-5-ones. <i>Journal of Organic Chemistry</i> , 2020, 85, 3560-3567.	3.2	15
21	CuI-catalyzed synthesis of (benzo)imidazo[2,1- <i>a</i>]isoquinolinone derivatives via successive $\hat{\pm}$ -arylation, deacylation and benzyl automatic oxidation. <i>Tetrahedron</i> , 2020, 76, 131200.	1.9	3
22	Cooperative Silver- and Base-Catalyzed Diastereoselective Cycloaddition of Nitrones with Methylene Isocyanides: Access to 2-Imidazolinones. <i>European Journal of Organic Chemistry</i> , 2020, 2020, 3475-3479.	2.4	10
23	Switchable Copper-Catalyzed Approach to Benzodithiole, Benzothiaselenole, and Dibenzodithiocine Skeletons. <i>Organic Letters</i> , 2020, 22, 3454-3459.	4.6	20
24	Synthesis of Structurally Diversified Benzo[<i>c</i>]chromene Derivatives under (An)aerobic Conditions Catalyzed by CuI. <i>Journal of Heterocyclic Chemistry</i> , 2019, 56, 2822-2830.	2.6	7
25	Copper-Catalyzed Synthesis of Dibenzo[<i>b,f</i>]imidazo[1,2- <i>d</i>][1,4]oxazepine Derivatives via a Double Ullmann Coupling Reaction. <i>Synthesis</i> , 2019, 51, 1662-1668.	2.3	9
26	An efficient synthesis of 6-hydroxy-6-methyl-5,6-dihydro-8 <i>H</i> -isoquinolino[1,2- <i>b</i>]quinazolin-8-ones via a CuI-catalyzed deacylation and no dehydration reaction. <i>Monatshefte für Chemie</i> , 2019, 150, 1305-1315.	1.8	2
27	Silver-Induced [3+2] Cycloaddition of Isocyanides with Acyl Chlorides: Regioselective Synthesis of 2,5-Disubstituted Oxazoles. <i>ChemCatChem</i> , 2019, 11, 4272-4275.	3.7	16
28	Silver-Mediated Synthesis of Substituted Benzofuran- and Indole-Pyrroles via Sequential Reaction of <i>ortho</i> -Alkynylaromatics with Methylene Isocyanides. <i>Journal of Organic Chemistry</i> , 2019, 84, 8998-9006.	3.2	17
29	Silver Triflate Catalyzed Synthesis of Isoquinolino[2,1- <i>a</i>]quinazolin-3(2- <i>c</i>)quinazoline Derivatives via Alkyne Hydroamination. <i>Synthesis</i> , 2019, 51, 3101-3108.	2.3	7
30	Copper/I-proline-catalyzed synthesis of 5-amino-2,3-diphenylimidazo[2,1- <i>a</i>]isoquinolines in the presence of Cs ₂ CO ₃ . <i>Monatshefte für Chemie</i> , 2019, 150, 681-689.	1.8	3
31	Copper-Catalyzed Synthesis of 13-Aminoisoquinolino[2,1- <i>a</i>]perimidine-12-carboxylates <i>via</i> $\hat{\pm}$ -Arylation with a High Chemoselectivity. <i>Journal of Heterocyclic Chemistry</i> , 2019, 56, 663-669.	2.6	2
32	Silver-Catalyzed Sequential Cascade Reaction of Isocyanides with 1-(2-Ethynylphenyl)prop-2-yn-1-ol: Access to Benzo[<i>b</i>]fluorenes and Benzofuran-Pyrroles. <i>Advanced Synthesis and Catalysis</i> , 2019, 361, 1543-1548.	4.3	20
33	An efficient synthesis of biaryl diamides <i>via</i> Ullmann coupling reaction catalyzed by CuI in the presence of Cs ₂ CO ₃ and TBAB. <i>Research on Chemical Intermediates</i> , 2018, 44, 5271-5283.	2.7	4
34	Study on the iodine-catalyzed reaction of 3-aminopyrazine-2-carbohydrazide and 2-(arylethynyl)benzaldehydes. <i>Tetrahedron</i> , 2018, 74, 1468-1475.	1.9	7
35	One-pot synthesis of 2,3-diphenyl-6,7-dihydroimidazo[1,2- <i>f</i>]phenanthridin-8(5 <i>H</i>)-ones catalyzed by CuI/I-proline. <i>Monatshefte für Chemie</i> , 2018, 149, 569-576.	1.8	7
36	An efficient synthesis of 16 <i>H</i> -dibenzo[2,3:6,7][1,4]oxazepino[5,4- <i>b</i>]quinazolin-16-ones <i>via</i> an Ullmann reaction catalyzed by CuI. <i>Organic and Biomolecular Chemistry</i> , 2018, 16, 1679-1685.	2.8	11

#	ARTICLE	IF	CITATIONS
37	One-Pot Four-Component Synthesis of 5,10-Diarylpyrido[4,3-b][1,6] Naphthyridine Derivatives in Ionic Liquids Catalyzed by TsOH. <i>Polycyclic Aromatic Compounds</i> , 2018, 38, 236-243.	2.6	2
38	Green Synthesis of 6-Aryl-5,6-dihydrobenzo[4,5]imidazo[1,2-c]quinazoline Derivatives in Ionic Liquid under Catalyst-free Conditions. <i>Journal of Heterocyclic Chemistry</i> , 2018, 55, 166-172.	2.6	11
39	[3 + 2] Cycloaddition of Isocyanides with Aryl Diazonium Salts: Catalyst-Dependent Regioselective Synthesis of 1,3- and 1,5-Disubstituted 1,2,4-Triazoles. <i>Organic Letters</i> , 2018, 20, 6930-6933.	4.6	58
40	Synthesis of Substituted 4-H-Thiochromen-4-imines via Copper-Catalyzed Cyclization Cascades of α -Bromobenzothioamides with Terminal Alkynes. <i>Journal of Organic Chemistry</i> , 2018, 83, 9504-9509.	3.2	6
41	Dioxane-involving reaction for the synthesis of 3-aryl-1-(2-(vinylloxy)ethoxy)isoquinolines catalyzed by AgOTf. <i>Organic and Biomolecular Chemistry</i> , 2018, 16, 6070-6076.	2.8	8
42	Iodine-catalyzed synthesis of 5-benzoyl-8H-phthalazino[1,2-b]quinazolin-8-one derivatives via a domino reaction involving a benzyl automatic oxidation by oxygen. <i>Tetrahedron</i> , 2018, 74, 4746-4753.	1.9	4
43	A Consecutive Condensation, Cyclization, and Dehydration for the Synthesis of Benzimidazopyrroloquinazolines Catalyzed by TsOH. <i>Journal of Heterocyclic Chemistry</i> , 2018, 55, 2325-2333.	2.6	2
44	The Chemoselective Reaction of 2-Amino-N-arylbzohydrazide and Ketonic Acid Catalyzed by Iodine for the Synthesis of Quinazoline Derivatives. <i>Journal of Heterocyclic Chemistry</i> , 2018, 55, 1906-1916.	2.6	2
45	Green Synthesis of Benzo or Cyclopenta[1,7]phenanthroline Derivatives in EtOH under Catalyst-free Conditions. <i>Journal of Heterocyclic Chemistry</i> , 2017, 54, 248-254.	2.6	1
46	One-Pot Ullmann C-N Coupling Cyclization Toward Domino Synthesis of Fused Hexacyclic Quinolinotriazoloacridinones Catalyzed by CuI/Proline. <i>Journal of Heterocyclic Chemistry</i> , 2017, 54, 986-992.	2.6	4
47	One-Pot Three-Component Synthesis of Pyrido[2,3-c]carbazole Derivatives in EtOH under Catalyst-free Conditions. <i>Journal of Heterocyclic Chemistry</i> , 2017, 54, 1378-1383.	2.6	3
48	Catalyst-free Synthesis of 5-Arylimidazo[1,2-c]quinazoline Derivatives in Ionic Liquids. <i>Journal of Heterocyclic Chemistry</i> , 2017, 54, 509-516.	2.6	6
49	N-Heterocyclic Carbene-Catalyzed [4 + 2] Cyclization of Saturated Carboxylic Acid with α -Quinone Methides through in Situ Activation: Enantioselective Synthesis of Dihydrocoumarins. <i>Journal of Organic Chemistry</i> , 2017, 82, 1790-1795.	3.2	58
50	Synthesis of Pyridophenanthrolines via a Three-Component Reaction Involving 1,10-Phenanthroline-5-Amine. <i>Journal of Heterocyclic Chemistry</i> , 2017, 54, 2266-2271.	2.6	1
51	Consecutive Sonogashira Coupling and Hydroamination Cyclization for the Synthesis of Isoindolo[1,2-b]quinazolin-10(12H)-ones Catalyzed by CuI/Proline. <i>Journal of Organic Chemistry</i> , 2017, 82, 4918-4923.	3.2	41
52	One-Pot Three-Component Synthesis of 6-H-chromeno[4,3-b] or Cyclopenta[b]furo[3,2-f]quinoline Derivatives. <i>Journal of Heterocyclic Chemistry</i> , 2017, 54, 2929-2934.	2.6	8
53	Structurally diversified synthesis of 2,3-dihydroquinazolin-4-(1H)-ones from 2-aminobenzamides and 1,2-dicarbonyl compounds in ionic liquids catalyzed by iodine. <i>Research on Chemical Intermediates</i> , 2017, 43, 2985-3005.	2.7	6
54	CuI-catalyzed Sonogashira reaction for the efficient synthesis of 1 H-imidazo[2,1-a]isoquinoline derivatives. <i>Tetrahedron</i> , 2017, 73, 4698-4705.	1.9	29

#	ARTICLE	IF	CITATIONS
55	An efficient synthesis of 6-arylbenzo[4,5]imidazo[2,1-a]isoquinolines via sequential $\hat{\pm}$ -arylation of carbonyl and deacylation catalyzed by CuI. <i>Organic and Biomolecular Chemistry</i> , 2017, 15, 5325-5331.	2.8	22
56	Copper(I)-catalyzed $\hat{\pm}$ -arylation of carbonyl cascade reaction leading to benzo[4, 5]imidazo[1,2-f]phenanthridin-4(1H)-one derivatives. <i>Research on Chemical Intermediates</i> , 2017, 43, 5995-6006.	2.7	4
57	Cu(OAc) ₂ -Catalyzed Aerobic Oxidative Dehydrogenation Coupling: Synthesis of Heptacyclic Quinolizino[3,4,5,6- <i>kl</i>]perimidines. <i>Journal of Organic Chemistry</i> , 2017, 82, 1817-1822.	3.2	40
58	An efficient green synthesis of 5 <i>H</i> -spiro[benzo[4,5]imidazo[1,2- <i>cd</i>]quinazoline-6,3 <i>â</i> -indolin]-2 <i>â</i> -ones catalyzed by iodine in ionic liquids. <i>Heterocyclic Communications</i> , 2017, 23, 385-388.	1.2	1
59	An Efficient Synthesis of Pyrrolo[1,2- <i>ac</i>] or Pyrido[1,2- <i>ac</i>]benzo[4,5]imidazo[1,2- <i>cd</i>]quinazoline Derivatives in Ionic Liquids Catalyzed by Iodine. <i>Journal of Heterocyclic Chemistry</i> , 2017, 54, 3440-3446.	2.6	7
60	An efficient synthesis of quinazoline or pyrrolo[1,2- <i>a</i>]quinazolin-5(1H)-one derivatives in ionic liquids catalyzed by iodine. <i>Research on Chemical Intermediates</i> , 2017, 43, 6787-6801.	2.7	4
61	Copper-catalyzed synthesis of arylcarboxamides from aldehydes and isocyanides: the isocyano group as an N1 synthon. <i>Organic and Biomolecular Chemistry</i> , 2017, 15, 6314-6317.	2.8	14
62	Green Synthesis of Spiro[indoline-3,4- <i>â</i> pyrazolo[3,4- <i>bc</i>][1,6]naphthyridine]-2,5- <i>â</i> (1 <i>â</i> - <i>H</i>)- <i>â</i> -diones Catalyzed by TsOH in Ionic Liquids. <i>Journal of Heterocyclic Chemistry</i> , 2016, 53, 1578-1583.	2.6	5
63	Copper-catalyzed synthesis of 1-amino-5-arylindazolo[3,2- <i>b</i>]quinazolin-7(5 <i>H</i>)-ones via a ring-opening reaction of 4-halogenated isatin. <i>Tetrahedron</i> , 2016, 72, 3844-3850.	1.9	10
64	Iodine-catalyzed synthesis of 5 <i>H</i> -phthalazino[1,2- <i>b</i>]quinazoline and isoindolo[2,1- <i>a</i>]quinazoline derivatives via a chemoselective reaction of 2-aminobenzohydrazide and 2-formylbenzoic acid in ionic liquids. <i>Tetrahedron Letters</i> , 2016, 57, 2515-2519.	1.4	18
65	Synthesis of spiro[pyrazole-4,8- <i>â</i> pyrazolo[3,4- <i>f</i>]quinolin]-5(1H)-ones by the reaction of aldehydes with 1 <i>H</i> -indazol-6-amine and 1 <i>H</i> -pyrazol-5(4H)-one. <i>Heterocyclic Communications</i> , 2016, 22, .	1.2	3
66	Parallel Synthesis of Pyrrolo[3,2- <i>f</i>]quinolines (PQQ Skeleton) Library via a One-Pot Three-Component Reaction under Catalyst-Free Conditions. <i>Polycyclic Aromatic Compounds</i> , 2016, 36, 683-696.	2.6	2
67	A Green Synthesis of Fused Polycyclic 5 <i>H</i> -Chromeno[3,2- <i>c</i>]quinoline-6,8(7 <i>H</i> ,9 <i>H</i>)-dione Derivatives Catalyzed by TsOH in Ionic Liquids. <i>Polycyclic Aromatic Compounds</i> , 2016, 36, 758-772.	2.6	8
68	An Efficient Synthesis of Fused Polycyclic Triazolo[4,5- <i>ac</i>]acridine Derivatives under Catalyst-Free Conditions with High Regioselectivity. <i>Polycyclic Aromatic Compounds</i> , 2016, 36, 671-682.	2.6	2
69	Copper-catalyzed Ullmann reaction for the synthesis of fused hexacyclic heterocycles containing naphthyridine, acridine, and pyrazole (imidazole) moieties. <i>Monatshefte für Chemie</i> , 2016, 147, 1233-1242.	1.8	4
70	An Enantioselective Assembly of Dihydropyranones through an NHC/LiCl ₄ -Mediated in situ Activation of $\hat{\pm}$ -Unsaturated Carboxylic Acids. <i>Chemistry - an Asian Journal</i> , 2016, 11, 678-681.	3.3	27
71	Synthesis of 6-aryl-5 <i>H</i> -quinazolino[4,3- <i>b</i>]quinazolin-8(6 <i>H</i>)-one derivatives in ionic liquids catalyzed by iodine. <i>Research on Chemical Intermediates</i> , 2016, 42, 1045-1055.	2.7	5
72	Iodine-catalyzed synthesis of dibenzo[<i>b,h</i>][1,6]naphthyridine-11-carboxamides via a domino reaction involving double elimination of hydrogen bromide. <i>Organic and Biomolecular Chemistry</i> , 2016, 14, 2774-2779.	2.8	14

#	ARTICLE	IF	CITATIONS
73	Copper(I)-catalyzed synthesis of thienopyrazoloquinazolinone derivatives under ligand-free conditions. <i>Research on Chemical Intermediates</i> , 2016, 42, 6769-6776.	2.7	4
74	Iodine-catalyzed synthesis of fused tetracyclic pyridazino[6,1-b]pyrrolo[1,2-a]quinazolin-9(1H)-one derivatives via a tandem reaction. <i>Tetrahedron</i> , 2016, 72, 2178-2185.	1.9	14
75	Iodine-Catalyzed Synthesis of Fused Polycyclic Heterocycles Containing Pyrazoloquinoline via Povarov Reaction. <i>Polycyclic Aromatic Compounds</i> , 2016, 36, 275-283.	2.6	1
76	Green Synthesis of Fused Polycyclic Pyrazolo[3,4- <i>b</i>][1,6]naphthyridine Derivatives in Ionic Liquids via Three-Component Reaction. <i>Polycyclic Aromatic Compounds</i> , 2016, 36, 478-489.	2.6	6
77	Formation of Csp ² -N bond under metal-catalyst-free conditions for the synthesis of pyridopyrazoloquinazoline derivatives. <i>Monatshefte für Chemie</i> , 2016, 147, 775-782.	1.8	4
78	A Convenient Synthesis of Spiro[isoxazole-pyrazoloquinoline] Derivatives under Catalyst-Free Conditions. <i>Synthesis</i> , 2015, 48, 65-72.	2.3	4
79	Enantioselective Assembly of Spirocyclic Oxindole-dihydropyranones through NHC-Catalyzed Cascade Reaction of Isatins with N-Hydroxybenzotriazole Esters of α,β -Unsaturated Carboxylic Acid. <i>Journal of Organic Chemistry</i> , 2015, 80, 3289-3294.	3.2	60
80	A highly regioselective synthesis of functionalized furo[3,2- <i>a</i>]acridine derivatives via a three-component reaction. <i>Research on Chemical Intermediates</i> , 2015, 41, 9917-9927.	2.7	2
81	Copper(I) Iodide Catalyzed Synthesis of Fused Hexacyclic Pyrazolo[4,5,1- <i>de</i>]quinolino[4,3,2- <i>mn</i>]acridin-14(11H)-ones under Ligand-Free Conditions. <i>Synthesis</i> , 2015, 47, 562-568.	2.3	10
82	Domino synthesis of fused pyrazolo[5,1- <i>b</i>]quinazolin-9(1 H)-ones catalyzed by CuI via subsequent Michael addition and elimination. <i>Tetrahedron</i> , 2015, 71, 8732-8737.	1.9	8
83	An efficient synthesis of 11-aryl-10-oxo-7,8,10,11-tetrahydro-1H-[1,2,3]triazolo[4- α^2 ,5- α^2 :3,4]benzo[1,2- <i>b</i>][1,6]naphthyridine derivatives under catalyst-free conditions. <i>Heterocyclic Communications</i> , 2015, 21, 377-380.	1.2	1
84	Green synthesis of polysubstituted quinoline and benzoquinoline derivatives in ionic liquid via a three-component reaction. <i>Research on Chemical Intermediates</i> , 2015, 41, 7393-7403.	2.7	5
85	Convenient synthesis of naphtho[1,6] naphthyridine derivatives under catalyst-free conditions. <i>Research on Chemical Intermediates</i> , 2015, 41, 1703-1714.	2.7	4
86	A Convenient Synthesis of Pyridophenanthroline Derivatives under Catalyst Free Conditions. <i>Journal of Heterocyclic Chemistry</i> , 2015, 52, 373-379.	2.6	2
87	Green synthesis of naphthyridine derivatives in ionic liquid via three-component reaction. <i>Research on Chemical Intermediates</i> , 2015, 41, 3873-3884.	2.7	3
88	A three-component domino reaction for efficient synthesis of functionalized pyrazolo[3,4- <i>f</i>]quinolines under catalyst-free conditions. <i>Research on Chemical Intermediates</i> , 2015, 41, 6339-6350.	2.7	4
89	An Efficient Synthesis of Polycyclic Heterocycles Containing Pyrazolo[3,4- <i>f</i>]quinoline or Benzo[<i>h</i>]indazolo[6,7- <i>b</i>][1,6]naphthyridine Under Catalyst-Free Conditions. <i>Polycyclic Aromatic Compounds</i> , 2014, 34, 606-619.	2.6	11
90	A Green Synthesis of Pyrido[1,2- <i>a</i>]quinazolin-1,6-dione Derivatives in Ionic Liquid Catalyzed by Iodine. <i>Journal of Heterocyclic Chemistry</i> , 2014, 51, E314.	2.6	4

#	ARTICLE	IF	CITATIONS
91	A Green Synthesis of Pyrrolo[1,2- <i>a</i>]quinazolin-5(1 <i>H</i>)-one Derivatives in Ionic Liquids Catalyzed by Iodine. <i>Journal of Heterocyclic Chemistry</i> , 2014, 51, 1472-1475.	2.6	7
92	Green synthesis of bis-quinazolinone derivatives catalyzed by iodine in ionic liquids. <i>Research on Chemical Intermediates</i> , 2014, 40, 2823-2835.	2.7	12
93	Iodine-catalyzed Povarov reaction for synthesis of cyclobuta[<i>c</i>]quinoline derivatives. <i>Research on Chemical Intermediates</i> , 2014, 40, 1103-1113.	2.7	3
94	Copper(I)-catalyzed synthesis of 1-arylpyrazolo[5,1- <i>b</i>]quinazolin-9(1 <i>H</i>)-one via intramolecular alkyne hydroamination. <i>Tetrahedron</i> , 2014, 70, 2889-2893.	1.9	25
95	Iodine-catalyzed synthesis of 2-arylpyrazolo[5,1- <i>b</i>]quinazolin-9(3 <i>H</i>)-one derivatives in ionic liquids via domino reaction. <i>Tetrahedron</i> , 2014, 70, 3440-3446.	1.9	19
96	Three-Component One-Pot Synthesis of Indolo[3,4- <i>a</i>]acridine Derivatives with High Regioselectivity under Catalyst-Free Conditions. <i>Journal of Heterocyclic Chemistry</i> , 2014, 51, E349.	2.6	7
97	Iodine-Catalyzed Synthesis of Cyclopenta[<i>c</i>]quinoline Derivatives via Imino Diels-Alder Reaction. <i>Journal of Heterocyclic Chemistry</i> , 2014, 51, 830-834.	2.6	10
98	An Efficient Method for the Synthesis of 3-Arylnaphtho[2,3- <i>a</i>]quinoline-1,2-dicarboxylate Derivatives Catalyzed by Yb(OTf) ₃ . <i>Journal of Heterocyclic Chemistry</i> , 2014, 51, 502-506.	2.6	6
99	Domino synthesis of fused hexacyclic imidazoquinolinoacridinones catalyzed by CuI/l-proline. <i>Tetrahedron</i> , 2014, 70, 8919-8924.	1.9	15
100	A Selective Method for the Synthesis of <i>N,N</i> -diarylbenzene-1,4-diamine and Dispirocyclic Quinazolinone Derivatives Catalyzed by Iodine. <i>Journal of Heterocyclic Chemistry</i> , 2014, 51, 1363-1368.	2.6	6
101	CuI-Catalyzed C-N Bond Formation and Cleavage for the Synthesis of Benzimidazo[1,2- <i>a</i>]quinazoline Derivatives. <i>Journal of Organic Chemistry</i> , 2014, 79, 5847-5851.	3.2	69
102	Synthesis of Isoindolo[2,1- <i>a</i>]quinazoline Derivatives in Ionic Liquid Catalyzed by Iodine. <i>Journal of Heterocyclic Chemistry</i> , 2014, 51, 630-634.	2.6	17
103	An Efficient Synthesis of Pyrrolo[1,2- <i>a</i>]quinazoline Derivatives in Ionic Liquid Catalyzed by Iodine. <i>Journal of Heterocyclic Chemistry</i> , 2014, 51, 841-845.	2.6	13
104	The N-H...X Hydrogen Bonds in the Crystal Structures of (Thio)isochromene Derivatives. <i>Journal of Chemical Crystallography</i> , 2013, 43, 26-30.	1.1	5
105	Combinatorial Synthesis of Pyrrolo[3,2- <i>f</i>]quinoline and Pyrrolo[3,2- <i>a</i>]acridine Derivatives via a Three-Component Reaction under Catalyst-Free Conditions. <i>ACS Combinatorial Science</i> , 2013, 15, 498-502.	3.8	30
106	Iodine-catalyzed synthesis of pyrrolo[1,2- <i>a</i>]quinazoline-3 <i>a</i> -carboxylic acid derivatives in ionic liquids. <i>Research on Chemical Intermediates</i> , 2013, 39, 3327-3335.	2.7	5
107	Iodine-catalyzed synthesis of pyrazolo[4,3- <i>f</i>]quinoline derivatives via a highly regio-selective Povarov reaction. <i>Research on Chemical Intermediates</i> , 2013, 39, 1781-1787.	2.7	7
108	Synthesis of bis-benzoquinoline derivatives catalyzed by iodine via ring-opening of furan. <i>Tetrahedron</i> , 2013, 69, 7045-7050.	1.9	17

#	ARTICLE	IF	CITATIONS
109	An Efficient Synthesis of Clopenta[b]pyrazolo[4,3-f]quinolin-9(3H)-one Derivatives by Three-component Reaction in Ionic Liquids. <i>Journal of Heterocyclic Chemistry</i> , 2013, 50, 937-940.	2.6	2
110	Structurally diversified products from the reactions of 2-aminobenzamides with 1,3-cyclohexanediones catalyzed by iodine. <i>Tetrahedron Letters</i> , 2013, 54, 757-760.	1.4	27
111	Combinatorial Synthesis of Fused Tetracyclic Heterocycles Containing [1,6]Naphthyridine Derivatives under Catalyst Free Conditions. <i>ACS Combinatorial Science</i> , 2013, 15, 267-272.	3.8	16
112	Copper(I)-Catalyzed Synthesis of 5-Arylindazolo[3,2-b]quinazolin-7(5H)-one via Ullmann-Type Reaction. <i>Journal of Organic Chemistry</i> , 2013, 78, 5700-5704.	3.2	49
113	Ionic Liquid as an Efficient and Recyclable Reaction Medium for the Synthesis of Pyrido[2,3-d]pyrimidines. <i>Journal of Heterocyclic Chemistry</i> , 2013, 50, 534-538.	2.6	12
114	Crystal structure of 1-methyl-5-(trifluoromethyl)-1H-benzo[d]imidazole-2(3H)-thione, C ₉ H ₇ F ₃ N ₂ S. <i>Zeitschrift Fur Kristallographie - New Crystal Structures</i> , 2013, 228, 187-188.	0.3	1
115	Combinatorial Synthesis of Pyrazoloquinoline and Pyrazoloacridine Derivatives with High Regioselectivity. <i>Combinatorial Chemistry and High Throughput Screening</i> , 2013, 16, 550-561.	1.1	5
116	Green Method for the Synthesis of Polysubstituted Chromene Derivatives in Ionic Liquids. <i>Synthetic Communications</i> , 2012, 42, 599-607.	2.1	8
117	Mild and Efficient One-Pot Three-Component Synthesis of Benzopyrimidoquinoline-Tetraone Derivatives in Ionic Liquids. <i>Journal of Chemical Research</i> , 2012, 36, 453-456.	1.3	11
118	Reaction of 2-Aminobenzamides with Indoline-2,3-Dione in an Ionic Liquid in the Presence and Absence of Iodine. <i>Journal of Chemical Research</i> , 2012, 36, 157-161.	1.3	2
119	Iodine-Catalysed Synthesis of Thiopyrano[3,4-c]Quinoline Derivatives via Imino-Diels-Alder Reaction. <i>Journal of Chemical Research</i> , 2012, 36, 318-321.	1.3	5
120	An efficient method for the synthesis of naphthoquinoline derivatives catalyzed by iodine. <i>Heterocyclic Communications</i> , 2012, 18, 17-21.	1.2	3
121	Iodine-catalyzed synthesis of 5-arylanthra[2,1-c][2,7]naphthyridine derivatives via three-component reaction. <i>Heterocyclic Communications</i> , 2012, 18, .	1.2	0
122	Crystal structure of 2-amino-5,6-dihydro-7(4H)-benzothiazolone, C ₇ H ₈ N ₂ O ₂ . <i>Zeitschrift Fur Kristallographie - New Crystal Structures</i> , 2012, 227, 525-526.	0.3	0
123	Crystal structure of 2,3-dihydro-2-phenyl-3-[(E)-(1-phenylethylidene) amino]-4(1H)-quinazolinone, C ₂₂ H ₁₉ N ₃ O. <i>Zeitschrift Fur Kristallographie - New Crystal Structures</i> , 2012, 227, 499-500.	0.3	0
124	Crystal structure of (Z)-N-[[amino(pyridin-2-yl)-methylene]-2-hydroxybenzohydrazide, C ₁₃ H ₁₂ N ₄ O ₂ . <i>Zeitschrift Fur Kristallographie - New Crystal Structures</i> , 2012, 227, 533-534.	0.3	3
125	Ionic Liquid-Mediated One-Pot Synthesis of 5-(Trifluoromethyl)-4,7-dihydro-1,5-diazapyrimidine Derivatives. <i>Synthetic Communications</i> , 2012, 42, 2728-2738.	2.1	22
126	An Efficient Method for the Synthesis of 3-Aryl-4,7-dihydrophenanthroline Derivatives Catalyzed by Iodine. <i>Journal of Heterocyclic Chemistry</i> , 2012, 49, 1239-1242.	2.6	3

#	ARTICLE	IF	CITATIONS
127	Yb(OTf) ₃ : An Efficient Catalyst for the Synthesis of 11-Aryl-7-cyclopenta[<i>b</i>]phenanthroline-10-one Derivatives. <i>Journal of Heterocyclic Chemistry</i> , 2012, 49, 1439-1442.	2.6	7
128	Green Synthesis of Quinazolinone Derivatives Catalyzed by Iodine in Ionic Liquid. <i>Synthetic Communications</i> , 2012, 42, 341-349.	2.1	33
129	A Green Method for the Synthesis of Cyclopenta[<i>b</i>]chromen-9-one Derivatives in Ionic Liquids. <i>Journal of the Chinese Chemical Society</i> , 2012, 59, 650-654.	1.4	6
130	An Efficient Method for The Synthesis of 4-Arylfuro[2,3- <i>a</i>][4,7]phenanthroline Derivatives Catalyzed by Iodine. <i>Journal of Heterocyclic Chemistry</i> , 2012, 49, 585-588.	2.6	4
131	A Stereoselective Povarov Reaction Leading to <i>exo</i> -Tetrahydroindolo[3,2- <i>c</i>]quinoline Derivatives Catalyzed by Iodine. <i>European Journal of Organic Chemistry</i> , 2012, 2012, 4811-4818.	2.4	28
132	Diverse Confirmations in the Crystal Structures of 2,3-Dihydro-2,2-dimethylquinazolin-4(1 <i>H</i>)-one Derivatives. <i>Journal of Chemical Crystallography</i> , 2012, 42, 701-705.	1.1	1
133	A highly selective method for the synthesis of 1,3-diarylbenzo[<i>f</i>]quinoline derivatives catalyzed by silver triflate. <i>Monatshefte für Chemie</i> , 2012, 143, 935-938.	1.8	9
134	Combinatorial Synthesis of 3-Arylideneaminoquinazolin-4(1 <i>H</i>)-one Derivatives Catalyzed by Iodine in Ionic Liquids. <i>ACS Combinatorial Science</i> , 2011, 13, 196-199.	3.8	46
135	Three-Six-Membered Rings with Diverse Conformations in the Structure of 9-(4-Methoxyphenyl)-3,3-Dimethyl-10-(4-Methylphenyl)-1,2,3,4,5,6,7,8,9,10-Decahydroacridin-1,8-Dione. <i>Journal of Chemical Crystallography</i> , 2011, 41, 439-442.	1.1	3
136	Unclassical Hydrogen Bonds of C-H \cdots N and C-H \cdots Cl in the Crystal Structures of 2-((<i>E</i>)-1,3-diarylallylidene)malononitriles. <i>Journal of Chemical Crystallography</i> , 2011, 41, 59-63.	1.1	6
137	A green method for the synthesis of thiochromene derivatives in ionic liquids. <i>Journal of Heterocyclic Chemistry</i> , 2011, 48, 1056-1060.	2.6	15
138	Efficient method for the synthesis of 2-(3-arylbenzo[<i>f</i>]quinolin-2-yl)ethanol derivatives through an unusual ring-opening of THF-involved reaction. <i>Tetrahedron Letters</i> , 2011, 52, 612-614.	1.4	12
139	Synthesis of 7-Aryl-9-Methyl-3 <i>H</i> -Pyrazolo[4,3- <i>f</i>]Quinoline Derivatives Catalysed by Iodine. <i>Journal of Chemical Research</i> , 2011, 35, 513-515.	1.3	3
140	An efficient and highly selective method for the synthesis of cryptotackiene derivatives catalyzed by iodine. <i>Journal of Heterocyclic Chemistry</i> , 2010, 47, 873-877.	2.6	9
141	Yb(OTf) ₃ : an efficient catalyst for the synthesis of 3-arylbenzo[<i>f</i>]quinoline-1,2-dicarboxylate derivatives via imino-Diels-Alder reaction. <i>Tetrahedron Letters</i> , 2010, 51, 5721-5723.	1.4	33
142	Efficient and Green Method for the Synthesis of Highly Substituted Cyclohexadiene Derivatives in Aqueous Media. <i>Synthetic Communications</i> , 2010, 40, 1065-1073.	2.1	8
143	Efficient and Highly Selective Method for the Synthesis of Benzo(naphtho)quinoline Derivatives Catalyzed by Iodine. <i>ACS Combinatorial Science</i> , 2010, 12, 266-269.	3.3	50
144	Facile Method for the Combinatorial Synthesis of 2,2-Disubstituted Quinazolin-4(1 <i>H</i>)-one Derivatives Catalyzed by Iodine in Ionic Liquids. <i>ACS Combinatorial Science</i> , 2010, 12, 417-421.	3.3	90

#	ARTICLE	IF	CITATIONS
145	Divergent Products Obtained from the Reactions of Salicylaldehyde and 4-Hydroxycoumarin in TEBAc-H ₂ O, KF-Al ₂ O ₃ -EtOH, and Ionic Liquid. <i>Synthetic Communications</i> , 2010, 40, 3332-3345.	2.1	13
146	Synthesis of 2-Arylquinazolin-4(3 <i>H</i>)-one Derivatives Catalyzed by Iodine in [bmim] ⁺ []. <i>Synthetic Communications</i> , 2010, 40, 2633-2646.	2.1	40
147	Facile and Green Method for the Synthesis of β -Aminoketone Derivatives in Aqueous Media. <i>Synthetic Communications</i> , 2010, 40, 964-972.	2.1	3
148	An efficient synthesis of 1,3-diarylbenzo[<i>f</i>]quinolines from 2-halogenated acetophenone, aromatic aldehyde, and naphthalen-2-amine catalyzed by iodine. <i>Journal of Heterocyclic Chemistry</i> , 2009, 46, 1222-1228.	2.6	16
149	A novel and efficient method for the synthesis of 5-arylnaphtho[2,1- <i>c</i>][2,7]naphthyridine derivatives catalyzed by iodine. <i>Journal of Heterocyclic Chemistry</i> , 2009, 46, 1229-1234.	2.6	15
150	A novel and green method for the synthesis of highly substituted isoquinoline derivatives in ionic liquid. <i>Journal of Heterocyclic Chemistry</i> , 2009, 46, 1355-1363.	2.6	12
151	Efficient Method for the Synthesis of Pyranoquinoline, Thiopyranoquinoline, Thienoquinoline, and Naphtho[2,7]naphthyridine Derivatives Catalyzed by Iodine. <i>ACS Combinatorial Science</i> , 2009, 11, 433-437.	3.3	61
152	Iodine-Catalyzed Synthesis of 3-Arylbenzoquinoline Derivatives by Three-Component Reactions. <i>Synthetic Communications</i> , 2009, 39, 702-715.	2.1	15
153	Green Method for the Synthesis of Benzo[<i>f</i>]pyrimido[4,5- <i>b</i>]quinoline Derivatives Catalyzed by Iodine in Aqueous Media. <i>Synthetic Communications</i> , 2009, 39, 3069-3080.	2.1	17
154	Green Method for the Synthesis of Highly Substituted Cyclohexa-1,3-diene, Polyhydroindene, Polyhydronaphthalene, Isochromene, Isothiochromene, and Isoquinoline Derivatives in Ionic Liquids. <i>ACS Combinatorial Science</i> , 2009, 11, 1011-1022.	3.3	45
155	Malononitrile-Catalyzed and Highly Selective Method for the Synthesis of 2-((E)-1,3-Diaryllallylidene)malononitriles in Ionic Liquid. <i>Synthetic Communications</i> , 2009, 39, 3045-3059.	2.1	5
156	A Green Method for the Synthesis of Novel benzo[<i>b</i>]pyran Derivatives in an Ionic Liquid. <i>Journal of Chemical Research</i> , 2009, 2009, 234-236.	1.3	5
157	An efficient synthesis of polyhydroacridine derivatives by the three-component reaction of aldehydes, amines and dimedone in ionic liquid. <i>Journal of Heterocyclic Chemistry</i> , 2008, 45, 653-660.	2.6	53
158	An Efficient synthesis of pyrimido[4,5- <i>b</i>]quinoline and indeno[2- <i>a</i> ,1- <i>e</i> :5,6]pyrido[2,3- <i>i</i>]pyrimidine derivatives via multicomponent reactions in ionic liquid. <i>Journal of Heterocyclic Chemistry</i> , 2008, 45, 693-702.	2.6	61
159	I ₂ -catalyzed reactions of schiff base and alkyl aldehyde towards benzo[<i>f</i>]quinoline derivatives. <i>Journal of Heterocyclic Chemistry</i> , 2008, 45, 1027-1031.	2.6	9
160	An Efficient Method for the Synthesis of Benzo[<i>f</i>]quinoline and Benzo[<i>a</i>]phenanthridine Derivatives Catalyzed by Iodine by a Three-Component Reaction of Arenecarbaldehyde, Naphthalen-2-amine, and Cyclic Ketone. <i>European Journal of Organic Chemistry</i> , 2008, 2008, 3513-3518.	2.4	66
161	Simple Procedure for the Synthesis of 5,7-Diarylpyrido[2,3- <i>d</i>]pyrimidine Derivatives catalyzed by KF-Alumina. <i>Synthetic Communications</i> , 2008, 38, 1896-1908.	2.1	11
162	A Novel and Green Method for the Synthesis of Indeno[2,1- <i>c</i>]pyridine Derivatives in Ionic Liquid Catalyzed by Malononitrile. <i>Synlett</i> , 2008, 2008, 1185-1188.	1.8	28

#	ARTICLE	IF	CITATIONS
163	An Efficient and Highly Selective Method for the Synthesis of 3-Arylbenzoquinoline Derivatives Catalyzed by Iodine via Three-Component Reactions. <i>Synthesis</i> , 2008, 2008, 1902-1910.	2.3	47
164	An Unexpected Triethylbenzylammonium Chloride Catalyzed Ring Opening of 2-Pyrones in the Synthesis of 1-Arylbenzo[f]quinoline-2-carboxamide Derivatives in Aqueous Media. <i>Synlett</i> , 2007, 2007, 3141-3144.	1.8	14
165	Unexpected Spiro-benzoquinolines in the Reaction of N-(Arylidene)naphthalen-2-amine, Arylaldehyde, and 1,3-Dimethylbarbituric Acid in Water. <i>Chemistry Letters</i> , 2007, 36, 450-451.	1.3	8
166	A Clean Synthesis of 1,4-Diarylquinoline Derivatives Catalyzed by TEAC in Aqueous Media. <i>Journal of the Chinese Chemical Society</i> , 2007, 54, 1033-1039.	1.4	9
167	An Efficient Synthesis of Pyrazolo[3,4-b]pyridine Derivatives in Aqueous Media. <i>Journal of the Chinese Chemical Society</i> , 2007, 54, 1341-1345.	1.4	8
168	Novel N,N'-Diacylhydrazine-Based Colorimetric Receptors for Selective Sensing of Fluoride and Acetate Anions. <i>Chinese Journal of Chemistry</i> , 2007, 25, 973-976.	4.9	6
169	A new synthesis method for benzo[f]quinolin-3-carbonyl urea and thiourea derivatives in aqueous media catalyzed by TEAC. <i>Journal of Heterocyclic Chemistry</i> , 2007, 44, 441-447.	2.6	6
170	Three-component green synthesis of N-arylquinoline derivatives in ionic liquid [Bmim][BF ₄]: reactions of arylaldehyde, 3-arylamino-5,5-dimethylcyclohex-2-enone, and active methylene compounds. <i>Tetrahedron</i> , 2007, 63, 4439-4449.	1.9	89
171	An improved and clean procedure for the synthesis of one-donor poly-acceptors systems containing 2,6-dicyanoamine moiety in aqueous media catalyzed by TEAC in the presence and absence of K ₂ CO ₃ . <i>Tetrahedron</i> , 2007, 63, 5265-5273.	1.9	56
172	Synthesis and Crystal Structures of 3,3,6,6-tetramethyl-9-(2,4-dichlorophenyl)-3,4,6,7,9,10-hexahydro-2H,5H-acridine-1,8-dione and 3,3,6,6-tetramethyl-9,10-di(4-methoxyphenyl)-3,4,6,7,9,10-hexahydro-2H,5H-acridine-1,8-dione. <i>Journal of Chemical Crystallography</i> , 2007, 37, 483-487.	1.1	3
173	Unexpected Ring-Opening of a 2-Pyrone Ring in the Synthesis of 3-[(Z)-1-Hydroxy-3-Oxobut-1-Enyl]-2H-chromen-2-One Derivatives Catalysed by Kf-Alumina. <i>Journal of Chemical Research</i> , 2006, 2006, 602-604.	1.3	7
174	An improved synthesis of reduced 9-arylacridine-1,8-diones from 3-amino-5,5-dimethylcyclohex-2-enone, arylaldehydes and 1,3-dicarbonyl compounds in aqueous medium. <i>Journal of Chemical Research</i> , 2006, 2006, 719-721.	1.3	6
175	Synthesis of 5,7-Diarylpyrido[2,3-d]Pyrimidine Derivatives catalysed by Kf-Alumina. <i>Journal of Chemical Research</i> , 2006, 2006, 440-442.	1.3	1
176	Crystal Structure of 7-(4-Fluorophenyl)-5,6,7,14-tetrahydroquinolino[4,3-b]-benzo[f]quinolin-6-one N,N-Dimethylformamide Solvate. <i>Analytical Sciences: X-ray Structure Analysis Online</i> , 2006, 22, X125-X126.	0.1	0
177	The hydrogen bonding in 2-amino-3-cyano-4-(3-nitrophenyl)-4,6-dihydro-5H-pyrano[3,2-c]quinolin-5-one N,N-dimethylformamide solvate monohydrate. <i>Journal of Chemical Crystallography</i> , 2006, 36, 697-701.	1.1	2
178	A clean synthesis of polyhydroacridine and indenoquinoline derivatives catalyzed by triethylbenzylammonium chloride in aqueous media. <i>Journal of Heterocyclic Chemistry</i> , 2006, 43, 989-995.	2.6	18
179	One-Pot Synthesis of Pyrano[2,3-D]Pyrimidine Derivatives in Ionic Liquid Medium. <i>Journal of Chemical Research</i> , 2006, 2006, 157-159.	1.3	10
180	Synthesis of 3-Amino-1-Aryl-9-Methoxy-5,6-Dihydro-1H-Benzo[f]chromene-2-Carbonitriles in Aqueous Media. <i>Journal of Chemical Research</i> , 2006, 2006, 225-227.	1.3	2

#	ARTICLE	IF	CITATIONS
181	A Convenient and Clean Procedure for the Synthesis of Pyran Derivatives in Aqueous Media Catalysed by Tebac. <i>Journal of Chemical Research</i> , 2006, 2006, 228-230.	1.3	13
182	An Improved and Benign Synthesis of 9,10-Diarylacridine-1,8-dione and Indenoquinoline Derivatives from 3-Anilino-5,5-dimethylcyclohex-2-enones, Benzaldehydes, and 1,3-Dicarbonyl Compounds in an Ionic Liquid Medium. <i>Synthesis</i> , 2006, 2006, 4187-4199.	2.3	49
183	The crystal structure and unclassical pyran conformation of 2-amino-7-methyl-4-(3-nitrophenyl)-5-oxo-4H,5H-pyran [4,3-b]pyran-3-carbonitrile. <i>Journal of Chemical Research</i> , 2005, 2005, 775-777.	1.3	1
184	A Clean Procedure for the Synthesis of Chromeno[4,3-b]benzo[f]quinoline and Quinolino[4,3-b]benzo[f]quinoline Derivatives in Aqueous Media. <i>Chemistry Letters</i> , 2005, 34, 1316-1317.	1.3	17
185	A simple and clean procedure for the synthesis of polyhydroacridine and quinoline derivatives: reaction of Schiff base with 1,3-dicarbonyl compounds in aqueous medium. <i>Tetrahedron Letters</i> , 2005, 46, 7169-7173.	1.4	77
186	One-pot Synthesis of N-Hydroxyacridine Derivatives in Water. <i>Chinese Journal of Chemistry</i> , 2005, 23, 1223-1227.	4.9	27
187	The structure of 2-amino-3-cyano-4-(4-methylphenyl)-6-methoxy-1,4,9,10-tetrahydrobenzo[f]chromene. <i>Journal of Chemical Crystallography</i> , 2005, 35, 243-247.	1.1	2
188	Unclassical hydrogen bonds of C-H...S and C-H...N in the crystals of 2-amino-3-cyano-4-(3,4-dichlorophenyl)-5-oxo-1,4,5,6-tetrahydro-4H-pyrano[2,3-d]pyrimidine. <i>Journal of Chemical Crystallography</i> , 2005, 35, 999-1004.	1.1	2
189	One Pot Three Component Synthesis of 9-aryl polyhydroacridine Derivatives in an Ionic Liquid Medium. <i>Journal of Chemical Research</i> , 2005, 2005, 600-602.	1.3	29
190	Synthesis of 4H,5H-pyrano[3,2-c]pyrano-5-ones in aqueous media. <i>Journal of Chemical Research</i> , 2005, 2005, 724-726.	1.3	3
191	A Convenient Synthesis of 2-Amino-3-Cyano-4-Aryl-9,10-Dihydrobenzo[f] Chromene Derivatives Catalysed by KF/Al ₂ O ₃ . <i>Journal of Chemical Research</i> , 2004, 2004, 679-680.	1.3	3
192	Michael Addition Reaction of Malononitrile with α,β -Unsaturated Cycloketones Catalyzed by KF/Al ₂ O ₃ . <i>Chinese Journal of Chemistry</i> , 2004, 22, 122-125.	4.9	8
193	A Convenient Synthesis of 5-Oxo-5,6,7,8-tetrahydro-4H-benzo-[b]-pyran Derivatives Catalyzed by KF-Alumina. <i>Synthetic Communications</i> , 2003, 33, 119-126.	2.1	150
194	Synthesis of 2-aminochromene derivatives catalyzed by KF/Al ₂ O ₃ . <i>Chinese Journal of Chemistry</i> , 2003, 21, 1114-1117.	4.9	17
195	A CONVENIENT SYNTHESIS OF 2,4-DIARYLPOLYHYDROQUINOLINE DERIVATIONS IN THE PRESENCE OF AMMONIUM ACETATE. <i>Synthetic Communications</i> , 2002, 32, 3449-3454.	2.1	19
196	A concise synthesis of 10-benzoyl-3,4-dihydroanthracen-1(2H)-one derivatives catalyzed by TfOH under metal-free conditions. <i>Synthetic Communications</i> , 0, , 1-9.	2.1	1
197	CuI / L-Proline Catalyzed Synthesis of Bis(2-(4,5-diaryl-1H-imidazol-2-yl) phenyl)sulfane Derivatives Using Potassium Ethylxanthate as a Sulphur Source. <i>Journal of Heterocyclic Chemistry</i> , 0, , .	2.6	0