

Anton Kaufmann

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2435962/publications.pdf>

Version: 2024-02-01

67
papers

2,539
citations

186265

28
h-index

197818

49
g-index

68
all docs

68
docs citations

68
times ranked

2058
citing authors

#	ARTICLE	IF	CITATIONS
1	Improving the QuEChERS Liquid/Liquid Extraction of Analytes With Widely Varying Physicochemical Properties: Example of 201 Veterinary Drugs in Milk. <i>Journal of AOAC INTERNATIONAL</i> , 2022, 105, 1030-1042.	1.5	10
2	Simplifying Nontargeted Analysis of PFAS in Complex Food Matrixes. <i>Journal of AOAC INTERNATIONAL</i> , 2022, 105, 1280-1287.	1.5	11
3	High resolution mass spectrometry-based detection and quantification of Î²-agonists at relevant trace levels in a variety of animal-based food matrices. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2021, 38, 1350-1363.	2.3	8
4	Partially overlapping sequential window acquisition of all theoretical mass spectra: A methodology to improve the spectra quality of veterinary drugs present at low concentrations in highly complex biological matrices. <i>Rapid Communications in Mass Spectrometry</i> , 2020, 34, e8638.	1.5	2
5	The use of UHPLC, IMS, and HRMS in multiresidue analytical methods: A critical review. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2020, 1158, 122369.	2.3	11
6	Does the ion mobility resolving power as provided by commercially available ion mobility quadrupole time-of-flight mass spectrometry instruments permit the unambiguous identification of small molecules in complex matrices?. <i>Analytica Chimica Acta</i> , 2020, 1107, 113-126.	5.4	21
7	High-resolution mass spectrometry for bioanalytical applications: Is this the new gold standard?. <i>Journal of Mass Spectrometry</i> , 2020, 55, e4533.	1.6	36
8	High-resolution mass spectrometry-based multi-residue method covering relevant steroids, stilbenes and resorcylic acid lactones in a variety of animal-based matrices. <i>Analytica Chimica Acta</i> , 2019, 1054, 59-73.	5.4	17
9	Analysis of a variety of inorganic and organic additives in food products by ion-pairing liquid chromatography coupled to high-resolution mass spectrometry. <i>Analytical and Bioanalytical Chemistry</i> , 2018, 410, 5629-5640.	3.7	13
10	Coalescence and self-bunching observed in commercial high-resolution mass spectrometry instrumentation. <i>Rapid Communications in Mass Spectrometry</i> , 2018, 32, 503-515.	1.5	16
11	Analytical performance of the various acquisition modes in Orbitrap MS and MS/MS. <i>Journal of Mass Spectrometry</i> , 2018, 53, 725-738.	1.6	31
12	Easy and Fast Method for the Determination of Biogenic Amines in Fish and Fish Products with Liquid Chromatography Coupled to Orbitrap Tandem Mass Spectrometry. <i>Journal of AOAC INTERNATIONAL</i> , 2018, 101, 336-341.	1.5	25
13	Practical application of <i>in silico</i> fragmentation based residue screening with ion mobility high-resolution mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2017, 31, 1147-1157.	1.5	18
14	Using In Silico Fragmentation to Improve Routine Residue Screening in Complex Matrices. <i>Journal of the American Society for Mass Spectrometry</i> , 2017, 28, 2705-2715.	2.8	19
15	Comparison of linear intrascan and interscan dynamic ranges of Orbitrap and ion-mobility time-of-flight mass spectrometers. <i>Rapid Communications in Mass Spectrometry</i> , 2017, 31, 1915-1926.	1.5	34
16	Product ion isotopologue pattern: A tool to improve the reliability of elemental composition elucidations of unknown compounds in complex matrices. <i>Rapid Communications in Mass Spectrometry</i> , 2016, 30, 791-799.	1.5	5
17	Extension of the Q Orbitrap intrascan dynamic range by using a dedicated customized scan. <i>Rapid Communications in Mass Spectrometry</i> , 2016, 30, 1087-1095.	1.5	19
18	Nested data independent MS/MS acquisition. <i>Analytical and Bioanalytical Chemistry</i> , 2016, 408, 5031-5040.	3.7	8

#	ARTICLE	IF	CITATIONS
19	High throughput screening of animal urine samples: It is fast but is it also reliable?. <i>Drug Testing and Analysis</i> , 2016, 8, 491-497.	2.6	4
20	Improved performance of multiplexed targeted tandem mass spectrometry scans using customized Q Orbitrap data acquisition. <i>Rapid Communications in Mass Spectrometry</i> , 2016, 30, 1131-1138.	1.5	6
21	Determination of nitrofurans and chloramphenicol residues by high resolution mass spectrometry versus tandem quadrupole mass spectrometry. <i>Analytica Chimica Acta</i> , 2015, 862, 41-52.	5.4	89
22	Reliability of veterinary drug residue confirmation: High resolution mass spectrometry versus tandem mass spectrometry. <i>Analytica Chimica Acta</i> , 2015, 856, 54-67.	5.4	47
23	Determination of Corticosteroids, Anabolic Steroids, and Basic Nonsteroidal Anti-Inflammatory Drugs in Milk and Animal Tissues. <i>Journal of AOAC INTERNATIONAL</i> , 2014, 97, 263-272.	1.5	10
24	Signal suppression can bias selected reaction monitoring ratios. Implications for the confirmation of positive findings in residue testing. <i>Rapid Communications in Mass Spectrometry</i> , 2014, 28, 899-907.	1.5	8
25	Multi-residue quantification of veterinary drugs in milk with a novel extraction and cleanup technique: Salting out supported liquid extraction (SOSLE). <i>Analytica Chimica Acta</i> , 2014, 820, 56-68.	5.4	52
26	Quantitative analysis of polypeptide antibiotic residues in a variety of food matrices by liquid chromatography coupled to tandem mass spectrometry. <i>Analytica Chimica Acta</i> , 2013, 797, 81-88.	5.4	47
27	Evaluation of the interrelationship between mass resolving power and mass error tolerances for targeted bioanalysis using liquid chromatography coupled to high-resolution mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2013, 27, 347-356.	1.5	14
28	Post-run target screening strategy for ultra high performance liquid chromatography coupled to Orbitrap based veterinary drug residue analysis in animal urine. <i>Journal of Chromatography A</i> , 2013, 1292, 104-110.	3.7	27
29	Study of High-Resolution Mass Spectrometry Technology as a Replacement for Tandem Mass Spectrometry in the Field of Quantitative Pesticide Residue Analysis. <i>Journal of AOAC INTERNATIONAL</i> , 2012, 95, 528-548.	1.5	39
30	Determination of microbial transglutaminase in meat and meat products. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2012, 29, 1364-1373.	2.3	30
31	Determination of aminoglycoside residues by liquid chromatography and tandem mass spectrometry in a variety of matrices. <i>Analytica Chimica Acta</i> , 2012, 711, 46-53.	5.4	100
32	High Mass Resolution Versus MS/MS. <i>Comprehensive Analytical Chemistry</i> , 2012, 58, 169-215.	1.3	8
33	Accuracy of relative isotopic abundance and mass measurements in a single-stage orbitrap mass spectrometer. <i>Rapid Communications in Mass Spectrometry</i> , 2012, 26, 1081-1090.	1.5	40
34	The current role of high-resolution mass spectrometry in food analysis. <i>Analytical and Bioanalytical Chemistry</i> , 2012, 403, 1233-1249.	3.7	174
35	Semi-targeted residue screening in complex matrices with liquid chromatography coupled to high resolution mass spectrometry: current possibilities and limitations. <i>Analyst</i> , 2011, 136, 1898.	3.5	50
36	Quantification of anthelmintic drug residues in milk and muscle tissues by liquid chromatography coupled to Orbitrap and liquid chromatography coupled to tandem mass spectrometry. <i>Talanta</i> , 2011, 85, 991-1000.	5.5	64

#	ARTICLE	IF	CITATIONS
37	Development of an improved high resolution mass spectrometry based multi-residue method for veterinary drugs in various food matrices. <i>Analytica Chimica Acta</i> , 2011, 700, 86-94.	5.4	133
38	Quantitative and confirmative performance of liquid chromatography coupled to high-resolution mass spectrometry compared to tandem mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2011, 25, 979-992.	1.5	93
39	Comprehensive comparison of liquid chromatography selectivity as provided by two types of liquid chromatography detectors (high resolution mass spectrometry and tandem mass spectrometry): "Where is the crossover point?". <i>Analytica Chimica Acta</i> , 2010, 673, 60-72.	5.4	178
40	Strategy for the elucidation of elemental compositions of trace analytes based on a mass resolution of 100 000 full width at half maximum. <i>Rapid Communications in Mass Spectrometry</i> , 2010, 24, 2035-2045.	1.5	32
41	Post-interface signal suppression, a phenomenon observed in a single-stage Orbitrap mass spectrometer coupled to an electrospray interfaced liquid chromatograph. <i>Rapid Communications in Mass Spectrometry</i> , 2010, 24, 2162-2170.	1.5	52
42	<i>Veterinary Drugs.</i> , 2010, , 687-706.		0
43	Are liquid chromatography/electrospray tandem quadrupole fragmentation ratios unequivocal confirmation criteria?. <i>Rapid Communications in Mass Spectrometry</i> , 2009, 23, 985-998.	1.5	73
44	False-positive liquid chromatography/tandem mass spectrometric confirmation of sebuthylazine residues using the identification points system according to EU directive 2002/657/EC due to a biogenic insecticide in tarragon. <i>Rapid Communications in Mass Spectrometry</i> , 2009, 23, 1196-1200.	1.5	64
45	<i>Veterinary Drugs.</i> , 2009, , 713-733.		0
46	Quantitative multiresidue method for about 100 veterinary drugs in different meat matrices by sub 2- $\frac{1}{4}$ m particulate high-performance liquid chromatography coupled to time of flight mass spectrometry. <i>Journal of Chromatography A</i> , 2008, 1194, 66-79.	3.7	184
47	Ultra-performance liquid chromatography coupled to time of flight mass spectrometry (UPLC-TOF): A novel tool for multiresidue screening of veterinary drugs in urine. <i>Analytica Chimica Acta</i> , 2007, 586, 13-21.	5.4	133
48	Determination of the elemental composition of trace analytes in complex matrices using exact masses of product ions and corresponding neutral losses. <i>Rapid Communications in Mass Spectrometry</i> , 2007, 21, 2003-2013.	1.5	23
49	Strategies to avoid false negative findings in residue analysis using liquid chromatography coupled to time-of-flight mass spectrometry. <i>Rapid Communications in Mass Spectrometry</i> , 2006, 20, 3566-3572.	1.5	54
50	Segmented post-column analyte addition; a concept for continuous response control of liquid chromatography/mass spectrometry peaks affected by signal suppression/enhancement. <i>Rapid Communications in Mass Spectrometry</i> , 2005, 19, 611-617.	1.5	17
51	Quantitative liquid chromatography/tandem mass spectrometry determination of chloramphenicol residues in food using sub-2 μ m particulate high-performance liquid chromatography columns for sensitivity and speed. <i>Rapid Communications in Mass Spectrometry</i> , 2005, 19, 3694-3700.	1.5	33
52	Determination of 11 Aminoglycosides in Meat and Liver by Liquid Chromatography with Tandem Mass Spectrometry. <i>Journal of AOAC INTERNATIONAL</i> , 2005, 88, 1118-1125.	1.5	59
53	Analysis of polyphosphates in fish and shrimps tissues by two different ion chromatography methods: Implications on false-negative and -positive findings. <i>Food Additives and Contaminants</i> , 2005, 22, 1073-1082.	2.0	24
54	Determination of 11 aminoglycosides in meat and liver by liquid chromatography with tandem mass spectrometry. <i>Journal of AOAC INTERNATIONAL</i> , 2005, 88, 1118-25.	1.5	9

#	ARTICLE	IF	CITATIONS
55	Trace level quantification of streptomycin in honey with liquid chromatography/tandem mass spectrometry. Rapid Communications in Mass Spectrometry, 2003, 17, 2575-2577.	1.5	28
56	Quantitative LC/MS-MS Determination of Sulfonamides and Some Other Antibiotics in Honey. Journal of AOAC INTERNATIONAL, 2002, 85, 853-860.	1.5	79
57	Quantitative LC/MS-MS determination of sulfonamides and some other antibiotics in honey. Journal of AOAC INTERNATIONAL, 2002, 85, 853-60.	1.5	8
58	Comparison of different methods to determine polar compounds in frying oils. European Food Research and Technology, 2001, 213, 377-380.	3.3	8
59	HPLC with evaporative light scattering detection for the determination of polar compounds in used frying oils. European Food Research and Technology, 2001, 213, 372-376.	3.3	12
60	Multiresidue analysis of tranquilizers and the beta-blocker Carazolol in meat by liquid chromatography/tandem mass spectrometry. Rapid Communications in Mass Spectrometry, 2001, 15, 1747-1751.	1.5	17
61	Prevention of Vapor Overflow in Splitless Injection by a Novel Injector Design. Journal of High Resolution Chromatography, 1998, 21, 258-262.	1.4	9
62	Lead in wine. Food Additives and Contaminants, 1998, 15, 437-445.	2.0	38
63	Fully Automated Determination of Pesticides in Wine. Journal of AOAC INTERNATIONAL, 1997, 80, 1302-1307.	1.5	16
64	Multivariate Statistics as a Classification Tool in the Food Laboratory. Journal of AOAC INTERNATIONAL, 1997, 80, 665-675.	1.5	23
65	Determination of polar analytes in aqueous matrices by purge and trap. Journal of High Resolution Chromatography, 1997, 20, 10-16.	1.4	7
66	Maximum transfer condition for splitless injection. Journal of High Resolution Chromatography, 1997, 20, 193-200.	1.4	9
67	Capabilities and Limitations of High-Resolution Mass Spectrometry (HRMS): Time-of-flight and Orbitrap. Journal of High Resolution Chromatography, 1997, 20, 93-139.		7