

Yasumichi Kuwahara

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2461427/publications.pdf>

Version: 2024-02-01

28
papers

636
citations

623734

14
h-index

610901

24
g-index

30
all docs

30
docs citations

30
times ranked

1271
citing authors

#	ARTICLE	IF	CITATIONS
1	Suppression of malignant rhabdoid tumors through Chb ² -mediated RUNX1 inhibition. <i>Pediatric Blood and Cancer</i> , 2021, 68, e28789.	1.5	3
2	Mycosis fungoides in a patient with ulcerative colitis on anti-tumor necrosis factor-alpha therapy. <i>Clinical Journal of Gastroenterology</i> , 2021, 14, 170-175.	0.8	2
3	Frequent breakpoints of focal deletion and uniparental disomy in 22q11.1 or 11.2 segmental duplication region reveal distinct tumorigenesis in rhabdoid tumor of the kidney. <i>Genes Chromosomes and Cancer</i> , 2021, 60, 546-558.	2.8	0
4	Diverse outcomes in extra-cranial rhabdoid tumors: A single institute experience. <i>Pediatric Hematology and Oncology</i> , 2021, , 1-8.	0.8	0
5	Tumor necrosis factor-related apoptosis-inducing ligand is a novel transcriptional target of runt-related transcription factor A1. <i>International Journal of Oncology</i> , 2021, 60, .	3.3	4
6	The Novel Histone Deacetylase Inhibitor, OBP-801, Induces Apoptosis in Rhabdoid Tumors by Releasing the Silencing of <i>NOXA</i> . <i>Molecular Cancer Therapeutics</i> , 2020, 19, 1992-2000.	4.1	5
7	Oncogenic role of HMGA2 in fusion-negative rhabdomyosarcoma cells. <i>Cancer Cell International</i> , 2020, 20, 192.	4.1	7
8	Detection of circulating fungal DNA by polymerase chain reaction in a fatal case of <i>Cunninghamella bertholletiae</i> infection. <i>IDCases</i> , 2020, 20, e00760.	0.9	5
9	Novel Two MRT Cell Lines Established from Multiple Sites of a Synchronous MRT Patient. <i>Anticancer Research</i> , 2020, 40, 6159-6170.	1.1	0
10	High Frequency of Ovarian Cyst Development in <i>Vhl</i> / <i>Snf5</i> Mice. <i>American Journal of Pathology</i> , 2018, 188, 1510-1516.	3.8	0
11	Therapeutic targeting of PGBD5-induced DNA repair dependency in pediatric solid tumors. <i>Science Translational Medicine</i> , 2017, 9, .	12.4	48
12	A <i>NOXA</i> / <i>MCL1</i> Imbalance Underlies Chemoresistance of Malignant Rhabdoid Tumor Cells. <i>Journal of Cellular Physiology</i> , 2016, 231, 1932-1940.	4.1	11
13	Residual tumor in cases of intermediate-risk neuroblastoma did not influence the prognosis. <i>Japanese Journal of Clinical Oncology</i> , 2016, 46, 661-666.	1.3	12
14	Diffuse Anterior Retinoblastoma with Sarcoidosis-Like Nodule. <i>Case Reports in Ophthalmology</i> , 2015, 6, 443-447.	0.7	10
15	The Chromatin-Modifying Protein HMGA2 Promotes Atypical Teratoid/Rhabdoid Cell Tumorigenicity. <i>Journal of Neuropathology and Experimental Neurology</i> , 2015, 74, 177-185.	1.7	26
16	Disrupting LIN28 in atypical teratoid rhabdoid tumors reveals the importance of the mitogen activated protein kinase pathway as a therapeutic target. <i>Oncotarget</i> , 2015, 6, 3165-3177.	1.8	66
17	<i>SNF5</i> / <i>INI1</i> Deficiency Redefines Chromatin Remodeling Complex Composition during Tumor Development. <i>Molecular Cancer Research</i> , 2014, 12, 1574-1585.	3.4	31
18	<i>SNF5</i> Reexpression in Malignant Rhabdoid Tumors Regulates Transcription of Target Genes by Recruitment of <i>SWI</i> / <i>SNF</i> Complexes and <i>RNAPII</i> to the Transcription Start Site of Their Promoters. <i>Molecular Cancer Research</i> , 2013, 11, 251-260.	3.4	33

#	ARTICLE	IF	CITATIONS
19	Establishment and characterization of MRT cell lines from genetically engineered mouse models and the influence of genetic background on their development. <i>International Journal of Cancer</i> , 2013, 132, 2767-2777.	5.1	3
20	Sensitivity of malignant rhabdoid tumor cell lines to PD 0332991 is inversely correlated with p16 expression. <i>Biochemical and Biophysical Research Communications</i> , 2011, 413, 62-68.	2.1	36
21	Reexpression of hSNF5 in Malignant Rhabdoid Tumor Cell Lines Causes Cell Cycle Arrest through a p21CIP1/WAF1-Dependent Mechanism. <i>Cancer Research</i> , 2010, 70, 1854-1865.	0.9	40
22	Inactivation of SNF5 cooperates with p53 loss to accelerate tumor formation in Snf5 ^{+/+} ; p53 ^{+/+} mice. <i>Molecular Carcinogenesis</i> , 2009, 48, 1139-1148.	2.7	23
23	Trastuzumab Activates Allogeneic or Autologous Antibody-Dependent Cellular Cytotoxicity against Malignant Rhabdoid Tumor Cells and Interleukin-2 Augments the Cytotoxicity. <i>Clinical Cancer Research</i> , 2008, 14, 1192-1199.	7.0	21
24	Induction of apoptosis by an inhibitor of EGFR in neuroblastoma cells. <i>Biochemical and Biophysical Research Communications</i> , 2007, 358, 226-232.	2.1	33
25	Establishment of a cell line from a malignant rhabdoid tumor of the liver lacking the function of two tumor suppressor genes, hSNF5/IN1 and p16. <i>Cancer Genetics and Cytogenetics</i> , 2005, 158, 172-179.	1.0	16
26	Prediction of MYC Amplification in Neuroblastoma Using Serum DNA and Real-Time Quantitative Polymerase Chain Reaction. <i>Journal of Clinical Oncology</i> , 2005, 23, 5205-5210.	1.6	89
27	Antitumor Activity of Gefitinib in Malignant Rhabdoid Tumor Cells In vitro and In vivo. <i>Clinical Cancer Research</i> , 2004, 10, 5940-5948.	7.0	42
28	Fenretinide induces sustained-activation of JNK/p38 MAPK and apoptosis in a reactive oxygen species-dependent manner in neuroblastoma cells. <i>International Journal of Cancer</i> , 2004, 112, 219-224.	5.1	69