
Jenni Hällfors

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2464841/publications.pdf Version: 2024-02-01



IENNI HÃO FORS

#	Article	IF	CITATIONS
1	Is Brain Derived Neurotrophic Factor (Bdnf) Associated With Smoking Initiation? Replication Using a Large Finnish Population Sample. Nicotine and Tobacco Research, 2020, 22, 293-296.	1.4	5
2	Genome-Wide Meta-Analyses of FTND and TTFC Phenotypes. Nicotine and Tobacco Research, 2020, 22, 900-909.	1.4	17
3	Genomeâ€wide association study in Finnish twins highlights the connection between nicotine addiction and neurotrophin signaling pathway. Addiction Biology, 2019, 24, 549-561.	1.4	17
4	<i>KLB</i> is associated with alcohol drinking, and its gene product β-Klotho is necessary for FGF21 regulation of alcohol preference. Proceedings of the National Academy of Sciences of the United States of America, 2016, 113, 14372-14377.	3.3	208
5	A Genome-Wide Association Study of a Biomarker of Nicotine Metabolism. PLoS Genetics, 2015, 11, e1005498.	1.5	107
6	Stratification by Smoking Status Reveals an Association of CHRNA5-A3-B4 Genotype with Body Mass Index in Never Smokers. PLoS Genetics, 2014, 10, e1004799.	1.5	45
7	Genetics and Smoking. Current Addiction Reports, 2014, 1, 75-82.	1.6	47
8	Distinct Loci in the <i>CHRNA5</i> / <i>CHRNA3</i> / <i>CHRNB4</i> Gene Cluster Are Associated With Onset of Regular Smoking. Genetic Epidemiology, 2013, 37, 846-859.	0.6	32
9	Scrutiny of the CHRNA5-CHRNA3-CHRNB4 smoking behavior locus reveals a novel association with alcohol use in a Finnish population based study. International Journal of Molecular Epidemiology and Genetics, 2013, 4, 109-19.	0.4	18
10	Increased Genetic Vulnerability to Smoking at CHRNA5 in Early-Onset Smokers. Archives of General Psychiatry, 2012, 69, 854.	13.8	71