

Jian-Yong Shao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2477006/publications.pdf>

Version: 2024-02-01

60
papers

2,883
citations

279798

23
h-index

168389

53
g-index

60
all docs

60
docs citations

60
times ranked

4353
citing authors

#	ARTICLE	IF	CITATIONS
1	Plasma SCIP1 methylation in diagnosis and prognosis prediction in hepatocellular carcinoma. <i>Neoplasma</i> , 2021, 68, 62-70.	1.6	5
2	The Diagnostic Value of Serum PIVKA-II Alone or in Combination with AFP in Chinese Hepatocellular Carcinoma Patients. <i>Disease Markers</i> , 2021, 2021, 1-9.	1.3	19
3	Diagnostic and Prognostic Characteristics of Circulating Free DNA Methylation Detected by the Electrochemical Method in Malignant Tumors. <i>Cancers</i> , 2021, 13, 664.	3.7	3
4	Performance of common genetic variants in risk prediction for colorectal cancer in Chinese: A two-stage and multicenter study. <i>Genomics</i> , 2021, 113, 867-873.	2.9	1
5	Classification of gastric cancer by EBV status combined with molecular profiling predicts patient prognosis. <i>Clinical and Translational Medicine</i> , 2020, 10, 353-362.	4.0	13
6	Identification of a Gene-Related Risk Signature in Melanoma Patients Using Bioinformatic Profiling. <i>Journal of Oncology</i> , 2020, 2020, 1-13.	1.3	8
7	Risk stratification for nasopharyngeal carcinoma: a real-world study based on locoregional extension patterns and Epstein-Barr virus DNA load. <i>Therapeutic Advances in Medical Oncology</i> , 2020, 12, 175883592093205.	3.2	17
8	Development of a Nomogram Model for Treatment of Nonmetastatic Nasopharyngeal Carcinoma. <i>JAMA Network Open</i> , 2020, 3, e2029882.	5.9	31
9	Development and validation of a prognostic nomogram for the pre-treatment prediction of early metachronous metastasis in endemic nasopharyngeal carcinoma: a big-data intelligence platform-based analysis. <i>Therapeutic Advances in Medical Oncology</i> , 2020, 12, 175883592097813.	3.2	6
10	The Percentage of Anaplastic Lymphoma Kinase-Positive Tumor Cells Has Clinical Implications for Patients with Non-Small Cell Lung Cancer. <i>Genetic Testing and Molecular Biomarkers</i> , 2019, 23, 589-597.	0.7	3
11	Prognostic Implications of Tripartite Motif Containing 24 Expression Levels in Patients with Solid Tumors: A Systematic Review and Meta-Analysis. <i>Genetic Testing and Molecular Biomarkers</i> , 2019, 23, 473-479.	0.7	1
12	<p>Mutation spectrum of germline cancer susceptibility genes among unselected Chinese colorectal cancer patients</p>. <i>Cancer Management and Research</i> , 2019, Volume 11, 3721-3739.	1.9	15
13	Past and Recent Salted Fish and Preserved Food Intakes Are Weakly Associated with Nasopharyngeal Carcinoma Risk in Adults in Southern China. <i>Journal of Nutrition</i> , 2019, 149, 1596-1605.	2.9	25
14	Interfering Expression of Chimeric Transcript<i> SEPT7P2-PSPH</i> Promotes Cell Proliferation in Patients with Nasopharyngeal Carcinoma. <i>Journal of Oncology</i> , 2019, 2019, 1-10.	1.3	9
15	Plasma Epstein-Barr Virus DNA Load After Induction Chemotherapy Predicts Outcome in Locoregionally Advanced Nasopharyngeal Carcinoma. <i>International Journal of Radiation Oncology Biology Physics</i> , 2019, 104, 355-361.	0.8	64
16	Prognostic implications of a molecular classifier derived from wholeâexome sequencing in nasopharyngeal carcinoma. <i>Cancer Medicine</i> , 2019, 8, 2705-2716.	2.8	13
17	Medical History, Medication Use, and Risk of Nasopharyngeal Carcinoma. <i>American Journal of Epidemiology</i> , 2018, 187, 2117-2125.	3.4	20
18	SPP1 rs4754 and its epistatic interactions with SPARC polymorphisms in gastric cancer susceptibility. <i>Gene</i> , 2018, 640, 43-50.	2.2	19

#	ARTICLE	IF	CITATIONS
19	Hypermethylation of APC2 Is a Predictive Epigenetic Biomarker for Chinese Colorectal Cancer. <i>Disease Markers</i> , 2018, 2018, 1-7.	1.3	9
20	Pretreatment quality of life as a predictor of survival for patients with nasopharyngeal carcinoma treated with IMRT. <i>BMC Cancer</i> , 2018, 18, 114.	2.6	13
21	The Prognostic Value of Treatment-Related Lymphopenia in Nasopharyngeal Carcinoma Patients. <i>Cancer Research and Treatment</i> , 2018, 50, 19-29.	3.0	56
22	Serum microRNA profiles as diagnostic biomarkers for HBV-positive hepatocellular carcinoma. <i>Liver International</i> , 2017, 37, 888-896.	3.9	56
23	Quantification of familial risk of nasopharyngeal carcinoma in a high-incidence area. <i>Cancer</i> , 2017, 123, 2716-2725.	4.1	54
24	Active and Passive Smoking and Risk of Nasopharyngeal Carcinoma: A Population-Based Case-Control Study in Southern China. <i>American Journal of Epidemiology</i> , 2017, 185, 1272-1280.	3.4	68
25	SPINK6 Promotes Metastasis of Nasopharyngeal Carcinoma via Binding and Activation of Epithelial Growth Factor Receptor. <i>Cancer Research</i> , 2017, 77, 579-589.	0.9	47
26	Concurrent chemoradiotherapy with or without cetuximab for stage II to IVb nasopharyngeal carcinoma: a case-control study. <i>BMC Cancer</i> , 2017, 17, 567.	2.6	29
27	Development of a population-based cancer case-control study in southern china. <i>Oncotarget</i> , 2017, 8, 87073-87085.	1.8	29
28	Implication of comorbidity on the initiation of chemotherapy and survival outcomes in patients with locoregionally advanced nasopharyngeal carcinoma. <i>Oncotarget</i> , 2017, 8, 10594-10601.	1.8	5
29	Predictors of Mastoiditis after Intensity-Modulated Radiotherapy in Nasopharyngeal Carcinoma: A Dose-Volume Analysis. <i>Journal of Cancer</i> , 2016, 7, 276-282.	2.5	4
30	Induction chemotherapy followed by concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone in stage III-IVb nasopharyngeal carcinoma patients with Epstein-Barr virus DNA ≥ 4000 copies/ml: a matched study. <i>Oncotarget</i> , 2016, 7, 29739-29748.	1.8	15
31	Oral Hygiene and Risk of Nasopharyngeal Carcinoma—A Population-Based Case-Control Study in China. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2016, 25, 1201-1207.	2.5	46
32	With or without reirradiation in advanced local recurrent nasopharyngeal carcinoma: a case-control study. <i>BMC Cancer</i> , 2016, 16, 774.	2.6	17
33	Comparison of KRAS mutation status between primary tumor and metastasis in Chinese colorectal cancer patients. <i>Medical Oncology</i> , 2016, 33, 71.	2.5	12
34	A new prognostic histopathologic classification of nasopharyngeal carcinoma. <i>Chinese Journal of Cancer</i> , 2016, 35, 41.	4.9	83
35	Serum microRNA profiles as prognostic biomarkers for HBV-positive hepatocellular carcinoma. <i>Oncotarget</i> , 2016, 7, 45637-45648.	1.8	22
36	Plasma Epstein-Barr viral DNA complements TNM classification of nasopharyngeal carcinoma in the era of intensity-modulated radiotherapy. <i>Oncotarget</i> , 2016, 7, 6221-6230.	1.8	37

#	ARTICLE	IF	CITATIONS
37	LOX expression in primary nasopharyngeal carcinoma: correlation with prognostic parameters and outcome. <i>Oncotarget</i> , 2016, 7, 8200-8207.	1.8	14
38	Prognostic effect of pregnancy on young female patients with nasopharyngeal carcinoma: results from a matched cohort analysis. <i>Oncotarget</i> , 2016, 7, 21913-21921.	1.8	4
39	Neoadjuvant chemotherapy in locally advanced nasopharyngeal carcinoma: Defining high-risk patients who may benefit before concurrent chemotherapy combined with intensity-modulated radiotherapy. <i>Scientific Reports</i> , 2015, 5, 16664.	3.3	34
40	Hepatitis B virus reactivation and hepatitis in diffuse large B-cell lymphoma patients with resolved hepatitis B receiving rituximab-containing chemotherapy: risk factors and survival. <i>Chinese Journal of Cancer</i> , 2015, 34, 225-34.	4.9	21
41	Clinicopathologic characteristics and therapeutic responses of Chinese patients with non-small cell lung cancer who harbor an anaplastic lymphoma kinase rearrangement. <i>Chinese Journal of Cancer</i> , 2015, 34, 404-12.	4.9	12
42	Identification of surrogate endpoints in patients with locoregionally advanced nasopharyngeal carcinoma receiving neoadjuvant chemotherapy plus concurrent chemoradiotherapy versus concurrent chemoradiotherapy alone. <i>BMC Cancer</i> , 2015, 15, 930.	2.6	6
43	The impact of the cumulative dose of cisplatin during concurrent chemoradiotherapy on the clinical outcomes of patients with advanced-stage nasopharyngeal carcinoma in an era of intensity-modulated radiotherapy. <i>BMC Cancer</i> , 2015, 15, 977.	2.6	21
44	Incidence of and Risk Factors for Mastoiditis after Intensity Modulated Radiotherapy in Nasopharyngeal Carcinoma. <i>PLoS ONE</i> , 2015, 10, e0131284.	2.5	4
45	High expression of Talin-1 is associated with poor prognosis in patients with nasopharyngeal carcinoma. <i>BMC Cancer</i> , 2015, 15, 332.	2.6	21
46	The Prognostic Value of Plasma Epstein-Barr Viral DNA and Tumor Response to Neoadjuvant Chemotherapy in Advanced-Stage Nasopharyngeal Carcinoma. <i>International Journal of Radiation Oncology Biology Physics</i> , 2015, 93, 862-869.	0.8	110
47	Comparison of the treatment outcomes of intensity-modulated radiotherapy and two-dimensional conventional radiotherapy in nasopharyngeal carcinoma patients with parapharyngeal space extension. <i>Radiotherapy and Oncology</i> , 2015, 116, 167-173.	0.6	14
48	Genome-Wide Identification of a Methylation Gene Panel as a Prognostic Biomarker in Nasopharyngeal Carcinoma. <i>Molecular Cancer Therapeutics</i> , 2015, 14, 2864-2873.	4.1	80
49	The Frequency and Clinical Implication of ROS1 and RET Rearrangements in Resected Stage IIIA-N2 Non-Small Cell Lung Cancer Patients. <i>PLoS ONE</i> , 2015, 10, e0124354.	2.5	17
50	TEL2 suppresses metastasis by down-regulating SERPINE1 in nasopharyngeal carcinoma. <i>Oncotarget</i> , 2015, 6, 29240-29253.	1.8	39
51	IDH1 mutation detection by droplet digital PCR in glioma. <i>Oncotarget</i> , 2015, 6, 39651-39660.	1.8	18
52	Inhibition of MiR-155 suppresses cell migration in nasopharyngeal carcinoma through targeting ZDHHC2. <i>International Journal of Clinical and Experimental Medicine</i> , 2015, 8, 8472-84.	1.3	7
53	Comparison of Long-Term Survival and Toxicity of Cisplatin Delivered Weekly versus Every Three Weeks Concurrently with Intensity-Modulated Radiotherapy in Nasopharyngeal Carcinoma. <i>PLoS ONE</i> , 2014, 9, e110765.	2.5	31
54	MicroRNA-30a promotes invasiveness and metastasis <i>in vitro</i> and <i>in vivo</i> through epithelial-mesenchymal transition and results in poor survival of nasopharyngeal carcinoma patients. <i>Experimental Biology and Medicine</i> , 2014, 239, 891-898.	2.4	29

#	ARTICLE	IF	CITATIONS
55	Knockdown of miR-214 Promotes Apoptosis and Inhibits Cell Proliferation in Nasopharyngeal Carcinoma. PLoS ONE, 2014, 9, e86149.	2.5	62
56	<i>ALK</i> gene copy number gain and its clinical significance in hepatocellular carcinoma. World Journal of Gastroenterology, 2014, 20, 183.	3.3	13
57	Plasma Epstein-Barr virus DNA level strongly predicts survival in metastatic/recurrent nasopharyngeal carcinoma treated with palliative chemotherapy. Cancer, 2011, 117, 3750-3757.	4.1	134
58	Eight-Signature Classifier for Prediction of Nasopharyngeal Carcinoma Survival. Journal of Clinical Oncology, 2011, 29, 4516-4525.	1.6	131
59	MicroRNA miR-21 overexpression in human breast cancer is associated with advanced clinical stage, lymph node metastasis and patient poor prognosis. Rna, 2008, 14, 2348-2360.	3.5	993
60	Comparison of plasma Epstein-Barr virus (EBV) DNA levels and serum EBV immunoglobulin A/virus capsid antigen antibody titers in patients with nasopharyngeal carcinoma. Cancer, 2004, 100, 1162-1170.	4.1	194