

Caroline Goupille

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2479516/publications.pdf>

Version: 2024-02-01

44
papers

1,113
citations

430874

18
h-index

414414

32
g-index

47
all docs

47
docs citations

47
times ranked

1675
citing authors

#	ARTICLE	IF	CITATIONS
1	Sensitization by docosahexaenoic acid (DHA) of breast cancer cells to anthracyclines through loss of glutathione peroxidase (GPx1) response. <i>Free Radical Biology and Medicine</i> , 2008, 44, 1483-1491.	2.9	101
2	Cancer cachexia is associated with a decrease in skeletal muscle mitochondrial oxidative capacities without alteration of ATP production efficiency. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2012, 3, 265-275.	7.3	89
3	Increase of rat colon carcinoma cells tumorigenicity by β 1,2 fucosyltransferase gene transfection. <i>Glycobiology</i> , 1997, 7, 221-229.	2.5	69
4	Sensitization by Dietary Docosahexaenoic Acid of Rat Mammary Carcinoma to Anthracycline: A Role for Tumor Vascularization. <i>Clinical Cancer Research</i> , 2006, 12, 5879-5886.	7.0	67
5	β 1,2Fucosyltransferase increases resistance to apoptosis of rat colon carcinoma cells. <i>Glycobiology</i> , 2000, 10, 375-382.	2.5	63
6	CCL5-enhanced human immature dendritic cell migration through the basement membrane in vitro depends on matrix metalloproteinase-9. <i>Journal of Leukocyte Biology</i> , 2006, 79, 767-778.	3.3	60
7	Long chain n-3 polyunsaturated fatty acids increase the efficacy of docetaxel in mammary cancer cells by downregulating Akt and PKC δ / β -induced ERK pathways. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2016, 1861, 380-390.	2.4	50
8	Efficiency of oxidative phosphorylation in liver mitochondria is decreased in a rat model of peritoneal carcinosis. <i>Journal of Hepatology</i> , 2011, 54, 320-327.	3.7	44
9	Activation of TRPV2 and BKCa channels by the LL-37 enantiomers stimulates calcium entry and migration of cancer cells. <i>Oncotarget</i> , 2016, 7, 23785-23800.	1.8	44
10	Sodium Channel Nav1.5 Controls Epithelial-to-Mesenchymal Transition and Invasiveness in Breast Cancer Cells Through its Regulation by the Salt-Inducible Kinase-1. <i>Scientific Reports</i> , 2019, 9, 18652.	3.3	43
11	ATP-dependent activity and mitochondrial localization of drug efflux pumps in doxorubicin-resistant breast cancer cells. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2017, 1861, 1075-1084.	2.4	40
12	Lipid metabolism and Calcium signaling in epithelial ovarian cancer. <i>Cell Calcium</i> , 2019, 81, 38-50.	2.4	36
13	IL-7 receptor is present on human microvascular endothelial cells. <i>Immunology Letters</i> , 2003, 86, 163-168.	2.5	30
14	α -Tocopherol Suppresses Mammary Tumor Sensitivity to Anthracyclines in Fish Oil-Fed Rats. <i>Nutrition and Cancer</i> , 2005, 51, 178-183.	2.0	28
15	Regulation of hepatic cardiolipin metabolism by TNF α : Implication in cancer cachexia. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2015, 1851, 1490-1500.	2.4	28
16	PPAR γ mRNA expression, reduced by n ω -3 PUFA diet in mammary tumor, controls breast cancer cell growth. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2013, 1831, 1618-1625.	2.4	26
17	Sulfated Glycoaminoglycans and Proteoglycan Syndecan-4 Are Involved in Membrane Fixation of LL-37 and Its Pro-Migratory Effect in Breast Cancer Cells. <i>Biomolecules</i> , 2019, 9, 481.	4.0	22
18	Increased tumorigenicity of rat colon carcinoma cells after β 1,2-fucosyltransferase anti-sense cDNA transfection. , 1999, 80, 606-611.		21

#	ARTICLE	IF	CITATIONS
19	Role for α 1,2-fucosyltransferase and histo-blood group antigen H type 2 in resistance of rat colon carcinoma cells to 5-fluorouracil. <i>International Journal of Cancer</i> , 2000, 85, 142-148.	5.1	20
20	The neonatal Fc receptor in cancer FcRn in cancer. <i>Cancer Medicine</i> , 2020, 9, 4736-4742.	2.8	19
21	Reducing endothelial NOS activation and interstitial fluid pressure with n ω -3 PUFA offset tumor chemoresistance. <i>Carcinogenesis</i> , 2012, 33, 260-267.	2.8	18
22	N-3 Polyunsaturated Fatty Acids of Marine Origin and Multifocality in Human Breast Cancer. <i>PLoS ONE</i> , 2016, 11, e0147148.	2.5	18
23	Role for α 1,2-fucosyltransferase and histo-blood group antigen H type 2 in resistance of rat colon carcinoma cells to 5-fluorouracil. <i>International Journal of Cancer</i> , 2000, 85, 142.	5.1	18
24	Low eicosapentaenoic acid and gamma-linolenic acid levels in breast adipose tissue are associated with inflammatory breast cancer. <i>Breast</i> , 2019, 45, 113-117.	2.2	17
25	Non-invasive quantification of tumor vascular architecture during docetaxel-chemotherapy. <i>Breast Cancer Research and Treatment</i> , 2012, 134, 1013-1025.	2.5	13
26	EPA and DHA Fatty Acids Induce a Remodeling of Tumor Vasculature and Potentiate Docetaxel Activity. <i>International Journal of Molecular Sciences</i> , 2020, 21, 4965.	4.1	13
27	Expression of A and H blood-group and of CD44 antigens during chemical rat colonic carcinogenesis. <i>Glycoconjugate Journal</i> , 1997, 14, 801-808.	2.7	12
28	Susceptibility of rat colon carcinoma cells to lymphokine activated killer-mediated cytotoxicity is decreased by α 1,2-fucosylation. , 2000, 86, 713-717.		12
29	N-3 PUFA-Enriched Diet Delays the Occurrence of Cancer Cachexia in Rat With Peritoneal Carcinosis. <i>Nutrition and Cancer</i> , 2010, 62, 343-350.	2.0	12
30	NMR-Based Lipidomic Approach To Evaluate Controlled Dietary Intake of Lipids in Adipose Tissue of a Rat Mammary Tumor Model. <i>Journal of Proteome Research</i> , 2016, 15, 868-878.	3.7	12
31	Increased BRCA1 protein in mammary tumours of rats fed marine ω -3 fatty acids. <i>Oncology Reports</i> , 2007, 17, 713.	2.6	11
32	A 1D High Performance Thin Layer Chromatography Method Validated to Quantify Phospholipids Including Cardiolipin and Monolysocardiolipin from Biological Samples. <i>European Journal of Lipid Science and Technology</i> , 2020, 122, 1900240.	1.5	10
33	Detection of a potential receptor for the H-blood-group antigen on rat colon-carcinoma cells and normal tissues. , 1998, 76, 136-140.		9
34	Low Levels of Omega-3 Long-Chain Polyunsaturated Fatty Acids Are Associated with Bone Metastasis Formation in Premenopausal Women with Breast Cancer: A Retrospective Study. <i>Nutrients</i> , 2020, 12, 3832.	4.1	8
35	Potassium and Calcium Channel Complexes as Novel Targets for Cancer Research. <i>Reviews of Physiology, Biochemistry and Pharmacology</i> , 2020, , 157-176.	1.6	6
36	Dietary docosahexaenoic acid proposed to sensitize breast tumors to locally delivered drug. <i>Clinical Lipidology</i> , 2010, 5, 233-243.	0.4	5

#	ARTICLE	IF	CITATIONS
37	Development of a Novel High-Performance Thin Layer Chromatography-Based Method for the Simultaneous Quantification of Clinically Relevant Lipids from Cells and Tissue Extracts. <i>Lipids</i> , 2020, 55, 403-412.	1.7	5
38	Accumulation of Arachidonic Acid, Precursor of Pro-Inflammatory Eicosanoids, in Adipose Tissue of Obese Women: Association with Breast Cancer Aggressiveness Indicators. <i>Biomedicines</i> , 2022, 10, 995.	3.2	5
39	Identification of a Positive Association between Mammary Adipose Cholesterol Content and Indicators of Breast Cancer Aggressiveness in a French Population. <i>Journal of Nutrition</i> , 2021, 151, 1119-1127.	2.9	3
40	A rat experimental model for the design of vaccines against tumor associated antigens Tn and Sialyl-Tn. <i>Glycoconjugate Journal</i> , 1999, 16, 681-684.	2.7	2
41	Total long-chain polyunsaturated n-3 fatty acids level is an independent predictive factor of breast cancer multifocality in women with positive hormone-receptors tumors. <i>Surgical Oncology</i> , 2021, 38, 101597.	1.6	2
42	S12.11 Involvement of histo-blood group antigens in the susceptibility of colon carcinoma cells to natural killer-mediated cytotoxicity. <i>Glycoconjugate Journal</i> , 1993, 10, 298-298.	2.7	0
43	Natural isotopic abundances as markers of compliance in clinical trials. <i>American Journal of Clinical Nutrition</i> , 2020, 111, 1109-1110.	4.7	0
44	Abstract 4100: Nutrition and breast cancer prevention: adipose tissue proton magnetic resonance spectroscopy (1H-NMR) as biomarker of past dietary intake of lipids. , 2014, , .		0