Liling Zhang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2489709/publications.pdf

Version: 2024-02-01

567281 315739 1,931 41 15 38 citations h-index g-index papers 47 47 47 3241 docs citations times ranked citing authors all docs

#	Article	IF	Citations
1	Detection and HER2 Expression of Circulating Tumor Cells: Prospective Monitoring in Breast Cancer Patients Treated in the Neoadjuvant GeparQuattro Trial. Clinical Cancer Research, 2010, 16, 2634-2645.	7.0	463
2	Meta-Analysis of the Prognostic Value of Circulating Tumor Cells in Breast Cancer. Clinical Cancer Research, 2012, 18, 5701-5710.	7.0	330
3	The role of interleukinâ€6 in monitoring severe case of coronavirus disease 2019. EMBO Molecular Medicine, 2020, 12, e12421.	6.9	293
4	Chidamide in relapsed or refractory peripheral T cell lymphoma: a multicenter real-world study in China. Journal of Hematology and Oncology, 2017, 10, 69.	17.0	155
5	A Single-Arm, Multicenter, Phase II Study of Camrelizumab in Relapsed or Refractory Classical Hodgkin Lymphoma. Clinical Cancer Research, 2019, 25, 7363-7369.	7.0	102
6	A lncRNA prognostic signature associated with immune infiltration and tumour mutation burden in breast cancer. Journal of Cellular and Molecular Medicine, 2020, 24, 12444-12456.	3 . 6	85
7	Disulfiram inhibits TGF-Î ² -induced epithelial-mesenchymal transition and stem-like features in breast cancer via ERK/NF-Î ² B/Snail pathway. Oncotarget, 2015, 6, 40907-40919.	1.8	75
8	lncRNA OSTN-AS1 May Represent a Novel Immune-Related Prognostic Marker for Triple-Negative Breast Cancer Based on Integrated Analysis of a ceRNA Network. Frontiers in Genetics, 2019, 10, 850.	2.3	42
9	Efficacy and safety of geptanolimab (GB226) for relapsed or refractory peripheral T cell lymphoma: an open-label phase 2 study (Gxplore-002). Journal of Hematology and Oncology, 2021, 14, 12.	17.0	40
10	Disulfiram/copper targets stem cellâ€like ALDH ⁺ population of multiple myeloma by inhibition of ALDH1A1 and Hedgehog pathway. Journal of Cellular Biochemistry, 2018, 119, 6882-6893.	2.6	39
11	Identification of Hub Genes and Key Pathways Associated with Two Subtypes of Diffuse Large B-Cell Lymphoma Based on Gene Expression Profiling via Integrated Bioinformatics. BioMed Research International, 2018, 2018, 1-14.	1.9	31
12	Interim PET/CT based on visual and semiquantitative analysis predicts survival in patients with diffuse large Bâ€cell lymphoma. Cancer Medicine, 2019, 8, 5012-5022.	2.8	24
13	Epidemiologic characteristics of malignant lymphoma in Hubei, China. Medicine (United States), 2018, 97, e12120.	1.0	22
14	Efficacy and Safety of a Pegasparaginase-Based Chemotherapy Regimen vs an L-asparaginase–Based Chemotherapy Regimen for Newly Diagnosed Advanced Extranodal Natural Killer/T-Cell Lymphoma. JAMA Oncology, 2022, 8, 1035.	7.1	21
15	RhoA/ROCK pathway inhibition by fasudil suppresses the vasculogenic mimicry of U2OS osteosarcoma cells in vitro. Anti-Cancer Drugs, 2017, 28, 514-521.	1.4	20
16	Identification of an Immune-Related Prognostic Signature Associated With Immune Infiltration in Melanoma. Frontiers in Genetics, 2020, 11 , 1002 .	2.3	16
17	Chidamide plus prednisone, etoposide, and thalidomide for untreated angioimmunoblastic Tâ€cell lymphoma in a Chinese population: A multicenter phase ⟨scp⟩ll⟨/scp⟩ trial. American Journal of Hematology, 2022, 97, 623-629.	4.1	15
18	Identification of hub genes and key pathways associated with angioimmunoblastic T-cell lymphoma using weighted gene co-expression network analysis. Cancer Management and Research, 2019, Volume 11, 5209-5220.	1.9	14

#	Article	IF	CITATIONS
19	Disulfiram Improves the Anti-PD-1 Therapy Efficacy by Regulating PD-L1 Expression via Epigenetically Reactivation of IRF7 in Triple Negative Breast Cancer. Frontiers in Oncology, 2021, 11, 734853.	2.8	14
20	A Primary Mediastinal Large B-Cell Lymphoma Patient With COVID-19 Infection After Intensive Immunochemotherapy: A Case Report. Frontiers in Oncology, 2020, 10, 924.	2.8	13
21	Treatment, Survival, and Prognosis of Advanced-Stage Natural Killer/T-Cell Lymphoma: An Analysis From the China Lymphoma Collaborative Group. Frontiers in Oncology, 2020, 10, 583050.	2.8	12
22	A Multi-Center, Real-World Study of Chidamide for Patients With Relapsed or Refractory Peripheral T-Cell Lymphomas in China. Frontiers in Oncology, 2021, 11, 750323.	2.8	12
23	Expression of Nischarin negatively correlates with estrogen receptor and alters apoptosis, migration and invasion in human breast cancer. Biochemical and Biophysical Research Communications, 2017, 484, 536-542.	2.1	11
24	Camrelizumab for relapsed or refractory classical Hodgkin lymphoma: extended followâ€up of the multicenter, singleâ€arm, phase 2 study. International Journal of Cancer, 2021, , .	5.1	11
25	Diagnosis and Treatment of Synchronous Lymphoma and Digestive System Carcinoma: Report of Four Cases and Literature Review. Frontiers in Oncology, 2019, 9, 1367.	2.8	10
26	Pegaspargase, gemcitabine, dexamethasone, and cisplatin (P-GDP) combined chemotherapy is effective for newly diagnosed extranodal NK/T-cell lymphoma: a retrospective study. Cancer Management and Research, 2018, Volume 10, 5061-5069.	1.9	9
27	Detection of circulating tumor cells by RT-PCR significantly associated with poor prognosis in breast cancer. Breast Cancer Research and Treatment, 2011, 130, 359-364.	2.5	8
28	miR-30a/SOX4 Double Negative Feedback Loop is modulated by Disulfiram and regulates EMT and Stem Cell-like properties in Breast Cancer. Journal of Cancer, 2021, 12, 5053-5065.	2.5	8
29	Primitive Neuroectodermal Tumor of the Kidney With Inferior Vena Cava Tumor Thrombus. Journal of the National Medical Association, 2009, 101, 1291-1294.	0.8	6
30	Complete Atelectasis of the Left Lung from Anaplastic Large-Cell Lymphoma. Scientific World Journal, The, 2010, 10, 1332-1333.	2.1	4
31	Long-term outcomes of upfront concurrent chemoradiotherapy followed by P-GDP regimen in newly diagnosed early stage extranodal nasal-type NK/T cell lymphoma. Medicine (United States), 2020, 99, e21705.	1.0	4
32	Serum metabolomic profiling based on GC/MS helped to discriminate Diffuse Large B-cell Lymphoma patients with different prognosis. Leukemia Research, 2021, 111, 106693.	0.8	4
33	Gemcitabine, cisplatin, and dexamethasone and ifosfamide, carboplatin, and etoposide regimens have similar efficacy as salvage treatment for relapsed/refractory aggressive lymphoma. Medicine (United) Tj ETQq1	1 0. 7& 4314	1 rgBT /Overlo
34	GEMSTONE-201: Preplanned primary analysis of a multicenter, single-arm, phase 2 study of sugemalimab (suge) in patients (pts) with relapsed or refractory extranodal natural killer/T cell lymphoma (R/R) Tj ETQq0 0 0	rgBTI/ © verl	ock#10 Tf 50 1
35	Beneficial effect of consolidative radiotherapy for patients with lymphoma and skeletal involvement. Medicine (United States), 2019, 98, e16688.	1.0	3
36	Case Report: Long-Term Response to Radiotherapy Combined With Targeted Therapy in Histiocytic Sarcoma Harboring Mutations in MAPK and PI3K/AKT Pathways. Frontiers in Oncology, 2021, 11, 755893.	2.8	3

#	Article	IF	CITATIONS
37	Ibrutinib as monotherapy versus combination therapy in Chinese patients with relapsed/refractory mantle cell lymphoma: A multicenter study. Cancer Medicine, 2022, 11, 4134-4145.	2.8	3
38	A novel prognostic nomogram for patients with extragastric mucosaâ€associated lymphoid tissue lymphoma: AÂmulticenter study. Cancer Medicine, 2022, 11, 3407-3416.	2.8	1
39	Successful Treatment of Chidamide and Cyclosporine for Refractory/Relapsed Angioimmunoblastic T Cell Lymphoma With Evans Syndrome: A Case Report With Long-Term Follow-Up. Frontiers in Oncology, 2020, 10, 1725.	2.8	O
40	Efficacy and Safety Analysis of Ibrutinib-Containing Therapy for Relapsed/Refractory (R/R) Mantle Cell Lymphoma (MCL): Results from a Real-World Study in China. Blood, 2020, 136, 1-1.	1.4	0
41	Phase 1 dose escalation study of MH048, a novel and non-covalent BTK inhibitor in patients with relapsed or refractory B-cell malignancies Journal of Clinical Oncology, 2022, 40, 7547-7547.	1.6	0