
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2510690/publications.pdf Version: 2024-02-01



ΗΙΡΟΙΙΙΜΑΤΑ

#	Article	IF	CITATIONS
1	Trastuzumab after Adjuvant Chemotherapy in HER2-Positive Breast Cancer. New England Journal of Medicine, 2005, 353, 1659-1672.	27.0	4,601
2	Atezolizumab and Nab-Paclitaxel in Advanced Triple-Negative Breast Cancer. New England Journal of Medicine, 2018, 379, 2108-2121.	27.0	3,097
3	Alpelisib for <i>PIK3CA</i> -Mutated, Hormone Receptor–Positive Advanced Breast Cancer. New England Journal of Medicine, 2019, 380, 1929-1940.	27.0	1,582
4	Fulvestrant plus palbociclib versus fulvestrant plus placebo for treatment of hormone-receptor-positive, HER2-negative metastatic breast cancer that progressed on previous endocrine therapy (PALOMA-3): final analysis of the multicentre, double-blind, phase 3 randomised controlled trial. Lancet Oncology, The, 2016, 17, 425-439.	10.7	1,344
5	Palbociclib in Hormone-Receptor–Positive Advanced Breast Cancer. New England Journal of Medicine, 2015, 373, 209-219.	27.0	1,239
6	Trastuzumab Deruxtecan in Previously Treated HER2-Positive Breast Cancer. New England Journal of Medicine, 2020, 382, 610-621.	27.0	1,143
7	Atezolizumab plus nab-paclitaxel as first-line treatment for unresectable, locally advanced or metastatic triple-negative breast cancer (IMpassion130): updated efficacy results from a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet Oncology, The, 2020, 21, 44-59.	10.7	826
8	Overall Survival with Palbociclib and Fulvestrant in Advanced Breast Cancer. New England Journal of Medicine, 2018, 379, 1926-1936.	27.0	805
9	Trastuzumab-Associated Cardiac Adverse Effects in the Herceptin Adjuvant Trial. Journal of Clinical Oncology, 2007, 25, 3859-3865.	1.6	505
10	Trastuzumab Deruxtecan versus Trastuzumab Emtansine for Breast Cancer. New England Journal of Medicine, 2022, 386, 1143-1154.	27.0	474
11	Antitumor Activity and Safety of Trastuzumab Deruxtecan in Patients With HER2-Low–Expressing Advanced Breast Cancer: Results From a Phase Ib Study. Journal of Clinical Oncology, 2020, 38, 1887-1896.	1.6	465
12	Neratinib after trastuzumab-based adjuvant therapy in HER2-positive breast cancer (ExteNET): 5-year analysis of a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet Oncology, The, 2017, 18, 1688-1700.	10.7	451
13	Buparlisib plus fulvestrant versus placebo plus fulvestrant in postmenopausal, hormone receptor-positive, HER2-negative, advanced breast cancer (BELLE-2): a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet Oncology, The, 2017, 18, 904-916.	10.7	427
14	Trastuzumab deruxtecan (DS-8201a) in patients with advanced HER2-positive breast cancer previously treated with trastuzumab emtansine: a dose-expansion, phase 1 study. Lancet Oncology, The, 2019, 20, 816-826.	10.7	252
15	Targeting HER2 with Trastuzumab Deruxtecan: A Dose-Expansion, Phase I Study in Multiple Advanced Solid Tumors. Cancer Discovery, 2020, 10, 688-701.	9.4	212
16	Palbociclib in Combination With Fulvestrant in Women With Hormone Receptor-Positive/HER2-Negative Advanced Metastatic Breast Cancer: Detailed Safety Analysis From a Multicenter, Randomized, Placebo-Controlled, Phase III Study (PALOMA-3). Oncologist, 2016, 21, 1165-1175.	3.7	183
17	Atezolizumab and <i>nab</i> -Paclitaxel in Advanced Triple-Negative Breast Cancer: Biomarker Evaluation of the IMpassion130 Study. Journal of the National Cancer Institute, 2021, 113, 1005-1016.	6.3	171
18	Trastuzumab deruxtecan (DS-8201a) in patients with advanced HER2-positive gastric cancer: a dose-expansion, phase 1 study. Lancet Oncology, The, 2019, 20, 827-836.	10.7	154

#	Article	IF	CITATIONS
19	KEYNOTE-355: Randomized, double-blind, phase III study of pembrolizumab + chemotherapy versus placebo + chemotherapy for previously untreated locally recurrent inoperable or metastatic triple-negative breast cancer Journal of Clinical Oncology, 2020, 38, 1000-1000.	1.6	135
20	Multi-center study evaluating circulating tumor cells as a surrogate for response to treatment and overall survival in metastatic breast cancer. Breast Cancer, 2010, 17, 199-204.	2.9	123
21	Phase III, Randomized Study of Dual Human Epidermal Growth Factor Receptor 2 (HER2) Blockade With Lapatinib Plus Trastuzumab in Combination With an Aromatase Inhibitor in Postmenopausal Women With HER2-Positive, Hormone Receptor–Positive Metastatic Breast Cancer: ALTERNATIVE. Journal of Clinical Oncology. 2018. 36. 741-748.	1.6	110
22	Phase II study of preoperative sequential FEC and docetaxel predicts of pathological response and disease free survival. Breast Cancer Research and Treatment, 2008, 110, 531-539.	2.5	97
23	PALOMA-3: Phase III Trial of Fulvestrant With or Without Palbociclib in Premenopausal and Postmenopausal Women With Hormone Receptor–Positive, Human Epidermal Growth Factor Receptor 2–Negative Metastatic Breast Cancer That Progressed on Prior Endocrine Therapy—Safety and Efficacy in Asian Patients. Iournal of Global Oncology. 2017. 3. 289-303.	0.5	94
24	The frequency of low HER2 expression in breast cancer and a comparison of prognosis between patients with HER2-low and HER2-negative breast cancer by HR status. Breast Cancer, 2022, 29, 234-241.	2.9	90
25	PD-L1 Immunohistochemistry Assay Comparison in Atezolizumab Plus <i>nab</i> -Paclitaxel–Treated Advanced Triple-Negative Breast Cancer. Journal of the National Cancer Institute, 2021, 113, 1733-1743.	6.3	83
26	Comparison of clinical outcomes between luminal invasive ductal carcinoma and luminal invasive lobular carcinoma. BMC Cancer, 2016, 16, 248.	2.6	78
27	Validation of the 21-gene test as a predictor of clinical response to neoadjuvant hormonal therapy for ER+, HER2-negative breast cancer: the TransNEOS study. Breast Cancer Research and Treatment, 2019, 173, 123-133.	2.5	77
28	Overall Survival with Palbociclib and Fulvestrant in Women with HR+/HER2â^' ABC: Updated Exploratory Analyses of PALOMA-3, a Double-blind, Phase III Randomized Study. Clinical Cancer Research, 2022, 28, 3433-3442.	7.0	65
29	Efficacy of everolimus with exemestane versus exemestane alone in Asian patients with HER2-negative, hormone-receptor-positive breast cancer in BOLERO-2. Breast Cancer, 2014, 21, 703-714.	2.9	57
30	Evaluating the 21-gene assay Recurrence Score® as a predictor of clinical response to 24Âweeks of neoadjuvant exemestane in estrogen receptor-positive breast cancer. International Journal of Clinical Oncology, 2014, 19, 607-613.	2.2	54
31	Docetaxel Followed by Fluorouracil/Epirubicin/Cyclophosphamide as Neoadjuvant Chemotherapy for Patients with Primary Breast Cancer. Japanese Journal of Clinical Oncology, 2011, 41, 867-875.	1.3	53
32	Randomized Controlled Trial of Trastuzumab With or Without Chemotherapy for HER2-Positive Early Breast Cancer in Older Patients. Journal of Clinical Oncology, 2020, 38, 3743-3752.	1.6	50
33	The Japanese Breast Cancer Society Clinical Practice Guidelines for systemic treatment of breast cancer, 2018 edition. Breast Cancer, 2020, 27, 322-331.	2.9	47
34	Ki67 index changes, pathological response and clinical benefits in primary breast cancer patients treated with 24 weeks of aromatase inhibition. Cancer Science, 2011, 102, 858-865.	3.9	44
35	Palbociclib in combination with letrozole in patients with estrogen receptor–positive, human epidermal growth factor receptor 2–negative advanced breast cancer: PALOMA-2 subgroup analysis of Japanese patients. International Journal of Clinical Oncology, 2019, 24, 274-287.	2.2	43
36	Palbociclib in combination with fulvestrant in patients with hormone receptor-positive, human epidermal growth factor receptor 2-negative advanced breast cancer: PALOMA-3 subgroup analysis of Japanese patients. International Journal of Clinical Oncology, 2019, 24, 262-273.	2.2	39

#	Article	IF	CITATIONS
37	Evaluation of Trastuzumab Without Chemotherapy as a Post-operative Adjuvant Therapy in HER2-positive Elderly Breast Cancer Patients: Randomized Controlled Trial [RESPECT (N-SAS BC07)]. Japanese Journal of Clinical Oncology, 2011, 41, 709-712.	1.3	38
38	Prognostic factors of HER2-positive breast cancer patients who develop brain metastasis: a multicenter retrospective analysis. Breast Cancer Research and Treatment, 2015, 149, 277-284.	2.5	32
39	Prognostic significance of subtype and pathologic response in operable breast cancer; a pooled analysis of prospective neoadjuvant studies of JBCRG. Breast Cancer, 2015, 22, 486-495.	2.9	29
40	Palbociclib in combination with letrozole as firstâ€line treatment for advanced breast cancer: A Japanese phase <scp>II</scp> study. Cancer Science, 2018, 109, 803-813.	3.9	29
41	Clinical development of CDK4/6 inhibitor for breast cancer. Breast Cancer, 2018, 25, 402-406.	2.9	29
42	Future treatment strategies for metastatic breast cancer: curable or incurable?. Breast Cancer, 2012, 19, 200-205.	2.9	26
43	Brain Metastases in Breast Cancer. Japanese Journal of Clinical Oncology, 2014, 44, 1133-1140.	1.3	26
44	Treatment Exposure and Discontinuation in the PALbociclib CoLlaborative Adjuvant Study of Palbociclib With Adjuvant Endocrine Therapy for Hormone Receptor–Positive/Human Epidermal Growth Factor Receptor 2–Negative Early Breast Cancer (PALLAS/AFT-05/ABCSG-42/BIG-14-03). Journal of Clinical Oncology, 2022, 40, 449-458.	1.6	25
45	Patient-Reported Outcomes in Patients With <i>PIK3CA</i> -Mutated Hormone Receptor–Positive, Human Epidermal Growth Factor Receptor 2–Negative Advanced Breast Cancer From SOLAR-1. Journal of Clinical Oncology, 2021, 39, 2005-2015.	1.6	23
46	Advances in treatment and care in metastatic breast cancer (MBC): are there MBC patients who are curable?. Chinese Clinical Oncology, 2018, 7, 23-23.	1.2	22
47	A phase II study of lapatinib for brain metastases in patients with HER2-overexpressing breast cancer following trastuzumab based systemic therapy and cranial radiotherapy: subset analysis of Japanese patients. International Journal of Clinical Oncology, 2013, 18, 621-628.	2.2	19
48	Comparison of different definitions of pathologic complete response in operable breast cancer: a pooled analysis of three prospective neoadjuvant studies of JBCRG. Breast Cancer, 2015, 22, 586-595.	2.9	19
49	Sentinel lymph node biopsy is not necessary in patients diagnosed with ductal carcinoma in situ of the breast by stereotactic vacuum-assisted biopsy. Breast Cancer, 2016, 23, 190-194.	2.9	18
50	Patient-reported outcomes and objective assessments with arm measurement and bioimpedance analysis for lymphedema among breast cancer survivors. Breast Cancer Research and Treatment, 2020, 179, 91-100.	2.5	16
51	Health-Related Quality of Life With Trastuzumab Monotherapy Versus Trastuzumab Plus Standard Chemotherapy as Adjuvant Therapy in Older Patients With HER2-Positive Breast Cancer. Journal of Clinical Oncology, 2021, 39, 2452-2462.	1.6	16
52	Evaluation of the Association of Polymorphisms With Palbociclib-Induced Neutropenia: Pharmacogenetic Analysis of PALOMA-2/-3. Oncologist, 2021, 26, e1143-e1155.	3.7	15
53	The Japanese Breast Cancer Society Clinical Practice Guidelines, 2018 edition: the tool for shared decision making between doctor and patient. Breast Cancer, 2020, 27, 1-3.	2.9	14
54	Circulating tumor cells as a prognostic marker for efficacy in the randomized phase III JO21095 trial in Japanese patients with HER2-negative metastatic breast cancer. Breast Cancer Research and Treatment, 2017, 162, 501-510.	2.5	13

#	Article	IF	CITATIONS
55	Comparison of sentinel lymph node biopsy between invasive lobular carcinoma and invasive ductal carcinoma. Breast Cancer, 2018, 25, 560-565.	2.9	13
56	Neoadjuvant endocrine therapy for postmenopausal patients with hormone receptor-positive early breast cancer: a new concept. Breast Cancer, 2011, 18, 92-97.	2.9	10
57	Current Status of Advance Care Planning and Endâ€ofâ€life Communication for Patients with Advanced and Metastatic Breast Cancer. Oncologist, 2021, 26, e686-e693.	3.7	10
58	Neutropenia management with palbociclib in Japanese patients with advanced breast cancer. Breast Cancer, 2019, 26, 637-650.	2.9	8
59	Compression therapy using surgical gloves does not prevent paclitaxel-induced peripheral neuropathy: results from a double-blind phase 2 trial. BMC Cancer, 2021, 21, 548.	2.6	8
60	Perspective of trastuzumab treatment. Breast Cancer, 2007, 14, 150-155.	2.9	7
61	Outcomes of trastuzumab therapy in HER2-positive early breast cancer patients. International Journal of Clinical Oncology, 2015, 20, 709-722.	2.2	7
62	Efficacy, safety, pharmacokinetics and biomarker findings in patients with HER2-positive advanced or metastatic breast cancer treated with lapatinib in combination with capecitabine: results from 51 Japanese patients treated in a clinical study. Breast Cancer, 2015, 22, 192-200.	2.9	7
63	Safety and pharmacokinetics of ramucirumab in combination with docetaxel in Japanese patients with locally advanced or metastatic breast cancer: a Phase Ib study. Japanese Journal of Clinical Oncology, 2016, 46, 1088-1094.	1.3	7
64	Efficacy and safety of low-dose capecitabine plus docetaxel versus single-agent docetaxel in patients with anthracycline-pretreated HER2-negative metastatic breast cancer: results from the randomized phase III JO21095 trial. Breast Cancer Research and Treatment, 2017, 161, 473-482.	2.5	7
65	Outcomes of trastuzumab therapy in HER2-positive early breast cancer patients: extended follow-up of JBCRG-cohort study 01. Breast Cancer, 2020, 27, 631-641.	2.9	6
66	Effects of ABCB1 and ABCG2 polymorphisms on the pharmacokinetics of abemaciclib. European Journal of Clinical Pharmacology, 2022, 78, 1239-1247.	1.9	6
67	Effects of neoadjuvant chemotherapy on operative adverse events and chemotherapy and radiotherapy in patients undergoing immediate breast reconstruction. Breast Cancer, 2020, 27, 716-723.	2.9	5
68	Phase I dose-finding study of eribulin and capecitabine for metastatic breast cancer: JBCRG-18 cape study. Breast Cancer, 2018, 25, 108-117.	2.9	4
69	Prediction of pathological margin status using preoperative contrast-enhanced MRI in patients with early breast cancer who underwent skin-sparing mastectomy. Breast Journal, 2019, 25, 202-206.	1.0	4
70	Study protocol for a nationwide questionnaire survey of physical activity among breast cancer survivors in Japan. BMJ Open, 2020, 10, e032871.	1.9	4
71	Palbociclib as an early-line treatment for Japanese patients with hormone receptor–positive/human epidermal growth factor receptor 2–negative advanced breast cancer: a review of clinical trial and real-world data. International Journal of Clinical Oncology, 2021, 26, 2179-2193.	2.2	4
72	Rainbow of KIBOU (ROK) study: a Breast Cancer Survivor Cohort in Japan. Breast Cancer, 2018, 25, 60-67.	2.9	3

#	Article	IF	CITATIONS
73	Management of breast cancer in older patients. Japanese Journal of Clinical Oncology, 2022, 52, 682-689.	1.3	3
74	Clinical Utility of Precision Medicine in Early Breast Cancer: What Is the Optimal Framework to Develop Precision Medicine?. Journal of Clinical Oncology, 2022, 40, 1962-1963.	1.6	3
75	Neo(adjuvant) trastuzumab treatment: current perspectives. Breast Cancer, 2009, 16, 288-294.	2.9	2
76	The Transition of Breast Cancer Treatment and Japan Clinical Oncology Group Research Over Two Decades. Japanese Journal of Clinical Oncology, 2012, 42, 14-20.	1.3	2
77	Impact of adjuvant endocrine therapy on prognosis in small hormone receptor-positive, HER2-negative early breast cancer. Breast Cancer, 2021, 28, 1087-1095.	2.9	2
78	Oncology care providers' awareness and practice related to physical activity promotion for breast cancer survivors and barriers and facilitators to such promotion: a nationwide cross-sectional web-based survey. Supportive Care in Cancer, 2022, 30, 3105-3118.	2.2	2
79	A single-arm, phase 2 study of steroid-containing mouthwash for the prevention of everolimus-associated stomatitis in multiple tumor types. International Journal of Clinical Oncology, 2019, 24, 1320-1327.	2.2	1
80	Reply to T.J.A. Dekker. Journal of Clinical Oncology, 2020, 38, 3351-3352.	1.6	0
81	Cost-Effectiveness of Trastuzumab With or Without Chemotherapy as Adjuvant Therapy in HER2-Positive Elderly Breast Cancer Patients: A Randomized, Open-Label Clinical Trial, the RESPECT Trial. Clinical Drug Investigation, 2022, 42, 253-262.	2.2	0