## Xiaohua Zhang

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2532607/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	<i>BRAF</i> and <i>TERT</i> promoter mutations in the aggressiveness of papillary thyroid carcinoma: a study of 653 patients. Oncotarget, 2016, 7, 18346-18355.	1.8	109
2	Expression profile analysis of long noncoding RNA in HER-2-enriched subtype breast cancer by next-generation sequencing and bioinformatics. OncoTargets and Therapy, 2016, 9, 761.	2.0	79
3	Osthole inhibits triple negative breast cancer cells by suppressing STAT3. Journal of Experimental and Clinical Cancer Research, 2018, 37, 322.	8.6	50
4	miR-27a regulates the sensitivity of breast cancer cells to cisplatin treatment via BAK-SMAC/DIABLO-XIAP axis. Tumor Biology, 2016, 37, 6837-6845.	1.8	47
5	Curcuminoid EF24 enhances the antiâ€ŧumour activity of Akt inhibitor MKâ€2206 through ROSâ€mediated endoplasmic reticulum stress and mitochondrial dysfunction in gastric cancer. British Journal of Pharmacology, 2017, 174, 1131-1146.	5.4	42
6	MAL2 promotes proliferation, migration, and invasion through regulating epithelial-mesenchymal transition in breast cancer cell lines. Biochemical and Biophysical Research Communications, 2018, 504, 434-439.	2.1	34
7	METTL7B promotes migration and invasion in thyroid cancer through epithelial-mesenchymal transition. Journal of Molecular Endocrinology, 2019, 63, 51-61.	2.5	34
8	<em>ITGA7 </em> functions as a tumor suppressor and regulates migration and invasion in breast cancer. Cancer Management and Research, 2018, Volume 10, 969-976.	1.9	32
9	Schisandrin B exhibits potent anticancer activity in triple negative breast cancer by inhibiting STAT3. Toxicology and Applied Pharmacology, 2018, 358, 110-119.	2.8	31
10	Alantolactone promotes ER stressâ€mediated apoptosis by inhibition of TrxR1 in tripleâ€negative breast cancer cell lines and in a mouse model. Journal of Cellular and Molecular Medicine, 2019, 23, 2194-2206.	3.6	28
11	(S)-crizotinib induces apoptosis in human non-small cell lung cancer cells by activating ROS independent of MTH1. Journal of Experimental and Clinical Cancer Research, 2017, 36, 120.	8.6	27
12	Preoperative endoscopic localization of colorectal cancer and tracing lymph nodes by using carbon nanoparticles in laparoscopy. World Journal of Surgical Oncology, 2016, 14, 231.	1.9	25
13	GABRB2 plays an important role in the lymph node metastasis of papillary thyroid cancer. Biochemical and Biophysical Research Communications, 2017, 492, 323-330.	2.1	21
14	KMT2A histone methyltransferase contributes to colorectal cancer development by promoting cathepsin Z transcriptional activation. Cancer Medicine, 2019, 8, 3544-3552.	2.8	21
15	LRP4 promotes proliferation, migration, and invasion in papillary thyroid cancer. Biochemical and Biophysical Research Communications, 2018, 503, 257-263.	2.1	20
16	H/ACA box small nucleolar RNA 7B acts as an oncogene and a potential prognostic biomarker in breast cancer. Cancer Cell International, 2019, 19, 125.	4.1	20
17	TEKT4 Promotes Papillary Thyroid Cancer Cell Proliferation, Colony Formation, and Metastasis through Activating PI3K/Akt Pathway. Endocrine Pathology, 2018, 29, 310-316.	9.0	18
18	A mono-carbonyl analog of curcumin induces apoptosis in drug-resistant EGFR-mutant lung cancer through the generation of oxidative stress and mitochondrial dysfunction. Cancer Management and Research, 2018, Volume 10, 3069-3082.	1.9	18

XIAOHUA ZHANG

#	Article	IF	CITATIONS
19	Synaptopodin-2 plays an important role in the metastasis of breast cancer via PI3K/Akt/mTOR pathway. Cancer Management and Research, 2018, Volume 10, 1575-1583.	1.9	17
20	Upregulation of LAGE3 correlates with prognosis and immune infiltrates in colorectal cancer: A bioinformatic analysis. International Immunopharmacology, 2020, 85, 106599.	3.8	17
21	Prediction of central lymph node metastasis in 392 patients with cervical lymph node-negative papillary thyroid carcinoma in Eastern China. Oncology Letters, 2015, 10, 2559-2564.	1.8	16
22	The Utility of Sentinel Lymph Node Biopsy in Papillary Thyroid Carcinoma with Occult Lymph Nodes. PLoS ONE, 2015, 10, e0129304.	2.5	16
23	ZCCHC12, a novel oncogene in papillary thyroid cancer. Journal of Cancer Research and Clinical Oncology, 2017, 143, 1679-1686.	2.5	15
24	Uridine phosphorylase 1 associates to biological and clinical significance in thyroid carcinoma cell lines. Journal of Cellular and Molecular Medicine, 2019, 23, 7438-7448.	3.6	15
25	Silencing of semaphorin 3C suppresses cell proliferation and migration in MCFâ€7 breast cancer cells. Oncology Letters, 2017, 14, 5913-5917.	1.8	14
26	Original tumour suppressor gene polycystic kidney and hepatic disease 1‑like 1 is associated with thyroid cancer cell progression. Oncology Letters, 2019, 18, 3227-3235.	1.8	13
27	The use of OK-432 to prevent seroma in extended latissimus dorsi flap donor site after breast reconstruction. Journal of Surgical Research, 2015, 193, 492-496.	1.6	12
28	<p>Lipase member H is a downstream molecular target of hypoxia inducible factor-1α and promotes papillary thyroid carcinoma cell migration in BCPAP and KTC-1 cell lines</p> . Cancer Management and Research, 2019, Volume 11, 931-941.	1.9	10
29	Downregulating integrin subunit alpha 7 (ITGA7) promotes proliferation, invasion, and migration of papillary thyroid carcinoma cells through regulating epithelial-to-mesenchymal transition. Acta Biochimica Et Biophysica Sinica, 2020, 52, 116-124.	2.0	10
30	Exploration of the Prognostic and Immunotherapeutic Value of B and T Lymphocyte Attenuator in Skin Cutaneous Melanoma. Frontiers in Oncology, 2020, 10, 592811.	2.8	10
31	Identification of a Pyroptosis-Related Prognostic Signature Combined With Experiments in Hepatocellular Carcinoma. Frontiers in Molecular Biosciences, 2022, 9, 822503.	3.5	10
32	Identification and validation of L Antigen Family Member 3 as an immune-related biomarker associated with the progression of papillary thyroid cancer. International Immunopharmacology, 2021, 90, 107267.	3.8	9
33	Clinicopathological characteristics and prognostic factors for primary thyroid lymphoma: report on 28 Chinese patients and results of a population-based study. Cancer Management and Research, 2018, Volume 10, 4411-4419.	1.9	8
34	Genomic Instability-Related LncRNA Signature Predicts the Prognosis and Highlights LINC01614 Is a Tumor Microenvironment-Related Oncogenic IncRNA of Papillary Thyroid Carcinoma. Frontiers in Oncology, 2021, 11, 737867.	2.8	8
35	CUX2 functions as an oncogene in papillary thyroid cancer. OncoTargets and Therapy, 2019, Volume 12, 217-224.	2.0	7
36	<clinical a="" advanced<br="" apatinib="" as="" chemo-refractory="" in="" of="" patients="" salvage="" treatment="" value="" with="">Cervical Cancer. OncoTargets and Therapy, 2019, Volume 12, 9707-9713.</clinical>	2.0	7

XIAOHUA ZHANG

#	Article	IF	CITATIONS
37	LEM domain containing 1 promotes thyroid cancer cell proliferation and migration by activating the Wnt/βâ€ʿcatenin signaling pathway and epithelialâ€ʿmesenchymal transition. Oncology Letters, 2021, 21, 442.	1.8	7
38	Identification of the prognostic and immunotherapeutic potential of L antigen family member 3 in malignant pleural mesothelioma. Clinical and Translational Medicine, 2020, 10, e207.	4.0	5
39	Major Vault Protein (MVP) Associated With BRAFV600E Mutation Is an Immune Microenvironment-Related Biomarker Promoting the Progression of Papillary Thyroid Cancer via MAPK/ERK and PI3K/AKT Pathways. Frontiers in Cell and Developmental Biology, 2021, 9, 688370.	3.7	5
40	T-Box Transcription Factor 22 Is an Immune Microenvironment-Related Biomarker Associated With the BRAFV600E Mutation in Papillary Thyroid Carcinoma. Frontiers in Cell and Developmental Biology, 2020, 8, 590898.	3.7	4
41	Papillary thyroid microcarcinoma with synchronous asymptomatic advanced esophageal squamous cell carcinoma: A case report and review of the literature. Oncology Letters, 2015, 9, 731-734.	1.8	3
42	β3-tubulin is a good predictor of sensitivity to taxane-based neoadjuvant chemotherapy in primary breast cancer. Clinical and Experimental Medicine, 2016, 16, 391-397.	3.6	3
43	Up-regulation of L Antigen Family Member 3 Associates With Aggressive Progression of Breast Cancer. Frontiers in Oncology, 2020, 10, 553628.	2.8	3
44	Treatment patterns for adjuvant docetaxel-based chemotherapy in early-stage breast cancer in China: A pooled retrospective analysis of four observational studies. Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association, Beijing Institute for Cancer Research, 2018, 30, 327-339.	2.2	3
45	Glucose-to-Lymphocyte Ratio (GLR) as a Predictor of Preoperative Central Lymph Node Metastasis in Papillary Thyroid Cancer Patients With Type 2 Diabetes Mellitus and Construction of the Nomogram. Frontiers in Endocrinology, 2022, 13, 829009.	3.5	3
46	Downstream Neighbor of Son Overexpression is Associated With Breast Cancer Progression and a Poor Prognosis. Journal of Breast Cancer, 2022, 25, 327.	1.9	3
47	The Prognostic Value of Combination of Plasma Fibrinogen and CA19-9 in Non-Distant Metastatic Breast Cancer Patients Undergoing Surgery. Cancer Management and Research, 2020, Volume 12, 8875-8886.	1.9	2
48	Subcutaneous Recurrences of Thyroid Cancer After Conventional Transcervical Thyroidectomy: A Case Report. Frontiers in Surgery, 2020, 7, 586106.	1.4	2
49	The ETNK2 gene promotes progression of papillary thyroid carcinoma through the HIPPO pathway. Journal of Cancer, 2022, 13, 508-516.	2.5	1
50	Development and Validation of a Nomogram for Preoperative Prediction of Central Compartment Lymph Node Metastasis in Patients with Papillary Thyroid Carcinoma and Type 2 Diabetes Mellitus. Cancer Management and Research, 2021, Volume 13, 2499-2513.	1.9	0
51	Treatment patterns and patient profiles for docetaxel-based adjuvant chemotherapy in early-stage breast cancer in China: A pooled analysis of four observational studies Journal of Clinical Oncology, 2017, 35, e12017-e12017.	1.6	0