

Xiaohua Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2532607/publications.pdf>

Version: 2024-02-01

51
papers

914
citations

471509

17
h-index

501196

28
g-index

51
all docs

51
docs citations

51
times ranked

1706
citing authors

#	ARTICLE	IF	CITATIONS
1	The ETNK2 gene promotes progression of papillary thyroid carcinoma through the HIPPO pathway. <i>Journal of Cancer</i> , 2022, 13, 508-516.	2.5	1
2	Identification of a Pyroptosis-Related Prognostic Signature Combined With Experiments in Hepatocellular Carcinoma. <i>Frontiers in Molecular Biosciences</i> , 2022, 9, 822503.	3.5	10
3	Glucose-to-Lymphocyte Ratio (GLR) as a Predictor of Preoperative Central Lymph Node Metastasis in Papillary Thyroid Cancer Patients With Type 2 Diabetes Mellitus and Construction of the Nomogram. <i>Frontiers in Endocrinology</i> , 2022, 13, 829009.	3.5	3
4	Downstream Neighbor of Son Overexpression is Associated With Breast Cancer Progression and a Poor Prognosis. <i>Journal of Breast Cancer</i> , 2022, 25, 327.	1.9	3
5	Identification and validation of L Antigen Family Member 3 as an immune-related biomarker associated with the progression of papillary thyroid cancer. <i>International Immunopharmacology</i> , 2021, 90, 107267.	3.8	9
6	Development and Validation of a Nomogram for Preoperative Prediction of Central Compartment Lymph Node Metastasis in Patients with Papillary Thyroid Carcinoma and Type 2 Diabetes Mellitus. <i>Cancer Management and Research</i> , 2021, Volume 13, 2499-2513.	1.9	0
7	LEM domain containing 1 promotes thyroid cancer cell proliferation and migration by activating the Wnt/ β -catenin signaling pathway and epithelial-mesenchymal transition. <i>Oncology Letters</i> , 2021, 21, 442.	1.8	7
8	Genomic Instability-Related LncRNA Signature Predicts the Prognosis and Highlights LINC01614 Is a Tumor Microenvironment-Related Oncogenic lncRNA of Papillary Thyroid Carcinoma. <i>Frontiers in Oncology</i> , 2021, 11, 737867.	2.8	8
9	Major Vault Protein (MVP) Associated With BRAFV600E Mutation Is an Immune Microenvironment-Related Biomarker Promoting the Progression of Papillary Thyroid Cancer via MAPK/ERK and PI3K/AKT Pathways. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 688370.	3.7	5
10	<p>The Prognostic Value of Combination of Plasma Fibrinogen and CA19-9 in Non-Distant Metastatic Breast Cancer Patients Undergoing Surgery</p>. <i>Cancer Management and Research</i> , 2020, Volume 12, 8875-8886.	1.9	2
11	Identification of the prognostic and immunotherapeutic potential of L antigen family member 3 in malignant pleural mesothelioma. <i>Clinical and Translational Medicine</i> , 2020, 10, e207.	4.0	5
12	Subcutaneous Recurrences of Thyroid Cancer After Conventional Transcervical Thyroidectomy: A Case Report. <i>Frontiers in Surgery</i> , 2020, 7, 586106.	1.4	2
13	Downregulating integrin subunit alpha 7 (ITGA7) promotes proliferation, invasion, and migration of papillary thyroid carcinoma cells through regulating epithelial-to-mesenchymal transition. <i>Acta Biochimica Et Biophysica Sinica</i> , 2020, 52, 116-124.	2.0	10
14	Exploration of the Prognostic and Immunotherapeutic Value of B and T Lymphocyte Attenuator in Skin Cutaneous Melanoma. <i>Frontiers in Oncology</i> , 2020, 10, 592811.	2.8	10
15	Up-regulation of L Antigen Family Member 3 Associates With Aggressive Progression of Breast Cancer. <i>Frontiers in Oncology</i> , 2020, 10, 553628.	2.8	3
16	Upregulation of LAGE3 correlates with prognosis and immune infiltrates in colorectal cancer: A bioinformatic analysis. <i>International Immunopharmacology</i> , 2020, 85, 106599.	3.8	17
17	T-Box Transcription Factor 22 Is an Immune Microenvironment-Related Biomarker Associated With the BRAFV600E Mutation in Papillary Thyroid Carcinoma. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 590898.	3.7	4
18	CUX2 functions as an oncogene in papillary thyroid cancer. <i>OncoTargets and Therapy</i> , 2019, Volume 12, 217-224.	2.0	7

#	ARTICLE	IF	CITATIONS
19	KMT2A histone methyltransferase contributes to colorectal cancer development by promoting cathepsin Z transcriptional activation. <i>Cancer Medicine</i> , 2019, 8, 3544-3552.	2.8	21
20	Uridine phosphorylase 1 associates to biological and clinical significance in thyroid carcinoma cell lines. <i>Journal of Cellular and Molecular Medicine</i> , 2019, 23, 7438-7448.	3.6	15
21	Original tumour suppressor gene polycystic kidney and hepatic disease 1-like 1 is associated with thyroid cancer cell progression. <i>Oncology Letters</i> , 2019, 18, 3227-3235.	1.8	13
22	H/ACA box small nucleolar RNA 7B acts as an oncogene and a potential prognostic biomarker in breast cancer. <i>Cancer Cell International</i> , 2019, 19, 125.	4.1	20
23	Lipase member H is a downstream molecular target of hypoxia inducible factor-1 α and promotes papillary thyroid carcinoma cell migration in BCPAP and KTC-1 cell lines. <i>Cancer Management and Research</i> , 2019, Volume 11, 931-941.	1.9	10
24	Clinical Value Of Apatinib As A Salvage Treatment In Patients With Chemo-Refractory Advanced Cervical Cancer. <i>OncoTargets and Therapy</i> , 2019, Volume 12, 9707-9713.	2.0	7
25	Alantolactone promotes ER stress-mediated apoptosis by inhibition of TrxR1 in triple-negative breast cancer cell lines and in a mouse model. <i>Journal of Cellular and Molecular Medicine</i> , 2019, 23, 2194-2206.	3.6	28
26	METTL7B promotes migration and invasion in thyroid cancer through epithelial-mesenchymal transition. <i>Journal of Molecular Endocrinology</i> , 2019, 63, 51-61.	2.5	34
27	Osthole inhibits triple negative breast cancer cells by suppressing STAT3. <i>Journal of Experimental and Clinical Cancer Research</i> , 2018, 37, 322.	8.6	50
28	TEKT4 Promotes Papillary Thyroid Cancer Cell Proliferation, Colony Formation, and Metastasis through Activating PI3K/Akt Pathway. <i>Endocrine Pathology</i> , 2018, 29, 310-316.	9.0	18
29	Clinicopathological characteristics and prognostic factors for primary thyroid lymphoma: report on 28 Chinese patients and results of a population-based study. <i>Cancer Management and Research</i> , 2018, Volume 10, 4411-4419.	1.9	8
30	A mono-carbonyl analog of curcumin induces apoptosis in drug-resistant EGFR-mutant lung cancer through the generation of oxidative stress and mitochondrial dysfunction. <i>Cancer Management and Research</i> , 2018, Volume 10, 3069-3082.	1.9	18
31	Schisandrin B exhibits potent anticancer activity in triple negative breast cancer by inhibiting STAT3. <i>Toxicology and Applied Pharmacology</i> , 2018, 358, 110-119.	2.8	31
32	MAL2 promotes proliferation, migration, and invasion through regulating epithelial-mesenchymal transition in breast cancer cell lines. <i>Biochemical and Biophysical Research Communications</i> , 2018, 504, 434-439.	2.1	34
33	ITGA7 functions as a tumor suppressor and regulates migration and invasion in breast cancer. <i>Cancer Management and Research</i> , 2018, Volume 10, 969-976.	1.9	32
34	Synaptopodin-2 plays an important role in the metastasis of breast cancer via PI3K/Akt/mTOR pathway. <i>Cancer Management and Research</i> , 2018, Volume 10, 1575-1583.	1.9	17
35	LRP4 promotes proliferation, migration, and invasion in papillary thyroid cancer. <i>Biochemical and Biophysical Research Communications</i> , 2018, 503, 257-263.	2.1	20
36	Treatment patterns for adjuvant docetaxel-based chemotherapy in early-stage breast cancer in China: A pooled retrospective analysis of four observational studies. <i>Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association</i> , Beijing Institute for Cancer Research, 2018, 30, 327-339.	2.2	3

#	ARTICLE	IF	CITATIONS
37	Curcuminoid EF24 enhances the anti-tumour activity of Akt inhibitor MK2206 through ROS-mediated endoplasmic reticulum stress and mitochondrial dysfunction in gastric cancer. <i>British Journal of Pharmacology</i> , 2017, 174, 1131-1146.	5.4	42
38	ZCCHC12, a novel oncogene in papillary thyroid cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 1679-1686.	2.5	15
39	GABRB2 plays an important role in the lymph node metastasis of papillary thyroid cancer. <i>Biochemical and Biophysical Research Communications</i> , 2017, 492, 323-330.	2.1	21
40	Silencing of semaphorin 3C suppresses cell proliferation and migration in MCF7 breast cancer cells. <i>Oncology Letters</i> , 2017, 14, 5913-5917.	1.8	14
41	(S)-crizotinib induces apoptosis in human non-small cell lung cancer cells by activating ROS independent of MTH1. <i>Journal of Experimental and Clinical Cancer Research</i> , 2017, 36, 120.	8.6	27
42	Treatment patterns and patient profiles for docetaxel-based adjuvant chemotherapy in early-stage breast cancer in China: A pooled analysis of four observational studies. <i>Journal of Clinical Oncology</i> , 2017, 35, e12017-e12017.	1.6	0
43	<i>BRAF</i> and <i>TERT</i> promoter mutations in the aggressiveness of papillary thyroid carcinoma: a study of 653 patients. <i>Oncotarget</i> , 2016, 7, 18346-18355.	1.8	109
44	Expression profile analysis of long noncoding RNA in HER-2-enriched subtype breast cancer by next-generation sequencing and bioinformatics. <i>OncoTargets and Therapy</i> , 2016, 9, 761.	2.0	79
45	Preoperative endoscopic localization of colorectal cancer and tracing lymph nodes by using carbon nanoparticles in laparoscopy. <i>World Journal of Surgical Oncology</i> , 2016, 14, 231.	1.9	25
46	miR-27a regulates the sensitivity of breast cancer cells to cisplatin treatment via BAK-SMAC/DIABLO-XIAP axis. <i>Tumor Biology</i> , 2016, 37, 6837-6845.	1.8	47
47	β 3-tubulin is a good predictor of sensitivity to taxane-based neoadjuvant chemotherapy in primary breast cancer. <i>Clinical and Experimental Medicine</i> , 2016, 16, 391-397.	3.6	3
48	Prediction of central lymph node metastasis in 392 patients with cervical lymph node-negative papillary thyroid carcinoma in Eastern China. <i>Oncology Letters</i> , 2015, 10, 2559-2564.	1.8	16
49	Papillary thyroid microcarcinoma with synchronous asymptomatic advanced esophageal squamous cell carcinoma: A case report and review of the literature. <i>Oncology Letters</i> , 2015, 9, 731-734.	1.8	3
50	The use of OK-432 to prevent seroma in extended latissimus dorsi flap donor site after breast reconstruction. <i>Journal of Surgical Research</i> , 2015, 193, 492-496.	1.6	12
51	The Utility of Sentinel Lymph Node Biopsy in Papillary Thyroid Carcinoma with Occult Lymph Nodes. <i>PLoS ONE</i> , 2015, 10, e0129304.	2.5	16