

# Armand Bensussan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2590499/publications.pdf>

Version: 2024-02-01

234  
papers

10,152  
citations

36303

51  
h-index

46799

89  
g-index

268  
all docs

268  
docs citations

268  
times ranked

11516  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Involvement of the CD39/CD73/adenosine pathway in T-cell proliferation and NK cell-mediated antibody-dependent cell cytotoxicity in SÅ©zary syndrome. <i>Blood</i> , 2022, 139, 2712-2716.               | 1.4 | 14        |
| 2  | CCR8 is a new therapeutic target in cutaneous T-cell lymphomas. <i>Blood Advances</i> , 2022, 6, 3507-3512.  | 5.2 | 6         |
| 3  | The soluble form of CD160 acts as a tumor mediator of immune escape in melanoma. <i>Cancer Immunology, Immunotherapy</i> , 2022, 71, 2731-2742.  | 4.2 | 6         |
| 4  | Bi38-3 is a novel CD38/CD3 bispecific T-cell engager with low toxicity for the treatment of multiple myeloma. <i>Haematologica</i> , 2021, 106, 1193-1197.   | 3.5 | 23        |
| 5  | The value of five blood markers in differentiating mycosis fungoides and SÅ©zary syndrome: a validation cohort. <i>British Journal of Dermatology</i> , 2021, 185, 405-411.                              | 1.5 | 7         |
| 6  | Expansion of Circulating CD49b+LAG3+ Type 1 Regulatory T Cells in Human Chronic Graft-Versus-Host Disease. <i>Journal of Investigative Dermatology</i> , 2021, 141, 193-197.e2.                          | 0.7 | 4         |
| 7  | PAK1-Dependent Antitumor Effect of AAC-11â€™ Derived Peptides on SÅ©zary Syndrome Malignant CD4+ T Lymphocytes. <i>Journal of Investigative Dermatology</i> , 2021, 141, 2261-2271.e5.                   | 0.7 | 3         |
| 8  | Secretomic and proteomic analysis of cutaneous T cell lymphoma-associated fibroblasts. <i>European Journal of Cancer</i> , 2021, 156, S5.  | 2.8 | 0         |
| 9  | Exploring the role of the skin microenvironment in cutaneous T-cell lymphoma using single cell RNA-sequencing. <i>European Journal of Cancer</i> , 2021, 156, S3-S4.                                     | 2.8 | 3         |
| 10 | ICOS is widely expressed in cutaneous T-cell lymphoma and its targeting promotes potent killing of malignant cells. <i>European Journal of Cancer</i> , 2021, 156, S23-S24.                              | 2.8 | 1         |
| 11 | Anti-tumor effect of anti-apoptosis clone 11 protein-derived peptides on SÅ©zary syndrome malignant CD4+ T lymphocytes. <i>European Journal of Cancer</i> , 2021, 156, S14.                              | 2.8 | 0         |
| 12 | Quantifying response to various treatments using the revisited blood staging of mycosis fungoides and SÅ©zary syndrome with the KIR3DL2 marker. <i>European Journal of Cancer</i> , 2021, 156, S6-S7.    | 2.8 | 0         |
| 13 | ICOS Is Widely Expressed in Cutaneous T-Cell Lymphoma and Its Targeting Promotes Potent Killing of Malignant Cells. <i>Blood</i> , 2021, 138, 790-790.   | 1.4 | 4         |
| 14 | Chimerized Anti-ICOS 314.8 Monoclonal Antibodies Inhibit Tumor Cells and Regulatory T Cells in Patients with SÅ©zary Syndrome. <i>Blood</i> , 2021, 138, 2260-2260.                                      | 1.4 | 0         |
| 15 | 857â€™...Selective Treg depletion in solid tumors with ALD2510, a novel humanized CD25-specific, IL-2 sparing monoclonal antibody. , 2021, 9, A898-A898.   |     | 3         |
| 16 | Revisiting the initial diagnosis and blood staging of mycosis fungoides and SÅ©zary syndrome with the <sc>KIR</sc> 3 <sc>DL</sc> 2 marker. <i>British Journal of Dermatology</i> , 2020, 182, 1415-1422. | 1.5 | 20        |
| 17 | Persistent deficiency of mucosal-associated invariant T cells during dermatomyositis. <i>Rheumatology</i> , 2020, 59, 2282-2286.   | 1.9 | 19        |
| 18 | ICOS is widely expressed in cutaneous T-cell lymphoma, and its targeting promotes potent killing of malignant cells. <i>Blood Advances</i> , 2020, 4, 5203-5214.   | 5.2 | 18        |

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 19 | MDA5+ Dermatomyositis Is Associated with Stronger Skin Type I Interferon Transcriptomic Signature with Upregulation of IFN- $\beta$ Transcript. Journal of Investigative Dermatology, 2020, 140, 1276-1279.e7.                         | 0.7  | 30        |
| 20 | In vivo anti-MUC1+ tumor activity and sequences of high-affinity anti-MUC1-SEA antibodies. Cancer Immunology, Immunotherapy, 2020, 69, 1337-1352.  | 4.2  | 15        |
| 21 | Effect of expression of ICOS in cutaneous T-cell lymphoma and its targeting on killing of malignant cells.. Journal of Clinical Oncology, 2020, 38, e20040-e20040.   | 1.6  | 2         |
| 22 | Atypical BRAF and NRAS Mutations in Mucosal Melanoma. Cancers, 2019, 11, 1133.   | 3.7  | 47        |
| 23 | Cutaneous T-cell lymphoma cells release proapoptotic Fas ligand in lysosomal secretory vesicles. European Journal of Cancer, 2019, 119, S17.   | 2.8  | 0         |
| 24 | 671 Study of the molecular and functional effects of wound dressings on human dermal fibroblasts. Journal of Investigative Dermatology, 2019, 139, S330.   | 0.7  | 0         |
| 25 | Identification of CD39 as a Marker for the Circulating Malignant T-Cell Clone of SÅ©zary Syndrome Patients. Journal of Investigative Dermatology, 2019, 139, 725-728.  | 0.7  | 6         |
| 26 | IPH4102, a first-in-class anti-KIR3DL2 monoclonal antibody, in patients with relapsed or refractory cutaneous T-cell lymphoma: an international, first-in-human, open-label, phase 1 trial. Lancet Oncology, The, 2019, 20, 1160-1170. | 10.7 | 119       |
| 27 | Blocking Antibodies Targeting the CD39/CD73 Immunosuppressive Pathway Unleash Immune Responses in Combination Cancer Therapies. Cell Reports, 2019, 27, 2411-2425.e9.  | 6.4  | 274       |
| 28 | Microenvironment tailors nTreg structure and function. Proceedings of the National Academy of Sciences of the United States of America, 2019, 116, 6298-6307.  | 7.1  | 22        |
| 29 | Extracellular Vesicles Released by Allogeneic Human Cardiac Stem/Progenitor Cells as Part of Their Therapeutic Benefit. Stem Cells Translational Medicine, 2019, 8, 911-924.   | 3.3  | 12        |
| 30 | Soluble Fc-Disabled Herpes Virus Entry Mediator Augments Activation and Cytotoxicity of NK Cells by Promoting Cross-Talk between NK Cells and Monocytes. Journal of Immunology, 2019, 202, 2057-2068.                                  | 0.8  | 0         |
| 31 | 664 Biological activities of traditional medicinal herbs on skin cells. Journal of Investigative Dermatology, 2019, 139, S329.   | 0.7  | 0         |
| 32 | Increased CD8+CD28- circulating T cells and high blood interferon score characterize the systemic inflammation of amyopathic dermatomyositis. Journal of the American Academy of Dermatology, 2019, 85, 755-758.                       | 1.2  | 1         |
| 33 | Increased expression of <sup>PD</sup>1 and <sup>CD</sup>39 on <sup>CD</sup>3<sup>+</sup><sup>CD</sup>4<sup>+</sup> skin T cells in the elderly. Experimental Dermatology, 2019, 28, 80-82.   | 2.9  | 10        |
| 34 | Truncating mutations of <i>TP53AIP1</i> gene predispose to cutaneous melanoma. Genes Chromosomes and Cancer, 2018, 57, 294-303.  | 2.8  | 8         |
| 35 | Cutaneous presentation of adult T-cell leukemia/lymphoma (ATLL). Single-center study on 37 patients in metropolitan France between 1996 and 2016. Annales De Dermatologie Et De Venereologie, 2018, 145, 405-412.                      | 1.0  | 10        |
| 36 | Impact of prednisone in patients with repeated embryo implantation failures: Beneficial or deleterious?. Journal of Reproductive Immunology, 2018, 127, 11-15.   | 1.9  | 36        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | The Interleukin-17 Family of Cytokines in Breast Cancer. International Journal of Molecular Sciences, 2018, 19, 3880.  | 4.1 | 50        |
| 38 | Cytokine levels in persistent skin lesions of adult-onset Still disease. Journal of the American Academy of Dermatology, 2018, 79, 947-949.  | 1.2 | 8         |
| 39 | Revisiting blood classification in Mycosis Fungoides and SÅ©zary syndrome with the KIR3DL2 marker. European Journal of Cancer, 2018, 101, S10-S11.   | 2.8 | 0         |
| 40 | KIR3DL2 expression in patients with adult T-cell lymphoma/leukaemia (ATLL). European Journal of Cancer, 2018, 101, S7-S8.  | 2.8 | 0         |
| 41 | Anti-CD160, Alone or in Combination With Bevacizumab, Is a Potent Inhibitor of Ocular Neovascularization in Rabbit and Monkey Models. , 2018, 59, 2687.  |     | 7         |
| 42 | Triple-negative and HER2-overexpressing breast cancer cell sialylation impacts tumor microenvironment T-lymphocyte subset recruitment: a possible mechanism of tumor escape. Cancer Management and Research, 2018, Volume 10, 1051-1059. | 1.9 | 8         |
| 43 | CD160 Expression in Retinal Vessels Is Associated With Retinal Neovascular Diseases. , 2018, 59, 2679.   |     | 6         |
| 44 | Argx-110 for Treatment of CD70-Positive Advanced Cutaneous T-Cell Lymphoma in a Phase 1/2 Clinical Trial. Blood, 2018, 132, 1627-1627.   | 1.4 | 6         |
| 45 | CD160. , 2018, , 846-852.  |     | 0         |
| 46 | PDE4D promotes FAK-mediated cell invasion in BRAF-mutated melanoma. Oncogene, 2017, 36, 3252-3262.   | 5.9 | 25        |
| 47 | Minimizing the risk of allo-sensitization to optimize the benefit of allogeneic cardiac-derived stem/progenitor cells. Scientific Reports, 2017, 7, 41125.   | 3.3 | 14        |
| 48 | Usefulness of KIR3DL2 to Diagnose, Follow-Up, and Manage the Treatment of Patients with SÅ©zary Syndrome. Clinical Cancer Research, 2017, 23, 3619-3627.   | 7.0 | 41        |
| 49 | Study of gene expression alteration in male androgenetic alopecia: evidence of predominant molecular signalling pathways. British Journal of Dermatology, 2017, 177, 1322-1336.  | 1.5 | 44        |
| 50 | KIR3DL2 expression in cutaneous T-cell lymphomas: expanding the spectrum for KIR3DL2 targeting. Blood, 2017, 130, 2900-2902.   | 1.4 | 30        |
| 51 | Chemotherapy treatment induces an increase of autophagy in the luminal breast cancer cell MCF7, but not in the triple-negative MDA-MB231. Scientific Reports, 2017, 7, 7201.   | 3.3 | 39        |
| 52 | Phase I Study of IPH4102, Anti-KIR3DL2 Mab, in Relapsed/Refractory Cutaneous T-Cell Lymphomas (CTCL): Dose-escalation Safety, Biomarker and Clinical Activity Results. Hematological Oncology, 2017, 35, 48-49.                          | 1.7 | 8         |
| 53 | Circulating and skin-derived SÅ©zary cells: clonal but with phenotypic plasticity. Blood, 2017, 130, 1468-1471.  | 1.4 | 44        |
| 54 | Dermatopulmonary Syndrome Associated With Anti-MDA5 Antibodies After Allogeneic Hematopoietic Stem Cell Transplantation. JAMA Dermatology, 2017, 153, 184.   | 4.1 | 17        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Uterine immune profiling for increasing live birth rate: A one-to-one matched cohort study. <i>Journal of Reproductive Immunology</i> , 2017, 119, 23-30.   | 1.9 | 47        |
| 56 | Therapeutic Antibodies to KIR3DL2 and Other Target Antigens on Cutaneous T-Cell Lymphomas. <i>Frontiers in Immunology</i> , 2017, 8, 1010.  | 4.8 | 18        |
| 57 | Human Cardiac-Derived Stem/Progenitor Cells Fine-Tune Monocyte-Derived Descendants Activities toward Cardiac Repair. <i>Frontiers in Immunology</i> , 2017, 8, 1413.  | 4.8 | 12        |
| 58 | Up-and-down immunity of pregnancy in humans. <i>F1000Research</i> , 2017, 6, 1216.  | 1.6 | 36        |
| 59 | The IL-17B-IL-17 receptor B pathway promotes resistance to paclitaxel in breast tumors through activation of the ERK1/2 pathway. <i>Oncotarget</i> , 2017, 8, 113360-113372.  | 1.8 | 33        |
| 60 | Interleukin 17 in the tumor microenvironment: A potent target for anticancer immunotherapy?. <i>Journal of Clinical Oncology</i> , 2017, 35, 115-115.   | 1.6 | 1         |
| 61 | Abstract 1602: Generation of anti-IL-17B antibodies neutralizing IL-17B-mediated alterations of the immune microenvironment, promotion of tumor cell initiating capacity and chemoresistance. , 2017, , .                 |     | 0         |
| 62 | TERT promoter mutations in melanoma render TERT expression dependent on MAPK pathway activation. <i>Oncotarget</i> , 2016, 7, 53127-53136.  | 1.8 | 54        |
| 63 | Targeting the Tumor Microenvironment: The Protumor Effects of IL-17 Related to Cancer Type. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1433.  | 4.1 | 104       |
| 64 | Evaluation of Immunophenotypic and Molecular Biomarkers for SÅ©zary Syndrome Using Standard Operating Procedures: A Multicenter Study of 59 Patients. <i>Journal of Investigative Dermatology</i> , 2016, 136, 1364-1372. | 0.7 | 78        |
| 65 | Evidence of <scp>T</scp>h1, <scp>T</scp>h17 and <scp>T</scp>c17 cells in psoriasiform chronic graft-versus-host disease. <i>Experimental Dermatology</i> , 2016, 25, 64-65.   | 2.9 | 10        |
| 66 | The Uterine Immune Profile May Help Women With Repeated Unexplained Embryo Implantation Failure After <i>In Vitro</i> Fertilization. <i>American Journal of Reproductive Immunology</i> , 2016, 75, 388-401.              | 1.2 | 143       |
| 67 | APRIL levels are associated with disease activity in human chronic graft-versus-host disease. <i>Haematologica</i> , 2016, 101, e312-e315.  | 3.5 | 9         |
| 68 | Phenotypic and functional changes in dermal primary fibroblasts isolated from intrinsically aged human skin. <i>Experimental Dermatology</i> , 2016, 25, 113-119.   | 2.9 | 46        |
| 69 | Intrinsically aged dermal fibroblasts fail to differentiate into adipogenic lineage. <i>Experimental Dermatology</i> , 2016, 25, 906-909.   | 2.9 | 1         |
| 70 | Expression of SÅ©zary Biomarkers in the Blood of Patients with Erythrodermic Mycosis Fungoides. <i>Journal of Investigative Dermatology</i> , 2016, 136, 317-320.   | 0.7 | 16        |
| 71 | Tremâ€ is not crucial in psoriasiform imiquimodâ€induced skin inflammation in mice. <i>Experimental Dermatology</i> , 2016, 25, 400-402.  | 2.9 | 6         |
| 72 | <i>PARKIN</i> Inactivation Links Parkinsonâ€™s Disease to Melanoma. <i>Journal of the National Cancer Institute</i> , 2016, 108, djv340.  | 6.3 | 56        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 73 | CD160. , 2016, , 1-7.   |     | 2         |
| 74 | First-in-Human, Multicenter Phase I Study of IPH4102, First-in-Class Humanized Anti-KIR3DL2 Monoclonal Antibody, in Relapsed/Refractory Cutaneous T-Cell Lymphomas: Preliminary Safety, Exploratory and Clinical Activity Results. <i>Blood</i> , 2016, 128, 1826-1826. | 1.4 | 6         |
| 75 | MUC1-ARF <sup>Δ</sup> A Novel MUC1 Protein That Resides in the Nucleus and Is Expressed by Alternate Reading Frame Translation of MUC1 mRNA. <i>PLoS ONE</i> , 2016, 11, e0165031.  | 2.5 | 11        |
| 76 | IL-17E synergizes with EGF and confers <i>in vitro</i> resistance to EGFR-targeted therapies in TNBC cells. <i>Oncotarget</i> , 2016, 7, 53350-53361.   | 1.8 | 23        |
| 77 | RICTOR involvement in the PI3K/AKT pathway regulation in melanocytes and melanoma. <i>Oncotarget</i> , 2015, 6, 28120-28131.  | 1.8 | 26        |
| 78 | CD39: A complementary target to immune checkpoints to counteract tumor-mediated immunosuppression. <i>Oncoimmunology</i> , 2015, 4, e1003015.   | 4.6 | 33        |
| 79 | Inhibition of CD39 Enzymatic Function at the Surface of Tumor Cells Alleviates Their Immunosuppressive Activity. <i>Cancer Immunology Research</i> , 2015, 3, 254-265.  | 3.4 | 190       |
| 80 | KIR3DL2/CpG ODN Interaction Mediates S <sup>Δ</sup> zary Syndrome Malignant T Cell Apoptosis. <i>Journal of Investigative Dermatology</i> , 2015, 135, 229-237.   | 0.7 | 14        |
| 81 | IL-17A and its homologs IL-25/IL-17E recruit the c-RAF/S6 kinase pathway and the generation of pro-oncogenic LMW-E in breast cancer cells. <i>Scientific Reports</i> , 2015, 5, 11874.  | 3.3 | 45        |
| 82 | Deficient regulatory B cells in human chronic graft-versus-host disease. <i>Oncoimmunology</i> , 2015, 4, e1016707.   | 4.6 | 11        |
| 83 | Authors' Reply. <i>American Journal of Pathology</i> , 2015, 185, 1168.   | 3.8 | 1         |
| 84 | CD24 <sup>hi</sup> CD27 <sup>+</sup> and plasmablast-like regulatory B cells in human chronic graft-versus-host disease. <i>Blood</i> , 2015, 125, 1830-1839.   | 1.4 | 144       |
| 85 | Colony Stimulating Factors 1, 2, 3 and early pregnancy steps: from bench to bedside. <i>Journal of Reproductive Immunology</i> , 2015, 109, 1-6.  | 1.9 | 33        |
| 86 | A novel targeted immunotherapy for CTCL is on its way: Anti-KIR3DL2 mAb IPH4102 is potent and safe in non-clinical studies. <i>Oncoimmunology</i> , 2015, 4, e1022306.  | 4.6 | 21        |
| 87 | CD Nomenclature 2015: Human Leukocyte Differentiation Antigen Workshops as a Driving Force in Immunology. <i>Journal of Immunology</i> , 2015, 195, 4555-4563.  | 0.8 | 125       |
| 88 | Genes involved in the <i>WNT</i> and vesicular trafficking pathways are associated with melanoma predisposition. <i>International Journal of Cancer</i> , 2015, 136, 2109-2119.   | 5.1 | 27        |
| 89 | CD158k Is a Reliable Marker for Diagnosis of S <sup>Δ</sup> zary Syndrome and Reveals an Unprecedented Heterogeneity of Circulating Malignant Cells. <i>Journal of Investigative Dermatology</i> , 2015, 135, 247-257.  | 0.7 | 56        |
| 90 | Autophagy is decreased in triple-negative breast carcinoma involving likely the MUC1-EGFR-NEU1 signalling pathway. <i>International Journal of Clinical and Experimental Pathology</i> , 2015, 8, 4344-55.  | 0.5 | 8         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 91  | T-Plastin Expression Downstream to the Calcineurin/NFAT Pathway Is Involved in Keratinocyte Migration. PLoS ONE, 2014, 9, e104700.  | 2.5 | 15        |
| 92  | IPH4102, a Humanized KIR3DL2 Antibody with Potent Activity against Cutaneous T-cell Lymphoma. Cancer Research, 2014, 74, 6060-6070.   | 0.9 | 65        |
| 93  | NKp46-Specific Expression on Skin-Resident CD4 + Lymphocytes in Mycosis Fungoides and SÅ©zary Syndrome. Journal of Investigative Dermatology, 2014, 134, 574-578.   | 0.7 | 3         |
| 94  | A Large French Case-Control Study Emphasizes the Role of Rare<i>Mc1R</i>Variants in Melanoma Risk. BioMed Research International, 2014, 2014, 1-10.   | 1.9 | 19        |
| 95  | Contribution of <sc>CD</sc>39 to the immunosuppressive microenvironment of acute myeloid leukaemia at diagnosis. British Journal of Haematology, 2014, 165, 722-725.  | 2.5 | 26        |
| 96  | Membrane expression of NK receptors CD160 and CD158k contributes to delineate a unique CD4<sup>+</sup>Tâ€lymphocyte subset in normal and mycosis fungoides skin. Cytometry Part A: the Journal of the International Society for Analytical Cytology, 2014, 85, 869-882. | 1.5 | 16        |
| 97  | Regulatory T-Cells in Pregnancy: Historical Perspective, State of the Art, and Burning Questions. Frontiers in Immunology, 2014, 5, 389.  | 4.8 | 79        |
| 98  | Lymphocyte-derived interleukin-17A adds another brick in the wall of inflammation-induced breast carcinogenesis. Oncoimmunology, 2014, 3, e28273.   | 4.6 | 29        |
| 99  | HACE1, a Potential Tumor Suppressor Gene on 6q21, Is Not Involved in Extranodal Natural Killer/T-Cell Lymphoma Pathophysiology. American Journal of Pathology, 2014, 184, 2899-2907.  | 3.8 | 13        |
| 100 | KIR3DL2 is a coinhibitory receptor on SÅ©zary syndrome malignant T cells that promotes resistance to activation-induced cell death. Blood, 2014, 124, 3330-3332.  | 1.4 | 22        |
| 101 | Granulocyte-Colony Stimulating Factor Related Pathways Tested on an Endometrial Ex-Vivo Model. PLoS ONE, 2014, 9, e102286.  | 2.5 | 53        |
| 102 | Seminal plasma peptides may determine maternal immune response that alters success or failure of pregnancy in the abortion-prone CBAxDBA/2 model. Journal of Reproductive Immunology, 2013, 99, 46-53.  | 1.9 | 28        |
| 103 | MUC1/CD227 IMMUNOHISTOCHEMISTRY IN ROUTINE PRACTICE IS A USEFUL BIOMARKER IN BREAST CANCERS. Journal of Immunoassay and Immunochemistry, 2013, 34, 232-245.   | 1.1 | 2         |
| 104 | Genetic variation at <sc><i>KIT</i></sc> locus may predispose to melanoma. Pigment Cell and Melanoma Research, 2013, 26, 88-96.   | 3.3 | 5         |
| 105 | IL-17A is produced by breast cancer TILs and promotes chemoresistance and proliferation through ERK1/2. Scientific Reports, 2013, 3, 3456.  | 3.3 | 119       |
| 106 | Inducible expression and pathophysiologic functions of T-plastin in cutaneous T-cell lymphoma. Blood, 2012, 120, 143-154.   | 1.4 | 33        |
| 107 | Active and Passive Anticytokine Immune Therapies: Current Status and Development. Advances in Immunology, 2012, 115, 187-227.   | 2.2 | 9         |
| 108 | TWEAK Affects Keratinocyte G2/M Growth Arrest and Induces Apoptosis through the Translocation of the AIF Protein to the Nucleus. PLoS ONE, 2012, 7, e33609.   | 2.5 | 41        |



| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 109 | CD158k/KIR3DL2 and NKp46 are frequently expressed in transformed mycosis fungoides. <i>Experimental Dermatology</i> , 2012, 21, 461-463.  | 2.9 | 36        |
| 110 | Active Chronic Sarcoidosis is Characterized by Increased Transitional Blood B Cells, Increased IL-10-Producing Regulatory B Cells and High BAFF Levels. <i>PLoS ONE</i> , 2012, 7, e43588.  | 2.5 | 78        |
| 111 | Assessment of tyrosinase variants and skin cancer risk in a large cohort of French subjects. <i>Journal of Dermatological Science</i> , 2011, 64, 127-133.  | 1.9 | 17        |
| 112 | CD101 Expression and Function in Normal and Rheumatoid Arthritis-affected Human T Cells and Monocytes/Macrophages. <i>Journal of Rheumatology</i> , 2011, 38, 419-428.  | 2.0 | 16        |
| 113 | Engagement of IL-1 receptor accessory protein (IL-1RAcP) with the monoclonal antibody AY19 provides co-activating signals and prolongs the CD2-induced proliferation of peripheral blood lymphocytes. <i>Immunology Letters</i> , 2011, 139, 52-57.                         | 2.5 | 5         |
| 114 | Death ligand TRAIL, secreted by CD1a+ and CD14+ cells in blister fluids, is involved in killing keratinocytes in toxic epidermal necrolysis. <i>Experimental Dermatology</i> , 2011, 20, 107-112.   | 2.9 | 35        |
| 115 | ERK and PDE4 cooperate to induce RAF isoform switching in melanoma. <i>Nature Structural and Molecular Biology</i> , 2011, 18, 584-591.   | 8.2 | 81        |
| 116 | Differential and tumor-specific expression of CD160 in B-cell malignancies. <i>Blood</i> , 2011, 118, 2174-2183.  | 1.4 | 47        |
| 117 | CD160: A unique activating NK cell receptor. <i>Immunology Letters</i> , 2011, 138, 93-96.  | 2.5 | 81        |
| 118 | IFN- $\gamma$ and CD46 stimulation are associated with active lupus and skew natural T regulatory cell differentiation to type 1 regulatory T (Tr1) cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 18995-19000. | 7.1 | 52        |
| 119 | A novel antiangiogenic and vascular normalization therapy targeted against human CD160 receptor. <i>Journal of Experimental Medicine</i> , 2011, 208, 973-986.  | 8.5 | 46        |
| 120 | Human endothelial cells generate Th17 and regulatory T cells under inflammatory conditions. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 2891-2896.  | 7.1 | 107       |
| 121 | Extranodal NK/T-Cell Lymphoma: Toward the Identification of Clinical Molecular Targets. <i>Journal of Biomedicine and Biotechnology</i> , 2011, 2011, 1-11.   | 3.0 | 19        |
| 122 | CD39/Adenosine Pathway Is Involved in AIDS Progression. <i>PLoS Pathogens</i> , 2011, 7, e1002110.  | 4.7 | 154       |
| 123 | Expression and Function of the Natural Cytotoxicity Receptor NKp46 on Circulating Malignant CD4+ T Lymphocytes of SÅ©zary Syndrome Patients. <i>Journal of Investigative Dermatology</i> , 2011, 131, 969-976.  | 0.7 | 41        |
| 124 | Human and Mouse Mast Cells Express and Secrete the GPI-Anchored Isoform of CD160. <i>Journal of Investigative Dermatology</i> , 2011, 131, 916-924.   | 0.7 | 23        |
| 125 | Two Domains of Vimentin Are Expressed on the Surface of Lymph Node, Bone and Brain Metastatic Prostate Cancer Lines along with the Putative Stem Cell Marker Proteins CD44 and CD133. <i>Cancers</i> , 2011, 3, 2870-2885.  | 3.7 | 36        |
| 126 | Histopathologic Diagnosis of Lymphomatous Versus Inflammatory Erythroderma: A Morphologic and Phenotypic Study on 47 Skin Biopsies. <i>American Journal of Dermatopathology</i> , 2010, 32, 755-763.  | 0.6 | 51        |



| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 127 | CD160 signaling mediates PI3K-dependent survival and growth signals in chronic lymphocytic leukemia. <i>Blood</i> , 2010, 115, 3079-3088.   | 1.4 | 48        |
| 128 | A novel KIR-associated function: evidence that CpG DNA uptake and shuttling to early endosomes is mediated by KIR3DL2. <i>Blood</i> , 2010, 116, 1637-1647.   | 1.4 | 83        |
| 129 | IL-10 produced by activated human B cells regulates CD4 <sup>+</sup> T cell activation <i>in vitro</i> . <i>European Journal of Immunology</i> , 2010, 40, 2686-2691.   | 2.9 | 216       |
| 130 | IFN- $\gamma$ kinoid vaccine-induced neutralizing antibodies prevent clinical manifestations in a lupus flare murine model. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 5294-5299.  | 7.1 | 205       |
| 131 | Identification and Characterization of a Transmembrane Isoform of CD160 (CD160-TM), a Unique Activating Receptor Selectively Expressed upon Human NK Cell Activation. <i>Journal of Immunology</i> , 2009, 182, 63-71.  | 0.8 | 58        |
| 132 | NKG2D Ligands Expression and NKG2D-Mediated NK Activity in Sezary Patients. <i>Journal of Investigative Dermatology</i> , 2009, 129, 359-364.   | 0.7 | 16        |
| 133 | Regulatory T cells differentially modulate the maturation and apoptosis of human CD8 <sup>+</sup> T-cell subsets. <i>Blood</i> , 2009, 113, 4556-4565.  | 1.4 | 54        |
| 134 | CD158K/KIR3DL2 Transcript Detection in Lesional Skin of Patients with Erythroderma Is a Tool for the Diagnosis of SÅ©zary Syndrome. <i>Journal of Investigative Dermatology</i> , 2008, 128, 465-472.   | 0.7 | 51        |
| 135 | NK cells and surveillance in humans. <i>Reproductive BioMedicine Online</i> , 2008, 16, 192-201.  | 2.4 | 14        |
| 136 | Critical and Differential Roles of NKp46- and NKp30-Activating Receptors Expressed by Uterine NK Cells in Early Pregnancy. <i>Journal of Immunology</i> , 2008, 181, 3009-3017.   | 0.8 | 125       |
| 137 | Cutting Edge: Selective Expression of Inhibitory or Activating Killer Cell Ig-Like Receptors in Circulating CD4 <sup>+</sup> T Lymphocytes. <i>Journal of Immunology</i> , 2008, 180, 2767-2771.  | 0.8 | 15        |
| 138 | A Soluble Form of the MHC Class I-Specific CD160 Receptor Is Released from Human Activated NK Lymphocytes and Inhibits Cell-Mediated Cytotoxicity. <i>Journal of Immunology</i> , 2007, 178, 1293-1300.   | 0.8 | 51        |
| 139 | Increased Number of Cytotoxic CD3 <sup>+</sup> CD28 <sup>â€</sup> T Cells in Peripheral Blood of Patients with Cutaneous Malignant Melanoma. <i>Dermatology</i> , 2007, 214, 283-288.   | 2.1 | 13        |
| 140 | VEGF kinoid vaccine, a therapeutic approach against tumor angiogenesis and metastases. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 2837-2842.   | 7.1 | 193       |
| 141 | CD160-activating NK cell effector functions depend on the phosphatidylinositol 3-kinase recruitment. <i>International Immunology</i> , 2007, 19, 401-409.   | 4.0 | 42        |
| 142 | Killer cell Ig-like receptors CD158a and CD158b display a coactivatory function, involving the c-Jun NH2-terminal protein kinase signaling pathway, when expressed on malignant CD4 <sup>+</sup> T cells from a patient with SÅ©zary syndrome. <i>Blood</i> , 2007, 109, 5064-5065. | 1.4 | 23        |
| 143 | Control of allergic reactions in mice by an active anti-murine IL-4 immunization. <i>Vaccine</i> , 2007, 25, 7206-7216.   | 3.8 | 21        |
| 144 | Identification of a Novel CD160 <sup>+</sup> CD4 <sup>+</sup> T-Lymphocyte Subset in the Skin: A Possible Role for CD160 in Skin Inflammation. <i>Journal of Investigative Dermatology</i> , 2007, 127, 1161-1166.  | 0.7 | 31        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 145 | Soluble HLA-G and control of angiogenesis. <i>Journal of Reproductive Immunology</i> , 2007, 76, 17-22.  | 1.9 | 44        |
| 146 | Recent discoveries in the genetics of melanoma and their therapeutic implications. <i>Archivum Immunologiae Et Therapiae Experimentalis</i> , 2007, 55, 363-372.   | 2.3 | 19        |
| 147 | Engagement of the CD160 activating NK cell receptor leads to its association with CD2 in circulating human NK cells. <i>Transplant Immunology</i> , 2006, 17, 36-38.   | 1.2 | 14        |
| 148 | Significance of circulating T-cell clones in Sezary syndrome. <i>Blood</i> , 2006, 107, 4030-4038.   | 1.4 | 69        |
| 149 | Soluble HLA-G1 inhibits angiogenesis through an apoptotic pathway and by direct binding to CD160 receptor expressed by endothelial cells. <i>Blood</i> , 2006, 108, 2608-2615.   | 1.4 | 181       |
| 150 | Heterogeneous Abnormalities of CCND1 and RB1 in Primary Cutaneous T-Cell Lymphomas Suggesting Impaired Cell Cycle Control in Disease Pathogenesis. <i>Journal of Investigative Dermatology</i> , 2006, 126, 1388-1395.   | 0.7 | 33        |
| 151 | Antiproliferative Effect of Semaphorin 3F on Human Melanoma Cell Lines. <i>Journal of Investigative Dermatology</i> , 2006, 126, 2343-2345.  | 0.7 | 19        |
| 152 | The co-expression of 2B4 (CD244) and CD160 delineates a subpopulation of human CD8+ T cells with a potent CD160-mediated cytolytic effector function. <i>European Journal of Immunology</i> , 2006, 36, 2359-2366.   | 2.9 | 55        |
| 153 | SC5 mAb Represents a Unique Tool for the Detection of Extracellular Vimentin as a Specific Marker of Sezary Cells. <i>Journal of Immunology</i> , 2006, 176, 652-659.  | 0.8 | 63        |
| 154 | CD molecules 2005: human cell differentiation molecules. <i>Blood</i> , 2005, 106, 3123-3126.  | 1.4 | 110       |
| 155 | Circulating Natural Killer Lymphocytes Are Potential Cytotoxic Effectors Against Autologous Malignant Cells in Sezary Syndrome Patients. <i>Journal of Investigative Dermatology</i> , 2005, 125, 1273-1278.   | 0.7 | 37        |
| 156 | The CD160+ CD8high cytotoxic T cell subset correlates with response to HAART in HIV-1+ patients. <i>Cellular Immunology</i> , 2005, 237, 96-105.   | 3.0 | 28        |
| 157 | T-cell: Section report. <i>Cellular Immunology</i> , 2005, 236, 3-5.   | 3.0 | 1         |
| 158 | SC3 monoclonal antibody defines a novel specific human B-cell surface antigen differentially expressed on B-cell leukaemias and lymphomas and involved in the proliferation of normal and malignant B lymphocytes. <i>Cellular Immunology</i> , 2005, 236, 92-100. | 3.0 | 0         |
| 159 | HLA Class I/NK Cell Receptor Interaction in Early Human Decidua basalis: Possible Functional Consequences. , 2005, 89, 72-83.  |     | 20        |
| 160 | Soluble CD100 functions on human monocytes and immature dendritic cells require plexin C1 and plexin B1, respectively. <i>International Immunology</i> , 2005, 17, 439-447.  | 4.0 | 84        |
| 161 | Cutting Edge: Engagement of CD160 by its HLA-C Physiological Ligand Triggers a Unique Cytokine Profile Secretion in the Cytotoxic Peripheral Blood NK Cell Subset. <i>Journal of Immunology</i> , 2004, 173, 5349-5354.  | 0.8 | 105       |
| 162 | CD158k/KIR3DL2 Is a New Phenotypic Marker of Sezary Cells: Relevance for the Diagnosis and Follow-Up of Sezary Syndrome. <i>Journal of Investigative Dermatology</i> , 2004, 122, 820-823.   | 0.7 | 135       |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 163 | Evaluation of the Potential Role of Cytokines in Toxic Epidermal Necrolysis. <i>Journal of Investigative Dermatology</i> , 2004, 123, 850-855.   | 0.7 | 152       |
| 164 | Sezary Syndrome Cells Unlike Normal Circulating T Lymphocytes Fail to Migrate Following Engagement of NT1 Receptor. <i>Journal of Investigative Dermatology</i> , 2004, 122, 111-118.  | 0.7 | 16        |
| 165 | Toxic epidermal necrolysis: Effector cells are drug-specific cytotoxic T cells. <i>Journal of Allergy and Clinical Immunology</i> , 2004, 114, 1209-1215.  | 2.9 | 353       |
| 166 | Semaphorin CD100 from Activated T Lymphocytes Induces Process Extension Collapse in Oligodendrocytes and Death of Immature Neural Cells. <i>Journal of Immunology</i> , 2004, 172, 1246-1255.  | 0.8 | 97        |
| 167 | Killer cell immunoglobulin-like receptor expression delineates in situ Sezary syndrome lymphocytes. <i>Journal of Pathology</i> , 2003, 199, 77-83.  | 4.5 | 47        |
| 168 | Functional and molecular characterization of a KIR3DL2/p140 expressing tumor-specific cytotoxic T lymphocyte clone infiltrating a human lung carcinoma. <i>Oncogene</i> , 2003, 22, 7192-7198.   | 5.9 | 22        |
| 169 | Immunopathogenesis of cutaneous T-cell lymphomas. <i>Hematology/Oncology Clinics of North America</i> , 2003, 17, 1313-1317.   | 2.2 | 7         |
| 170 | Polymorphic expression of CD158k/p140/KIR3DL2 in Sezary patients. <i>Blood</i> , 2003, 101, 1203-1203.   | 1.4 | 9         |
| 171 | Structure and Function of the Immune Semaphorin CD100/SEMA4D. <i>Critical Reviews in Immunology</i> , 2003, 23, 65-81.   | 0.5 | 35        |
| 172 | Engagement of CD160 receptor by HLA-C is a triggering mechanism used by circulating natural killer (NK) cells to mediate cytotoxicity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 16963-16968. | 7.1 | 128       |
| 173 | CD Antigens 2001. <i>Modern Pathology</i> , 2002, 15, 71-76.   | 5.5 | 7         |
| 174 | BY55/CD160 acts as a coreceptor in TCR signal transduction of a human circulating cytotoxic effector T lymphocyte subset lacking CD28 expression. <i>International Immunology</i> , 2002, 14, 445-451.   | 4.0 | 88        |
| 175 | Engagement of ILT2/CD85j in Sezary syndrome cells inhibits their CD3/TCR signaling. <i>Blood</i> , 2002, 100, 1019-1025.   | 1.4 | 31        |
| 176 | Drug Specific Cytotoxic T-Cells in the Skin Lesions of a Patient with Toxic Epidermal Necrolysis. <i>Journal of Investigative Dermatology</i> , 2002, 118, 728-733.  | 0.7 | 260       |
| 177 | CD4+ cutaneous T-cell lymphoma cells express the p140 killer cell immunoglobulin-like receptor. <i>Blood</i> , 2001, 97, 1388-1391.  | 1.4 | 119       |
| 178 | Increased Expression of a Novel Early Activation Surface Membrane Receptor in Cutaneous T Cell Lymphoma Cells. <i>Journal of Investigative Dermatology</i> , 2001, 116, 731-738.   | 0.7 | 27        |
| 179 | Functional Characterization of Neurotensin Receptors in Human Cutaneous T Cell Lymphoma Malignant Lymphocytes. <i>Journal of Investigative Dermatology</i> , 2001, 117, 687-693.   | 0.7 | 18        |
| 180 | CD antigens 2001. <i>European Journal of Immunology</i> , 2001, 31, 2841-2847.   | 2.9 | 3         |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 181 | Biological Activity of Soluble CD100. II. Soluble CD100, Similarly to H-Semalll, Inhibits Immune Cell Migration. <i>Journal of Immunology</i> , 2001, 166, 4348-4354.  | 0.8 | 154       |
| 182 | Multiple co-stimulatory signals are required for triggering proliferation of T cells from human secondary lymphoid tissue. <i>International Immunology</i> , 2001, 13, 441-450.  | 4.0 | 7         |
| 183 | Biological Activity of Soluble CD100. I. The Extracellular Region of CD100 Is Released from the Surface of T Lymphocytes by Regulated Proteolysis. <i>Journal of Immunology</i> , 2001, 166, 4341-4347.  | 0.8 | 130       |
| 184 | CD antigens 2001. <i>International Immunology</i> , 2001, 13, 1095-1098.   | 4.0 | 3         |
| 185 | Triggering CD101 molecule on human cutaneous dendritic cells inhibits T cell proliferation via IL-10 production. <i>European Journal of Immunology</i> , 2000, 30, 3132-3139.  | 2.9 | 35        |
| 186 | Functional Inhibitory Receptors Expressed by a Cutaneous T Cell Lymphoma-Specific Cytolytic Clonal T Cell Population. <i>Journal of Investigative Dermatology</i> , 2000, 115, 994-999.  | 0.7 | 9         |
| 187 | Cutaneous T Cell Lymphoma Reactive CD4+ Cytotoxic T Lymphocyte Clones Display a Th1 Cytokine Profile and Use a Fas-Independent Pathway for Specific Tumor Cell Lysis. <i>Journal of Investigative Dermatology</i> , 2000, 115, 74-80.                          | 0.7 | 80        |
| 188 | No Modulation of Circulating Natural Killer Cell and Natural Killer Receptor Bearing Memory T Cell Subsets in Patients with Atopic Dermatitis. <i>Journal of Investigative Dermatology</i> , 2000, 115, 1160-1162.   | 0.7 | 4         |
| 189 | Functional characterization of an IL-7-dependent CD4+CD8 <sup>+</sup> Th3-type malignant cell line derived from a patient with a cutaneous T-cell lymphoma. <i>Blood</i> , 2000, 96, 1056-1063.  | 1.4 | 29        |
| 190 | Cutting Edge: Soluble HLA-G1 Triggers CD95/CD95 Ligand-Mediated Apoptosis in Activated CD8+ Cells by Interacting with CD8. <i>Journal of Immunology</i> , 2000, 164, 6100-6104.  | 0.8 | 422       |
| 191 | Switch in the protein tyrosine phosphatase associated with human CD100 semaphorin at terminal B-cell differentiation stage. <i>Blood</i> , 2000, 95, 965-972.  | 1.4 | 36        |
| 192 | Functional characterization of an IL-7-dependent CD4+CD8 <sup>+</sup> Th3-type malignant cell line derived from a patient with a cutaneous T-cell lymphoma. <i>Blood</i> , 2000, 96, 1056-1063.  | 1.4 | 3         |
| 193 | Production and characterization of 22 monoclonal antibodies directed against S 20499, a new potent 5-HT1A chiral agonist: influence of the hapten structure on specificity and stereorecognition. <i>Pharmaceutical Research</i> , 1999, 16, 725-735.          | 3.5 | 1         |
| 194 | Soluble CD14 acts as a negative regulator of human T cell activation and function. <i>European Journal of Immunology</i> , 1999, 29, 265-276.  | 2.9 | 111       |
| 195 | Do Primary Cutaneous Non-T Non-B CD4+CD56+ Lymphomas Belong to the Myelo-Monocytic Lineage?. <i>Journal of Investigative Dermatology</i> , 1998, 111, 1242-1244.   | 0.7 | 26        |
| 196 | T Cell Receptor (TCR) Interacting Molecule (TRIM), A Novel Disulfide-linked Dimer Associated with the TCR-CD3 $\zeta$ Complex, Recruits Intracellular Signaling Proteins to the Plasma Membrane. <i>Journal of Experimental Medicine</i> , 1998, 188, 561-575. | 8.5 | 121       |
| 197 | Isolation of Tumor-Specific Cytotoxic CD4+ and CD4+CD8dim+ T-Cell Clones Infiltrating a Cutaneous T-Cell Lymphoma. <i>Blood</i> , 1998, 91, 4331-4341.   | 1.4 | 128       |
| 198 | Isolation of Tumor-Specific Cytotoxic CD4+ and CD4+CD8dim+ T-Cell Clones Infiltrating a Cutaneous T-Cell Lymphoma. <i>Blood</i> , 1998, 91, 4331-4341.   | 1.4 | 13        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 199 | The Human Semaphorin-like Leukocyte Cell Surface Molecule CD100 Associates with a Serine Kinase Activity. <i>Journal of Biological Chemistry</i> , 1997, 272, 23515-23520.  | 3.4 | 54        |
| 200 | Interferon- $\beta$ rescues HLA class Ia cell surface expression in term villous trophoblast cells by inducing synthesis of TAP proteins. <i>European Journal of Immunology</i> , 1997, 27, 45-54.  | 2.9 | 46        |
| 201 | Endothelial cells in chorionic fetal vessels of first trimester placenta express HLA- $\beta$ . <i>European Journal of Immunology</i> , 1997, 27, 3380-3388.  | 2.9 | 152       |
| 202 | Functional Role of CD101 on Skin Dendritic Cells. <i>Advances in Experimental Medicine and Biology</i> , 1997, 417, 227-232.  | 1.6 | 0         |
| 203 | Enhanced CD3 Monoclonal Antibody Induced Proliferation of Colonic Mucosal T Lymphocytes in Crohn's Disease Patients Free of Corticosteroid or Immunosuppressor Treatment. <i>Clinical Immunology and Immunopathology</i> , 1996, 79, 20-24.                   | 2.0 | 2         |
| 204 | Activation signals are delivered through two distinct epitopes of CD100, a unique 150 kDa human lymphocyte surface structure previously defined by BB18 mAb. <i>International Immunology</i> , 1995, 7, 1-8.  | 4.0 | 61        |
| 205 | Differential proliferative responses in subsets of human CD28+ cells delineated by BB27 mAb. <i>International Immunology</i> , 1994, 6, 423-430.  | 4.0 | 19        |
| 206 | A Monoclonal Antibody to the Hodgkin's Disease-Associated Antigen CD30 Induces Activation and Long-Term Growth of Human Autoreactive $\beta$ T Cell Clone. <i>Cellular Immunology</i> , 1994, 156, 230-239.   | 3.0 | 14        |
| 207 | Identification of a novel functional 85-kD glycoprotein restricted to long-term dividing human lymphocytic lines. <i>Human Immunology</i> , 1993, 37, 31-38.  | 2.4 | 3         |
| 208 | TCR $\beta$ Bearing T Lymphocytes Infiltrating Human Primary Cutaneous Melanomas. <i>Journal of Investigative Dermatology</i> , 1992, 98, 369-374.  | 0.7 | 45        |
| 209 | The CD39 molecule defines distinct cytotoxic subsets within alloactivated human CD8-positive cells. <i>European Journal of Immunology</i> , 1992, 22, 2681-2685.  | 2.9 | 30        |
| 210 | Modulation of allogenic reaction by ticlopidine treatment. <i>International Journal of Immunopharmacology</i> , 1991, 13, 101-105.  | 1.1 | 0         |
| 211 | Proliferation of resting lymphocytes is induced by triggering T cells through an epitope common to the three CD18/CD11 leukocyte adhesion molecules. <i>Cellular Immunology</i> , 1991, 136, 519-524.   | 3.0 | 18        |
| 212 | Activation of the CD3/T cell receptor (TcR) complex or of protein kinase C potentiate adenylyl cyclase stimulation in a tumoral T cell line: involvement of two distinct intracellular pathways. <i>European Journal of Immunology</i> , 1991, 21, 2877-2882. | 2.9 | 22        |
| 213 | Identification of CD3 Associated T Cell Receptor as a Diagnostic Tool in T Cell Acute Lymphoblastic Lymphoma or Leukemia. <i>Leukemia and Lymphoma</i> , 1991, 4, 187-192.  | 1.3 | 1         |
| 214 | Immunodeficiency after Bone Marrow Transplantation can be associated with Autoreactive T-Cell Receptor gammaomega-bearing Lymphocytes. <i>Immunological Reviews</i> , 1990, 116, 5-13.  | 6.0 | 11        |
| 215 | Correlation between T cell receptor $\beta$ isotypic forms and cytotoxic activity: Analysis with human T cell clones and lines. <i>Cellular Immunology</i> , 1990, 125, 315-325.  | 3.0 | 12        |
| 216 | Selective induction of autolytic activity through the CD3 molecule. <i>European Journal of Immunology</i> , 1990, 20, 2615-2619.  | 2.9 | 10        |

| #   | ARTICLE  | IF   | CITATIONS |
|-----|--|------|-----------|
| 217 | Detection of a T cell receptor $\hat{\alpha}$ chain with an anti-TCR $\hat{\alpha}$ chain serum. <i>Clinical Immunology and Immunopathology</i> , 1990, 55, 56-66.   | 2.0  | 0         |
| 218 | Rearranging sequence located in the intron of the human T cell receptor $\hat{\beta}$ chain gene constant region. <i>European Journal of Immunology</i> , 1989, 19, 637-642.   | 2.9  | 3         |
| 219 | Immunoactive products of human placenta. II. Direct inhibition of non-MHC restricted cytolytic activity of human CD3 alpha-beta but not CD3 gamma-delta expressing T cell clones. <i>Journal of Reproductive Immunology</i> , 1989, 16, 137-150. | 1.9  | 17        |
| 220 | Molecular and Functional Characterization of Human T-Cell Clones Expressing TCR- $\hat{\beta}$ Antigen Receptor. , 1989, , 554-554.  |      | 1         |
| 221 | CD4 cytotoxic T lymphocyte differentiation. <i>Biochimie</i> , 1988, 70, 937-941.  | 2.6  | 0         |
| 222 | Nucleotide sequence of a cDNA corresponding to a new human variable region of a functionally rearranged T cell receptor gamma chain. <i>Nucleic Acids Research</i> , 1987, 15, 10059-10059.  | 14.5 | 9         |
| 223 | In vitro allostimulation of peripheral blood mononuclear cells generates autologous reactive T lymphocytes: Analysis at the clonal level. <i>Human Immunology</i> , 1986, 17, 54-60.   | 2.4  | 2         |
| 224 | What is a T-cell clone? Effect of rIFN on T-cell clone function and T-cell receptor gene rearrangement. <i>Human Immunology</i> , 1986, 17, 214-223.   | 2.4  | 10        |
| 225 | Allospecific proliferative human T-cell clones acquire the cytotoxic effector function after three months in culture, in IL-2 conditioned medium. <i>Human Immunology</i> , 1986, 17, 30-36.   | 2.4  | 6         |
| 226 | Human T lymphocyte clones with killer or natural killer activity. <i>Journal of Immunological Methods</i> , 1986, 90, 215-219.   | 1.4  | 3         |
| 227 | Interferon: a cytotoxic T lymphocyte differentiation signal. <i>European Journal of Immunology</i> , 1986, 16, 767-770.  | 2.9  | 113       |
| 228 | The human T-cell receptor. <i>Journal of Clinical Immunology</i> , 1985, 5, 141-157.   | 3.8  | 39        |
| 229 | Production and characterization of antibody probes directed at constant regions of the $\hat{\alpha}$ and $\hat{\beta}$ subunit of the human T cell receptor. <i>European Journal of Immunology</i> , 1985, 15, 821-827.                         | 2.9  | 28        |
| 230 | T3-Ti receptor triggering of T8+ suppressor T cells leads to unresponsiveness to interleukin-2. <i>Nature</i> , 1984, 311, 565-567.  | 27.8 | 49        |
| 231 | Clonotypic Surface Structure on Human T Lymphocytes: Functional and Biochemical Analysis of the Antigen Receptor Complex. <i>Immunological Reviews</i> , 1984, 81, 95-130.   | 6.0  | 82        |
| 232 | Differential reactivity of human lymphocytes allosensitized in vitro in hormonally defined medium or medium supplemented with plasma. <i>Cellular Immunology</i> , 1983, 81, 441-446.  | 3.0  | 2         |
| 233 | Regulation of the human allogeneic proliferative response in vitro. <i>Immunogenetics</i> , 1981, 14, 107-116.   | 2.4  | 3         |
| 234 | Regulation of the human allogeneic proliferative response in vitro. <i>Immunogenetics</i> , 1981, 14, 117-127.   | 2.4  | 7         |