

Young Rim Song

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2593952/publications.pdf>

Version: 2024-02-01

53
papers

876
citations

567281

15
h-index

526287

27
g-index

54
all docs

54
docs citations

54
times ranked

1362
citing authors

#	ARTICLE	IF	CITATIONS
1	Serum cystatin C to creatinine ratio is associated with sarcopenia in non-dialysis-dependent chronic kidney disease. <i>Kidney Research and Clinical Practice</i> , 2022, 41, 580-590.	2.2	10
2	Longitudinal changes in body composition are associated with all-cause mortality in patients on peritoneal dialysis. <i>Clinical Nutrition</i> , 2021, 40, 120-126.	5.0	29
3	Comparison of dynamic arterial and venous pressure between metal needles and plastic cannulas in incident hemodialysis patients with arteriovenous graft. <i>Journal of Vascular Access</i> , 2021, 22, 42-47.	0.9	4
4	Fatal anaphylaxis Due to Nafamostat Mesylate During Hemodialysis. <i>Allergy, Asthma and Immunology Research</i> , 2021, 13, 517.	2.9	5
5	Neutrophil extracellular traps and heparin-induced antibodies contribute to vascular access thrombosis in hemodialysis patients. <i>Kidney Research and Clinical Practice</i> , 2021, 40, 712-723.	2.2	1
6	Late stage 3 chronic kidney disease is an independent risk factor for sarcopenia, but not proteinuria. <i>Scientific Reports</i> , 2021, 11, 18472.	3.3	16
7	Efficacy and Safety of Plastic Cannulae Compared with Metal Needles in the Initial Use of an Arteriovenous Fistulae in Incident Hemodialysis Patients: A Randomized Controlled Study. <i>American Journal of Nephrology</i> , 2021, 52, 479-486.	3.1	5
8	When and why to start continuous renal replacement therapy in critically ill patients with acute kidney injury. <i>Kidney Research and Clinical Practice</i> , 2021, 40, 566-577.	2.2	4
9	Uremic serum damages endothelium by provoking excessive neutrophil extracellular trap formation. <i>Scientific Reports</i> , 2021, 11, 21439.	3.3	6
10	Prognostic role of circulating neutrophil extracellular traps levels for long-term mortality in new end-stage renal disease patients. <i>Clinical Immunology</i> , 2020, 210, 108263.	3.2	21
11	Serum levels of protein carbonyl, a marker of oxidative stress, are associated with overhydration, sarcopenia and mortality in hemodialysis patients. <i>BMC Nephrology</i> , 2020, 21, 281.	1.8	36
12	P0762 THE SERUM URIC ACID HAS AN INDEPENDENT ASSOCIATION WITH 1-YEAR RENAL OUTCOMES IN CHRONIC KIDNEY DISEASE PATIENT WITH HYPERTENSION. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
13	P1580 LONGITUDINAL CHANGES IN BODY COMPOSITION ARE ASSOCIATED WITH ALL-CAUSE MORTALITY IN PATIENTS ON PERITONEAL DIALYSIS. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
14	Role of peribrachial fat as a key determinant of brachial artery dilatation for successful arteriovenous fistula maturation in hemodialysis patients. <i>Scientific Reports</i> , 2020, 10, 3841.	3.3	0
15	Fat to muscle ratio as a predictor of insulin resistance and metabolic syndrome in Korean adults. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2020, 11, 710-725.	7.3	50
16	Impact of sarcopenia on long-term mortality and cardiovascular events in patients undergoing hemodialysis. <i>Korean Journal of Internal Medicine</i> , 2019, 34, 599-607.	1.7	79
17	SP407 Serum protein carbonyl levels are associated with sarcopenia and long-term outcomes in hemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.7	0
18	FP728 LOW LEAN TISSUE AND HIGH FAT TISSUE INDEX ARE ASSOCIATED WITH DECREASED QUALITY OF LIFE IN DIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.7	0

#	ARTICLE	IF	CITATIONS
19	Custom-made Doppler ultrasound flow simulator for dialysis access using continuous renal replacement therapy machine. <i>Journal of Vascular Access</i> , 2019, 20, 701-705.	0.9	1
20	Effects of Excessive Body Fat Accumulation on Long-Term Outcomes during Peritoneal Dialysis. <i>Peritoneal Dialysis International</i> , 2019, 39, 268-275.	2.3	13
21	Association of Blood Pressure at Specific Time-Points with 1-Year Renal Outcomes in Patients with Diabetic Chronic Kidney Disease. <i>Electrolyte and Blood Pressure</i> , 2019, 17, 36.	1.8	0
22	Anatomical variants of upper arm veins on preoperative mapping venography for hemodialysis access in Korean adults. <i>Journal of Vascular Access</i> , 2019, 20, 270-275.	0.9	7
23	Outcomes of vascular access in hemodialysis patients: Analysis based on the Korean National Health Insurance Database from 2008 to 2016. <i>Kidney Research and Clinical Practice</i> , 2019, 38, 391-398.	2.2	13
24	Repeated Bioimpedance Measurements Predict Prognosis of Peritoneal Dialysis Patients. <i>American Journal of Nephrology</i> , 2018, 47, 120-129.	3.1	18
25	Comparison of tolvaptan treatment between patients with the SIADH and congestive heart failure: a single-center experience. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 561-567.	1.7	7
26	The relationship between autophagy, increased neutrophil extracellular traps formation and endothelial dysfunction in chronic kidney disease. <i>Clinical Immunology</i> , 2018, 197, 189-197.	3.2	16
27	Fat-to-Lean Mass Ratio Can Predict Cardiac Events and All-Cause Mortality in Patients Undergoing Hemodialysis. <i>Annals of Nutrition and Metabolism</i> , 2018, 73, 241-249.	1.9	21
28	Long-term prognosis of end-stage renal disease patients with normal myocardial perfusion as determined by single photon emission computed tomography. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 148-156.	1.7	3
29	Current treatment status and medical costs for hemodialysis vascular access based on analysis of the Korean Health Insurance Database. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 1160-1168.	1.7	11
30	Sex differences in the association between obesity and albuminuria among Korean adults: a cross-sectional study using the Korea National Health and Nutrition Examination Survey data. <i>Clinical and Experimental Nephrology</i> , 2017, 21, 27-34.	1.6	7
31	Impact of rapid ultrafiltration rate on changes in the echocardiographic left atrial volume index in patients undergoing haemodialysis: a longitudinal observational study. <i>BMJ Open</i> , 2017, 7, e013990.	1.9	8
32	Vitamin D: a possible modifying factor linking obesity to vascular calcification in hemodialysis patients. <i>Nutrition and Metabolism</i> , 2017, 14, 27.	3.0	7
33	Increased Neutrophil Extracellular Trap Formation in Uremia Is Associated with Chronic Inflammation and Prevalent Coronary Artery Disease. <i>Journal of Immunology Research</i> , 2017, 2017, 1-10.	2.2	28
34	Acute kidney injury can predict in-hospital and long-term mortality in elderly patients undergoing hip fracture surgery. <i>PLoS ONE</i> , 2017, 12, e0176259.	2.5	28
35	Physician perceptions of blood pressure control in patients with chronic kidney disease and target blood pressure achievement rate. <i>Kidney Research and Clinical Practice</i> , 2017, 36, 349-357.	2.2	4
36	Overhydration Negatively Affects Quality of Life in Peritoneal Dialysis Patients: Evidence from a Prospective Observational Study. <i>International Journal of Medical Sciences</i> , 2016, 13, 686-695.	2.5	19

#	ARTICLE	IF	CITATIONS
37	SP475 RAPID ULTRAFILTRATION RATE IS ASSOCIATED WITH THE INCREASE OF ECHOCARDIOGRAPHIC LEFT ATRIAL VOLUME INDEX AMONG HEMODIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i251-i252.	0.7	0
38	SP622 LOW VITAMIN D AS A MODIFYING FACTOR IN THE RELATIONSHIP BETWEEN OBESITY AND VASCULAR CALCIFICATION IN HEMODIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i302-i303.	0.7	0
39	Leptin, pre-existing vascular disease, and increased arteriovenous fistula maturation failure in dialysis patients. <i>Journal of Vascular Surgery</i> , 2016, 64, 402-410.e1.	1.1	10
40	Excessive Weight Gain during the First Year of Peritoneal Dialysis Is Associated with Inflammation, Diabetes Mellitus, and a Rapid Decrease in Residual Renal Function. <i>PLoS ONE</i> , 2015, 10, e0139033.	2.5	16
41	Obesity-related decrease in intraoperative blood flow is associated with maturation failure of radiocephalic arteriovenous fistula. <i>Journal of Vascular Surgery</i> , 2015, 62, 1010-1017.e1.	1.1	23
42	Four-Year Changes in Visceral Fat Mass and the Risk of Developing Proteinuria in the General Population. <i>PLoS ONE</i> , 2015, 10, e0131119.	2.5	10
43	Proton-pump inhibitors for prevention of upper gastrointestinal bleeding in patients undergoing dialysis. <i>World Journal of Gastroenterology</i> , 2015, 21, 4919.	3.3	8
44	Increased body fat rather than body weight has harmful effects on 4-year changes of renal function in the general elderly population with a normal or mildly impaired renal function. <i>Clinical Interventions in Aging</i> , 2014, 9, 1277.	2.9	9
45	Comparison of the Efficacy and Safety Profile of Morning Administration of Controlled-release Simvastatin Versus Evening Administration of Immediate-release Simvastatin in Chronic Kidney Disease Patients With Dyslipidemia. <i>Clinical Therapeutics</i> , 2014, 36, 1182-1190.	2.5	12
46	B-type natriuretic peptide predicts an ischemic etiology of acute heart failure in patients with stage 4-5 chronic kidney disease. <i>Clinical Biochemistry</i> , 2014, 47, 344-348.	1.9	7
47	Prevalence of and factors associated with sarcopenia in elderly patients with end-stage renal disease. <i>Clinical Nutrition</i> , 2014, 33, 64-68.	5.0	212
48	Association between cardiac valvular calcification and myocardial ischemia in asymptomatic high-risk patients with end-stage renal disease. <i>Atherosclerosis</i> , 2013, 229, 369-373.	0.8	19
49	Left atrial volume index is a predictor of silent myocardial ischemia in high-risk patients with end-stage renal disease. <i>International Journal of Cardiovascular Imaging</i> , 2013, 29, 1433-1439.	1.5	9
50	Serum S100B protein is associated with depressive symptoms in patients with end-stage renal disease. <i>Clinical Biochemistry</i> , 2012, 45, 1573-1577.	1.9	15
51	Left Ventricular Diastolic Dysfunction as a Predictor of Rapid Decline of Residual Renal Function in Patients with Peritoneal Dialysis. <i>Journal of the American Society of Echocardiography</i> , 2012, 25, 411-420.	2.8	21
52	Cardiac risk assessment by gated single-photon emission computed tomography in asymptomatic end-stage renal disease patients at the start of dialysis. <i>Journal of Nuclear Cardiology</i> , 2012, 19, 438-447.	2.1	28
53	A Case of Hemocholecystitis in a Patient with Microscopic Polyangiitis. <i>Korean Journal of Medicine</i> , 2012, 83, 127.	0.3	0