## David F Porinchu

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2631561/publications.pdf

Version: 2024-02-01

36 papers 1,072 citations

331670 21 h-index 32 g-index

40 all docs

40 docs citations

times ranked

40

1211 citing authors

#	Article	IF	Citations
1	A lake sediment–based paleoecological reconstruction of late Holocene fire history and vegetation change in Great Basin National Park, Nevada, USA. Quaternary Research, 2021, 104, 28-42.	1.7	2
2	Sedimentological and Grain Size Characteristics of Two Lake Cores from Himachal Pradesh, India. Journal of Climate Change, 2021, 7, 35-51.	0.5	2
3	Temperature change as a driver of spatial patterns and longâ€ŧerm trends in chironomid (Insecta:) Tj ETQq1 1 0.	784314 r	gBT <sub>3</sub> /Overlo <mark>ck</mark>
4	A high-resolution sedimentary charcoal- and geochemistry-based reconstruction of late Holocene fire regimes in the páramo of Chirripό National Park, Costa Rica. Quaternary Research, 2020, 93, 314-329.	1.7	3
5	Mountain Temperature Changes From Embedded Sensors Spanning 2000 m in Great Basin National Park, 2006–2018. Frontiers in Earth Science, 2020, 8, .	1.8	5
6	A global database of Holocene paleotemperature records. Scientific Data, 2020, 7, 115.	<b>5.</b> 3	112
7	Evidence of abrupt climate change at 9.3 ka and 8.2 ka in the central Canadian Arctic: Connection to the North Atlantic and Atlantic Meridional Overturning Circulation. Quaternary Science Reviews, 2019, 219, 204-217.	3.0	10
8	Late Holocene hydroclimate variability in Costa Rica: Signature of the terminal classic drought and the Medieval Climate Anomaly in the northern tropical Americas. Quaternary Science Reviews, 2019, 215, 144-159.	3.0	15
9	Holocene hydroclimate and environmental change inferred from a high-resolution multi-proxy record from Lago Ditkebi, Chirripó National Park, Costa Rica. Palaeogeography, Palaeoclimatology, Palaeoecology, 2019, 518, 172-186.	2.3	8
10	A chironomid-based reconstruction of late-Holocene climate and environmental change for southern Pacific Costa Rica. Holocene, 2017, 27, 73-84.	1.7	14
11	Prolonged California aridity linked to climate warming and Pacific sea surface temperature. Scientific Reports, 2016, 6, 33325.	3.3	36
12	Chironomid-environment relations in northern North America. Journal of Paleolimnology, 2015, 54, 223-237.	1.6	40
13	The modern distribution of chironomid sub-fossils (Insecta: Diptera) in Costa Rica and the development of a regional chironomid-based temperature inference model. Hydrobiologia, 2015, 742, 107-127.	2.0	9
14	Historical trends of mercury and spheroidal carbonaceous particle deposition in sub-alpine lakes in the Great Basin, United States. Journal of Paleolimnology, 2014, 52, 405-418.	1.6	8
15	A quantitative midge-based reconstruction of mean July air temperature from a high-elevation site in central Colorado, USA, for MIS 6 and 5. Quaternary Research, 2014, 82, 580-591.	1.7	10
16	A 2000-yr reconstruction of air temperature in the Great Basin of the United States with specific reference to the Medieval Climatic Anomaly. Quaternary Research, 2014, 82, 309-317.	1.7	14
17	High-elevation paleoenvironmental change during MIS 6–4 in the central Rockies of Colorado as determined from pollen analysis. Quaternary Research, 2014, 82, 542-552.	1.7	22
18	Regional Climate Change Evidenced by Recent Shifts in Chironomid Community Composition in Subalpine and Alpine Lakes in the Great Basin of the United States. Arctic, Antarctic, and Alpine Research, 2014, 46, 600-615.	1,1	7

#	Article	IF	CITATIONS
19	Application of a midge-based inference model for air temperature reveals evidence of late-20th century warming in sub-alpine lakes in the central Great Basin, United States. Quaternary International, 2010, 215, 15-26.	1.5	14
20	A multi-proxy paleolimnological reconstruction of Holocene climate conditions in the Great Basin, United States. Quaternary Research, 2009, 72, 347-358.	1.7	47
21	The use of high-resolution gridded climate data in the development of chironomid-based inference models from remote areas. Journal of Paleolimnology, 2009, 41, 343-348.	1.6	3
22	Development of a chironomid-based air temperature inference model for the central Canadian Arctic. Journal of Paleolimnology, 2009, 41, 349-368.	1.6	41
23	Paleolimnological evidence of the response of the central Canadian treeline zone to radiative forcing and hemispheric patterns of temperature change over the past 2000Âyears. Journal of Paleolimnology, 2009, 41, 129-141.	1.6	32
24	A 2000Âyear midge-based paleotemperature reconstruction from the Canadian Arctic archipelago. Journal of Paleolimnology, 2009, 41, 177-188.	1.6	26
25	Evidence of temperature depression and hydrological variations in the eastern Sierra Nevada during the Younger Dryas Stade. Quaternary Research, 2008, 70, 131-140.	1.7	49
26	Subfossil Chironomids As Indicators Of Recent Climate Change In Sierra Nevada, California, Lakes. Arctic, Antarctic, and Alpine Research, 2007, 39, 286-296.	1.1	23
27	Development of a Midge-Based Summer Surface Water Temperature Inference Model for the Great Basin of the Western United States. Arctic, Antarctic, and Alpine Research, 2007, 39, 566-577.	1.1	17
28	A late Quaternary chironomid-inferred temperature record from the Sierra Nevada, California, with connections to northeast Pacific sea surface temperatures. Quaternary Research, 2006, 66, 356-363.	1.7	39
29	Late Pleistocene and early Holocene climate and limnological changes in the Sierra Nevada, California, USA inferred from midges (Insecta: Diptera: Chironomidae). Palaeogeography, Palaeoclimatology, Palaeoecology, 2003, 198, 403-422.	2.3	40
30	The use and application of freshwater midges (Chironomidae: Insecta: Diptera) in geographical research. Progress in Physical Geography, 2003, 27, 378-422.	3.2	73
31	The use and application of freshwater midges (Chironomidae: Insecta: Diptera) in geographical research. Progress in Physical Geography, 2003, 27, 378-422.	3.2	68
32	Title is missing!. Journal of Paleolimnology, 2002, 27, 59-69.	1.6	51
33	Title is missing!. Journal of Paleolimnology, 2002, 28, 355-375.	1.6	42
34	The Distribution of Freshwater Chironomidae (Insecto: Diptera) across Treeline near the Lower Lena River, Northeast Siberia, Russia. Arctic, Antarctic, and Alpine Research, 2000, 32, 429-437.	1.1	30
35	The Distribution of Freshwater Chironomidae (Insecta: Diptera) across Treeline near the Lower Lena River, Northeast Siberia, Russia. Arctic, Antarctic, and Alpine Research, 2000, 32, 429.	1.1	41
36	Development of a Midge-Based Summer Surface Water Temperature Inference Model for the Great Basin of the Western United States. , 0, .		1