

Ian Pallister

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2667613/publications.pdf>

Version: 2024-02-01

45
papers

1,055
citations

471509

17
h-index

414414

32
g-index

46
all docs

46
docs citations

46
times ranked

1083
citing authors

#	ARTICLE	IF	CITATIONS
1	Randomized Trial of Hydroxyethyl Starch versus Gelatine for Trauma Resuscitation. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1999, 47, 1114.	2.4	100
2	What is the best clinical test for Achilles tendinopathy?. <i>Foot and Ankle Surgery</i> , 2013, 19, 112-117.	1.7	98
3	Close Contact Casting vs Surgery for Initial Treatment of Unstable Ankle Fractures in Older Adults. <i>JAMA - Journal of the American Medical Association</i> , 2016, 316, 1455.	7.4	94
4	Increased neutrophil migratory activity after major trauma: A factor in the etiology of acute respiratory distress syndrome?. <i>Critical Care Medicine</i> , 2002, 30, 1717-1721.	0.9	78
5	The Ankle Injury Management (AIM) trial: a pragmatic, multicentre, equivalence randomised controlled trial and economic evaluation comparing close contact casting with open surgical reduction and internal fixation in the treatment of unstable ankle fractures in patients aged over 60 years. <i>Health Technology Assessment</i> . 2016. 20. 1-158.	2.8	73
6	Effect of Locking Plate Fixation vs Intramedullary Nail Fixation on 6-Month Disability Among Adults With Displaced Fracture of the Distal Tibia. <i>JAMA - Journal of the American Medical Association</i> , 2017, 318, 1767.	7.4	55
7	Prediction of Posttraumatic Adult Respiratory Distress Syndrome by Albumin Excretion Rate Eight Hours after Admission. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1997, 42, 1056-1061.	2.4	51
8	Immediate application of improvised pelvic binder as first step in extended resuscitation from life-threatening hypovolaemic shock in conscious patients with unstable pelvic injuries. <i>Injury</i> , 2007, 38, 125-128.	1.7	48
9	A qualitative approach to recovery after open tibial fracture: The road to a novel, patient-derived recovery scale. <i>Injury</i> , 2012, 43, 1071-1078.	1.7	41
10	Enhanced neutrophil migratory activity following major blunt trauma. <i>Injury</i> , 2005, 36, 956-962.	1.7	35
11	Cholelithoptysis and empyema formation after laparoscopic cholecystectomy. <i>Annals of Thoracic Surgery</i> , 1995, 60, 1100-1102.	1.3	31
12	Neutrophil Priming for Elastase Release in Adult Blunt Trauma Patients. <i>Journal of Trauma</i> , 2006, 60, 590-596.	2.3	29
13	The UK Burden of Injury Study â€” a protocol. [National Research Register number: M0044160889]. <i>BMC Public Health</i> , 2007, 7, 317.	2.9	26
14	The Effects of Surgical Fracture Fixation on the Systemic Inflammatory Response to Major Trauma. <i>Journal of the American Academy of Orthopaedic Surgeons</i> , The, 2005, 13, 93-100.	2.5	26
15	Lactoferrin inhibits dexamethasone-induced chondrocyte impairment from osteoarthritic cartilage through up-regulation of extracellular signal-regulated kinase 1/2 and suppression of FASL, FAS, and Caspase 3. <i>Biochemical and Biophysical Research Communications</i> , 2013, 441, 249-255.	2.1	25
16	Alteration of Polymorphonuclear Neutrophil Surface Receptor Expression and Migratory Activity After Isolation: Comparison of Whole Blood and Isolated PMN Preparations from Normal and Postfracture Trauma Patients. <i>Journal of Trauma</i> , 2006, 60, 844-850.	2.3	20
17	Psychological factors contributing to perceptions pain intensity after acute orthopaedic injury. <i>Injury</i> , 2011, 42, 1214-1218.	1.7	20
18	The use of secure anonymised data linkage to determine changes in healthcare utilisation following severe open tibial fractures. <i>Injury</i> , 2015, 46, 1287-1292.	1.7	16

#	ARTICLE	IF	CITATIONS
19	Intramedullary nail fixation versus locking plate fixation for adults with a fracture of the distal tibia: the UK FixDT RCT. <i>Health Technology Assessment</i> , 2018, 22, 1-148.	2.8	16
20	Early post-traumatic acute respiratory distress syndrome and albumin excretion rate: a prospective evaluation of a "point-of-care" predictive test. <i>Injury</i> , 2001, 32, 177-181.	1.7	15
21	Ankle Injury Management (AIM): design of a pragmatic multi-centre equivalence randomised controlled trial comparing Close Contact Casting (CCC) to Open surgical Reduction and Internal Fixation (ORIF) in the treatment of unstable ankle fractures in patients over 60 years. <i>BMC Musculoskeletal Disorders</i> , 2014, 15, 79.	1.9	14
22	The stove-in chest: a complex flail chest injury. <i>Injury</i> , 2004, 35, 490-493.	1.7	12
23	Pilot study assessing functional outcome of tibial pilon fractures using the VSTORM method. <i>Injury</i> , 2013, 44, 1112-1116.	1.7	12
24	Indirect reduction using a simple quadrilateral frame in the application of distal tibial LCP technical tips. <i>Injury</i> , 2005, 36, 1138-1142.	1.7	11
25	Chemiluminescence: Comparison of Whole Blood with Isolated Polymorphonuclear Leukocytes after Major Trauma. <i>Journal of Trauma</i> , 2004, 57, 347-351.	2.3	10
26	Catastrophic myonecrosis following circumferential pelvic binding after massive crush injury: A case report. <i>Injury Extra</i> , 2009, 40, 84-86.	0.2	10
27	Finite element analysis modelling of proximal femoral fractures, including post-fixation periprosthetic fractures. <i>Injury</i> , 2013, 44, 791-795.	1.7	10
28	The use of personalised patient information leaflets to improve patients' perceived understanding following open fractures. <i>European Journal of Orthopaedic Surgery and Traumatology</i> , 2019, 29, 537-543.	1.4	10
29	Arthrodesis for osteoarthritis of the manubriosternal joint. <i>European Journal of Cardio-thoracic Surgery</i> , 2006, 29, 119-121.	1.4	9
30	High-Fidelity Simulation Model of Pelvic Hemorrhagic Trauma: The Future for Military Surgical Skills Training?. <i>Military Medicine</i> , 2016, 181, 1407-1409.	0.8	9
31	Intra-pelvic pressure changes after pelvic fracture: A cadaveric study quantifying the effect of a pelvic binder and limb bandaging over a bolster. <i>Injury</i> , 2017, 48, 833-840.	1.7	9
32	Current concepts of the inflammatory response after major trauma: an update. <i>Injury</i> , 2005, 36, 227-229.	1.7	8
33	Finite element analysis in trauma & orthopaedics – an introduction to clinically relevant simulation & its limitations. <i>Orthopaedics and Trauma</i> , 2012, 26, 410-416.	0.4	7
34	The development of a novel patient-derived recovery scale for open tibial fractures. <i>Bone and Joint Journal</i> , 2020, 102-B, 17-25.	4.4	7
35	Is physiotherapy effective for patients with a chronic mid-body Achilles tendinopathy? A systematic review of non-surgical and non-pharmacological interventions. <i>International Musculoskeletal Medicine</i> , 2011, 33, 152-160.	0.1	5
36	A Comparison of Improvised and Commercially Available Point-of-Wounding Tourniquets in Simulated Traumatic Amputation with Catastrophic Hemorrhage. <i>Military Medicine</i> , 2020, 185, e1536-e1541.	0.8	4

#	ARTICLE	IF	CITATIONS
37	(iii) An update on the systemic response to trauma. Orthopaedics and Trauma, 2010, 24, 24-28.	0.4	3
38	Simulation education for lower limb fasciotomy: improving surgical trainee confidence in executing a time-critical limb-saving procedure. BMJ Simulation and Technology Enhanced Learning, 2019, 5, 238-239.	0.7	3
39	A novel method to correctly place the fasciotomy incision for decompression of the anterior and peroneal compartments of the leg. Injury, 2016, 47, 962-968.	1.7	2
40	Modulation of Interleukin-8-Mediated Neutrophil Migration Following Major Lower-Limb Fracture and Operative Stabilization. European Journal of Trauma and Emergency Surgery, 2005, 31, 575-582.	0.3	1
41	Authors' Reply to the Response to Our Article (A Comparison of Improvised and Commercially) Tj ETQq1 1 0.784314 rgBT /Over	0.8	1
42	Surgical simulation training for escharotomy: A novel course, improving candidate's confidence in a time critical procedure. Burns, 2023, 49, 783-787.	1.9	1
43	Trauma assessment and follow up with integration of modern communication devices (TRAFIC). A pilot study. Injury, 2018, 49, 2013-2017.	1.7	0
44	Measuring recovery after open lower limb fractures: combined objective functional tests and global perceived recovery outperform narrower metrics and a standard generic patient reported outcome measure. BMC Musculoskeletal Disorders, 2021, 22, 539.	1.9	0
45	Damage Control Orthopaedics in the Low Resource Environment. , 2016, , 315-323.		0