

Iordanis N Papadopoulos

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2691696/publications.pdf>

Version: 2024-02-01

41
papers

864
citations

394421

19
h-index

477307

29
g-index

41
all docs

41
docs citations

41
times ranked

1033
citing authors

#	ARTICLE	IF	CITATIONS
1	Auditing 655 Fatalities with Pelvic Fractures by Autopsy as a Basis to Evaluate Trauma Care. <i>Journal of the American College of Surgeons</i> , 2006, 203, 30-43.	0.5	88
2	Early Diagnosis and Surgical Treatment for Necrotizing Fasciitis: A Multicenter Study. <i>Frontiers in Surgery</i> , 2017, 4, 5.	1.4	61
3	Clinical significance of kallikrein-related peptidase (KLK10) mRNA expression in colorectal cancer. <i>Clinical Biochemistry</i> , 2013, 46, 1453-1461.	1.9	48
4	High miR-96 levels in colorectal adenocarcinoma predict poor prognosis, particularly in patients without distant metastasis at the time of initial diagnosis. <i>Tumor Biology</i> , 2016, 37, 11815-11824.	1.8	44
5	Quantitative expression analysis and prognostic significance of the novel apoptosis-related gene <i>BCL2L12</i> in colon cancer. <i>Biological Chemistry</i> , 2008, 389, 1467-1475.	2.5	40
6	miR-15a-5p, A Novel Prognostic Biomarker, Predicting Recurrent Colorectal Adenocarcinoma. <i>Molecular Diagnosis and Therapy</i> , 2017, 21, 453-464.	3.8	40
7	Enhanced miR-182 transcription is a predictor of poor overall survival in colorectal adenocarcinoma patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2014, 52, 1217-27.	2.3	39
8	miR-224 overexpression is a strong and independent prognosticator of short-term relapse and poor overall survival in colorectal adenocarcinoma. <i>International Journal of Oncology</i> , 2015, 46, 849-859.	3.3	38
9	Molecular analysis and prognostic impact of the novel apoptotic gene <i>BCL2L12</i> in gastric cancer. <i>Biochemical and Biophysical Research Communications</i> , 2010, 391, 214-218.	2.1	34
10	Kallikrein-related peptidase-6 (KLK6) mRNA expression is an independent prognostic tissue biomarker of poor disease-free and overall survival in colorectal adenocarcinoma. <i>Tumor Biology</i> , 2014, 35, 4673-4685.	1.8	34
11	Upregulated miR-16 expression is an independent indicator of relapse and poor overall survival of colorectal adenocarcinoma patients. <i>Clinical Chemistry and Laboratory Medicine</i> , 2017, 55, 737-747.	2.3	33
12	Elevated expression of miR-24-3p is a potentially adverse prognostic factor in colorectal adenocarcinoma. <i>Clinical Biochemistry</i> , 2017, 50, 285-292.	1.9	31
13	Kallikrein-related peptidase 4 (KLK4) mRNA predicts short-term relapse in colorectal adenocarcinoma patients. <i>Cancer Letters</i> , 2013, 330, 106-112.	7.2	30
14	<i>KLK11</i> mRNA expression predicts poor disease-free and overall survival in colorectal adenocarcinoma patients. <i>Biomarkers in Medicine</i> , 2014, 8, 671-685.	1.4	30
15	High microRNA-28-5p expression in colorectal adenocarcinoma predicts short-term relapse of node-negative patients and poor overall survival of patients with non-metastatic disease. <i>Clinical Chemistry and Laboratory Medicine</i> , 2018, 56, 990-1000.	2.3	28
16	Kallikrein-related peptidase 13 (KLK13) gene expressional status contributes significantly in the prognosis of primary gastric carcinomas. <i>Clinical Biochemistry</i> , 2010, 43, 1205-1211.	1.9	27
17	miR-34a overexpression predicts poor prognostic outcome in colorectal adenocarcinoma, independently of clinicopathological factors with established prognostic value. <i>Clinical Biochemistry</i> , 2017, 50, 918-924.	1.9	25
18	The effect of metronomic versus standard chemotherapy on the regulatory to effector T-cell equilibrium in cancer patients. <i>Experimental Hematology and Oncology</i> , 2014, 3, 3.	5.0	23

#	ARTICLE	IF	CITATIONS
19	High clusterin (CLU) mRNA expression levels in tumors of colorectal cancer patients predict a poor prognostic outcome. <i>Clinical Biochemistry</i> , 2020, 75, 62-69.	1.9	23
20	TYMS/KRAS/BRAF molecular profiling predicts survival following adjuvant chemotherapy in colorectal cancer. <i>World Journal of Gastrointestinal Oncology</i> , 2019, 11, 551-566.	2.0	20
21	Osteoporosis-pseudoglioma syndrome: clinical, genetic, and treatment-response study of 10 new cases in Greece. <i>European Journal of Pediatrics</i> , 2019, 178, 323-329.	2.7	17
22	Heat shock protein beta 3 (HSPB3) is an unfavorable molecular biomarker in colorectal adenocarcinoma. <i>Molecular Carcinogenesis</i> , 2020, 59, 116-125.	2.7	17
23	Status and perspectives of hospital mortality in a public urban Hellenic hospital, based on a five-year review. <i>BMC Public Health</i> , 2008, 8, 28.	2.9	15
24	A structured autopsy-based audit of 370 firearm fatalities: Contribution to inform policy decisions and the probability of the injured arriving alive at a hospital and receiving definitive care. <i>Accident Analysis and Prevention</i> , 2013, 50, 667-677.	5.7	13
25	High Expression of a tRNAPro Derivative Associates with Poor Survival and Independently Predicts Colorectal Cancer Recurrence. <i>Biomedicines</i> , 2022, 10, 1120.	3.2	12
26	Alcohol and psychoactive drugs increased the pre-hospital mortality in 655 fall-related fatalities in Greece: A call for management protocols. <i>Injury</i> , 2012, 43, 1522-1526.	1.7	9
27	Asymptomatic omental granuloma following spillage of gallstones during laparoscopic cholecystectomy protects patients and influences surgeons' decisions: a review. <i>BMJ Case Reports</i> , 2012, 2012, bcr1020114980-bcr1020114980.	0.5	8
28	l-Dopa decarboxylase (DDC) constitutes an emerging biomarker in predicting patients' survival with stomach adenocarcinomas. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013, 139, 297-306.	2.5	8
29	Motor vehicle collision fatalities involving alcohol and illicit drugs in Greece: the need for management protocols and a reassessment of surveillance. <i>Addiction</i> , 2010, 105, 1952-1961.	3.3	6
30	Patients with pelvic fractures due to falls: A paradigm that contributed to autopsy-based audit of trauma in Greece. <i>Journal of Trauma Management and Outcomes</i> , 2011, 5, 2.	0.9	6
31	Primary epithelial splenic cyst with micro-rupture and raised carbohydrate antigen CA 19-9: a proposal for management. <i>BMJ Case Reports</i> , 2010, 2010, bcr0620103125-bcr0620103125.	0.5	4
32	Primary epithelial splenic cyst with micro-rupture and raised carbohydrate antigen CA 19-9: a paradigm of management. <i>BMJ Case Reports</i> , 2010, 2010, bcr0120102636-bcr0120102636.	0.5	4
33	Perforated mucinous cystadenoma of the vermiform appendix: an overview in reasoning clinical decisions. <i>BMJ Case Reports</i> , 2011, 2011, bcr0220113902-bcr0220113902.	0.5	3
34	Delayed reaction to the Dacron buttress used in Stamey bladder neck suspension. <i>International Urology and Nephrology</i> , 2006, 38, 269-272.	1.4	2
35	Fibroepithelial polyp in an anal fistulous track: a sign of chronic pathology. <i>BMJ Case Reports</i> , 2010, 2010, bcr0820092169-bcr0820092169.	0.5	2
36	Enterocolitis with multiple ulcers of ileum and right colon in a patient with leukaemia attributed to cytomegalovirus. <i>BMJ Case Reports</i> , 2012, 2012, bcr0120125651-bcr0120125651.	0.5	1

#	ARTICLE	IF	CITATIONS
37	The long-term need for calcium supplementation after incidental parathyroidectomy. Journal of Taibah University Medical Sciences, 2022, 17, 214-219.	0.9	1
38	Acute abdominal pain due to intussusception of the small intestine in a 17-year-old patient. Surgery, 2014, 156, 1282-1283.	1.9	0
39	Femoral fractures are an indicator of increased severity of injury for road traffic collision victims: an autopsy-based case-control study on 4895 fatalities. Archives of Orthopaedic and Trauma Surgery, 2021, , 1.	2.4	0
40	Asymptomatic bladder neurofibroma in neurofibromatosis type I. Minerva Pediatrics, 2020, , .	0.4	0
41	Incidence, risk factors and potential timing of occurrence of pulmonary embolism in fatal trauma: An autopsy based case-control study on 2705 victims. Journal of the Royal College of Surgeons of Edinburgh, 2022, , .	1.8	0