

# Nishat Passricha

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2726899/publications.pdf>

Version: 2024-02-01

168  
papers

21,279  
citations

36203

51  
h-index

9839

141  
g-index

173  
all docs

173  
docs citations

173  
times ranked

19933  
citing authors

#	ARTICLE	IF	CITATIONS
1	Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. <i>Plant Physiology and Biochemistry</i> , 2010, 48, 909-930.	2.8	8,238
2	Cold, salinity and drought stresses: An overview. <i>Archives of Biochemistry and Biophysics</i> , 2005, 444, 139-158.	1.4	2,295
3	Biofertilizers function as key player in sustainable agriculture by improving soil fertility, plant tolerance and crop productivity. <i>Microbial Cell Factories</i> , 2014, 13, 66.	1.9	747
4	Polyamines and abiotic stress tolerance in plants. <i>Plant Signaling and Behavior</i> , 2010, 5, 26-33.	1.2	606
5	Mechanisms of High Salinity Tolerance in Plants. <i>Methods in Enzymology</i> , 2007, 428, 419-438.	0.4	585
6	Signaling through MAP kinase networks in plants. <i>Archives of Biochemistry and Biophysics</i> , 2006, 452, 55-68.	1.4	331
7	Cadmium stress tolerance in crop plants. <i>Plant Signaling and Behavior</i> , 2011, 6, 215-222.	1.2	311
8	Calcium Signaling Network in Plants. <i>Plant Signaling and Behavior</i> , 2007, 2, 79-85.	1.2	310
9	Cadmium at high dose perturbs growth, photosynthesis and nitrogen metabolism while at low dose it up regulates sulfur assimilation and antioxidant machinery in garden cress ( <i>Lepidium sativum</i> L.). <i>Plant Science</i> , 2012, 182, 112-120.	1.7	293
10	Superoxide dismutaseâ€™ mentor of abiotic stress tolerance in crop plants. <i>Environmental Science and Pollution Research</i> , 2015, 22, 10375-10394.	2.7	247
11	Molecular Mechanisms of DNA Damage and Repair: Progress in Plants. <i>Critical Reviews in Biochemistry and Molecular Biology</i> , 2001, 36, 337-397.	2.3	238
12	Importance of nitric oxide in cadmium stress tolerance in crop plants. <i>Plant Physiology and Biochemistry</i> , 2013, 63, 254-261.	2.8	228
13	Pea DNA helicase 45 overexpression in tobacco confers high salinity tolerance without affecting yield. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 509-514.	3.3	216
14	Antioxidant enzyme activities in maize plants colonized with <i>Piriformospora indica</i> . <i>Microbiology (United Kingdom)</i> , 2009, 155, 780-790.	0.7	214
15	The CRISPR/Cas Genome-Editing Tool: Application in Improvement of Crops. <i>Frontiers in Plant Science</i> , 2016, 7, 506.	1.7	196
16	Genotoxic stress in plants: Shedding light on DNA damage, repair and DNA repair helicases. <i>Mutation Research - Reviews in Mutation Research</i> , 2009, 681, 134-149.	2.4	183
17	Unraveling DNA helicases. Motif, structure, mechanism and function. <i>FEBS Journal</i> , 2004, 271, 1849-1863.	0.2	172
18	Nucleolin: A Multifunctional Major Nucleolar Phosphoprotein. <i>Critical Reviews in Biochemistry and Molecular Biology</i> , 1998, 33, 407-436.	2.3	166

#	ARTICLE	IF	CITATIONS
19	Signaling through G protein coupled receptors. <i>Plant Signaling and Behavior</i> , 2009, 4, 942-947.	1.2	165
20	Ku Autoantigen: A Multifunctional DNA-Binding Protein. <i>Critical Reviews in Biochemistry and Molecular Biology</i> , 2000, 35, 1-33.	2.3	164
21	Dose-dependent response of <i>Trichoderma harzianum</i> in improving drought tolerance in rice genotypes. <i>Planta</i> , 2016, 243, 1251-1264.	1.6	146
22	ATP-sulfurylase, sulfur-compounds, and plant stress tolerance. <i>Frontiers in Plant Science</i> , 2015, 6, 210.	1.7	145
23	Prokaryotic and eukaryotic DNA helicases. Essential molecular motor proteins for cellular machinery. <i>FEBS Journal</i> , 2004, 271, 1835-1848.	0.2	139
24	Reactive Oxygen Species Generation-Scavenging and Signaling during Plant-Arbuscular Mycorrhizal and <i>Piriformospora indica</i> Interaction under Stress Condition. <i>Frontiers in Plant Science</i> , 2016, 7, 1574.	1.7	133
25	Knights in Action: Lectin Receptor-Like Kinases in Plant Development and Stress Responses. <i>Molecular Plant</i> , 2013, 6, 1405-1418.	3.9	132
26	<sc>O</sc>s<sc>SUV</sc>3 dual helicase functions in salinity stress tolerance by maintaining photosynthesis and antioxidant machinery in rice (<i>Oryza sativa</i> cv.) Tj ETQq0 0 0 rgBT /Overlook 10 Tf 50 457 Td	1.2	131
27	Genome-wide analysis of lectin receptor-like kinase family from <i>Arabidopsis</i> and rice. <i>Plant Molecular Biology</i> , 2012, 80, 365-388.	2.0	129
28	Recent advances in development of marker-free transgenic plants: Regulation and biosafety concern. <i>Journal of Biosciences</i> , 2012, 37, 167-197.	0.5	128
29	Stress responsive DEAD-box helicases: A new pathway to engineer plant stress tolerance. <i>Journal of Photochemistry and Photobiology B: Biology</i> , 2006, 84, 150-160.	1.7	126
30	Differential cadmium stress tolerance in five Indian mustard (<i>Brassica juncea</i>L.) cultivars. <i>Plant Signaling and Behavior</i> , 2011, 6, 293-300.	1.2	124
31	Heterotrimeric G-protein complex and G-protein-coupled receptor from a legume ( <i>Pisum sativum</i> ): role in salinity and heat stress and cross-talk with phospholipaseâ€fC. <i>Plant Journal</i> , 2007, 51, 656-669.	2.8	122
32	Metal/metalloid stress tolerance in plants: role of ascorbate, its redox couple, and associated enzymes. <i>Protoplasma</i> , 2014, 251, 1265-1283.	1.0	121
33	Cold- and salinity stress-induced bipolar pea DNA helicase 47 is involved in protein synthesis and stimulated by phosphorylation with protein kinase C. <i>Plant Journal</i> , 2005, 44, 76-87.	2.8	107
34	<i>Os<sc>ACA</sc>6</i>, a Pa€type <sc>ILB</sc> Ca<sup>2+</sup> <sc>ATP</sc>ase promotes salinity and drought stress tolerance in tobacco by <sc>ROS</sc> scavenging and enhancing the expression of stressâ€responsive genes. <i>Plant Journal</i> , 2013, 76, 997-1015.	2.8	97
35	Fungal association and utilization of phosphate by plants: success, limitations, and future prospects. <i>Frontiers in Microbiology</i> , 2015, 6, 984.	1.5	96
36	A new DEAD-box helicase ATP-binding protein (OsABP) from rice is responsive to abiotic stress. <i>Plant Signaling and Behavior</i> , 2012, 7, 1138-1143.	1.2	95

#	ARTICLE	IF	CITATIONS
37	Plant signaling in stress. <i>Plant Signaling and Behavior</i> , 2008, 3, 79-86.	1.2	86
38	Genome-wide analysis of helicase gene family from rice and Arabidopsis: a comparison with yeast and human. <i>Plant Molecular Biology</i> , 2010, 73, 449-465.	2.0	86
39	A DNA helicase from <i>Pisum sativum</i> is homologous to translation initiation factor and stimulates topoisomerase I activity. <i>Plant Journal</i> , 2000, 24, 219-229.	2.8	82
40	A DESD-box helicase functions in salinity stress tolerance by improving photosynthesis and antioxidant machinery in rice ( <i>Oryza sativa</i> L. cv. PB1). <i>Plant Molecular Biology</i> , 2013, 82, 1-22.	2.0	79
41	Overexpression of Pea DNA Helicase 45 (PDH45) imparts tolerance to multiple abiotic stresses in chili ( <i>Capsicum annum</i> L.). <i>Scientific Reports</i> , 2017, 7, 2760.	1.6	77
42	A single subunit MCM6 from pea promotes salinity stress tolerance without affecting yield. <i>Plant Molecular Biology</i> , 2011, 76, 19-34.	2.0	75
43	Plant MCM proteins: role in DNA replication and beyond. <i>Plant Molecular Biology</i> , 2011, 77, 537-545.	2.0	73
44	Chaperones and foldases in endoplasmic reticulum stress signaling in plants. <i>Plant Signaling and Behavior</i> , 2011, 6, 232-236.	1.2	73
45	Phenotypic and molecular characterization of native <i>Azospirillum</i> strains from rice fields to improve crop productivity. <i>Protoplasma</i> , 2014, 251, 943-953.	1.0	66
46	Pea p68, a DEAD-Box Helicase, Provides Salinity Stress Tolerance in Transgenic Tobacco by Reducing Oxidative Stress and Improving Photosynthesis Machinery. <i>PLoS ONE</i> , 2014, 9, e98287.	1.1	65
47	Over-expression of a DEAD-box helicase, PDH45, confers both seedling and reproductive stage salinity tolerance to rice ( <i>Oryza sativa</i> L.). <i>Molecular Breeding</i> , 2012, 30, 345-354.	1.0	61
48	Unraveling DNA Repair in Human: Molecular Mechanisms and Consequences of Repair Defect. <i>Critical Reviews in Biochemistry and Molecular Biology</i> , 2001, 36, 261-290.	2.3	60
49	NPKS uptake, sensing, and signaling and miRNAs in plant nutrient stress. <i>Protoplasma</i> , 2016, 253, 767-786.	1.0	59
50	Pea lectin receptor-like kinase functions in salinity adaptation without yield penalty, by alleviating osmotic and ionic stresses and upregulating stress-responsive genes. <i>Plant Molecular Biology</i> , 2015, 88, 193-206.	2.0	58
51	Rice heterotrimeric G-protein gamma subunits (RGG1 and RGG2) are differentially regulated under abiotic stress. <i>Plant Signaling and Behavior</i> , 2012, 7, 733-740.	1.2	57
52	Potassium: A key modulator for cell homeostasis. <i>Journal of Biotechnology</i> , 2020, 324, 198-210.	1.9	57
53	Genome-wide analysis of glutathione reductase (GR) genes from rice and Arabidopsis. <i>Plant Signaling and Behavior</i> , 2013, 8, e23021.	1.2	54
54	High frequency regeneration via direct somatic embryogenesis and efficient <i>Agrobacterium</i> -mediated genetic transformation of tobacco. <i>Plant Signaling and Behavior</i> , 2013, 8, e24354.	1.2	54

#	ARTICLE	IF	CITATIONS
55	Pea lectin receptor-like kinase promotes high salinity stress tolerance in bacteria and expresses in response to stress in planta. <i>Glycoconjugate Journal</i> , 2010, 27, 133-150.	1.4	51
56	Function of heterotrimeric G-protein $\beta$ subunit RGG1 in providing salinity stress tolerance in rice by elevating detoxification of ROS. <i>Planta</i> , 2017, 245, 367-383.	1.6	51
57	Plant DNA helicases: the long unwinding road. <i>Journal of Experimental Botany</i> , 2003, 54, 2201-2214.	2.4	49
58	Development of Agrobacterium-mediated transformation technology for mature seed-derived callus tissues of indica rice cultivar IR64. <i>GM Crops and Food</i> , 2012, 3, 123-128.	2.0	49
59	Genome-wide analysis of plant-type II Ca <sup>2+</sup> -ATPases gene family from rice and <i>Arabidopsis</i> : Potential role in abiotic stresses. <i>Plant Physiology and Biochemistry</i> , 2013, 65, 32-47.	2.8	49
60	Genome wide analysis of Cyclophilin gene family from rice and <i>Arabidopsis</i> and its comparison with yeast. <i>Plant Signaling and Behavior</i> , 2012, 7, 1653-1666.	1.2	48
61	PDH45 transgenic rice maintain cell viability through lower accumulation of Na <sup>+</sup> , ROS and calcium homeostasis in roots under salinity stress. <i>Journal of Plant Physiology</i> , 2016, 191, 1-11.	1.6	46
62	Introduction of Pea DNA Helicase 45 into Sugarcane ( <i>Saccharum</i> spp. Hybrid) Enhances Cell Membrane Thermostability and Upregulation of Stress-Responsive Genes Leads to Abiotic Stress Tolerance. <i>Molecular Biotechnology</i> , 2015, 57, 475-488.	1.3	45
63	Insights into the functional characteristics of geminivirus rolling-circle replication initiator protein and its interaction with host factors affecting viral DNA replication. <i>Archives of Virology</i> , 2015, 160, 375-387.	0.9	42
64	Simultaneous expression of regulatory genes associated with specific drought adaptive traits improves drought adaptation in peanut. <i>Plant Biotechnology Journal</i> , 2016, 14, 1008-1020.	4.1	42
65	Purification and Characterization of a DNA Helicase from Pea Chloroplast that Translocates in the 3'-to-5' Direction. <i>FEBS Journal</i> , 1996, 238, 54-63.	0.2	41
66	A novel <i>Azotobacter vinelandii</i> (SRI Az <sup>3</sup> ) functions in salinity stress tolerance in rice. <i>Plant Signaling and Behavior</i> , 2014, 9, e29377.	1.2	41
67	Overexpression of a Pea DNA Helicase (PDH45) in Peanut ( <i>Arachis hypogaea</i> L.) Confers Improvement of Cellular Level Tolerance and Productivity Under Drought Stress. <i>Molecular Biotechnology</i> , 2014, 56, 111-125.	1.3	41
68	Pea DNA helicase 45 promotes salinity stress tolerance in IR64 rice with improved yield. <i>Plant Signaling and Behavior</i> , 2012, 7, 1042-1046.	1.2	40
69	Human DNA helicase V, a novel DNA unwinding enzyme from HeLa cells. <i>Nucleic Acids Research</i> , 1993, 21, 2323-2329.	6.5	39
70	Multiple abiotic stress responsive rice cyclophilin: (OsCYP-25) mediates a wide range of cellular responses. <i>Communicative and Integrative Biology</i> , 2013, 6, e25260.	0.6	38
71	Field performance of bacterial inoculants to alleviate water stress effects in wheat ( <i>Triticum</i> ) Tj ETQq1 1 0.784314 rrgBT /Overlock 10 Tff	1.8	38
72	microRNAs as promising tools for improving stress tolerance in rice. <i>Plant Signaling and Behavior</i> , 2012, 7, 1296-1301.	1.2	36

#	ARTICLE	IF	CITATIONS
73	A pea homologue of human DNA helicase I is localized within the dense fibrillar component of the nucleolus and stimulated by phosphorylation with CK2 and cdc2 protein kinases. <i>Plant Journal</i> , 2001, 25, 9-17.	2.8	36
74	OsSUV3 transgenic rice maintains higher endogenous levels of plant hormones that mitigates adverse effects of salinity and sustains crop productivity. <i>Rice</i> , 2014, 7, 17.	1.7	35
75	Genome-wide analysis and transcriptional expression pattern-assessment of superoxide dismutase (SOD) in rice and Arabidopsis under abiotic stresses. <i>Plant Gene</i> , 2019, 17, 100165.	1.4	34
76	OsACA6, a P-type 2B Ca <sup>2+</sup> ATPase functions in cadmium stress tolerance in tobacco by reducing the oxidative stress load. <i>Planta</i> , 2014, 240, 809-824.	1.6	33
77	Assessing zygosity in progeny of transgenic plants: current methods and perspectives. <i>Journal of Biological Methods</i> , 2016, 3, e46.	1.0	32
78	Cloning and characterisation of a gene encoding an antiviral protein from <i>Clerodendrum aculeatum</i> L. <i>Plant Molecular Biology</i> , 1997, 33, 745-751.	2.0	31
79	Further Characterization of Calcineurin B-Like Protein and Its Interacting Partner CBL-Interacting Protein Kinase from <i>Pisum sativum</i> . <i>Plant Signaling and Behavior</i> , 2007, 2, 358-361.	1.2	31
80	Rice heterotrimeric G-protein alpha subunit (RGA1): In silico analysis of the gene and promoter and its upregulation under abiotic stress. <i>Plant Physiology and Biochemistry</i> , 2013, 63, 262-271.	2.8	31
81	Different expression of miRNAs targeting helicases in rice in response to low and high dose rate <sup>137</sup> Cs treatments. <i>Plant Signaling and Behavior</i> , 2013, 8, e25128.	1.2	30
82	Salt stress triggers augmented levels of Na <sup>+</sup> , Ca <sup>2+</sup> and ROS and alter stress-responsive gene expression in roots of CBL9 and CIPK23 knockout mutants of <i>Arabidopsis thaliana</i> . <i>Environmental and Experimental Botany</i> , 2019, 161, 265-276.	2.0	30
83	Forisomes: calcium-powered protein complexes with potential as smart biomaterials. <i>Trends in Biotechnology</i> , 2010, 28, 102-110.	4.9	27
84	Helicases as molecular motors: An insight. <i>Physica A: Statistical Mechanics and Its Applications</i> , 2006, 372, 70-83.	1.2	24
85	A single subunit MCM6 from pea forms homohexamer and functions as DNA helicase. <i>Plant Molecular Biology</i> , 2010, 74, 327-336.	2.0	23
86	<i>Pisum sativum</i> p68 DEAD-box protein is ATP-dependent RNA helicase and unique bipolar DNA helicase. <i>Plant Molecular Biology</i> , 2014, 85, 639-651.	2.0	23
87	Rice lectin receptor-like kinase provides salinity tolerance by ion homeostasis. <i>Biotechnology and Bioengineering</i> , 2020, 117, 498-510.	1.7	23
88	Stress induced beta subunit of heterotrimeric G-proteins from <i>Pisum sativum</i> interacts with mitogen activated protein kinase. <i>Plant Signaling and Behavior</i> , 2011, 6, 287-292.	1.2	22
89	Emerging Importance of Helicases in Plant Stress Tolerance: Characterization of <i>Oryza sativa</i> Repair Helicase XPB2 Promoter and Its Functional Validation in Tobacco under Multiple Stresses. <i>Frontiers in Plant Science</i> , 2015, 6, 1094.	1.7	22
90	Pea p68 Imparts Salinity Stress Tolerance in Rice by Scavenging of ROS-Mediated H <sub>2</sub> O <sub>2</sub> and Interacts with Argonaute. <i>Plant Molecular Biology Reporter</i> , 2015, 33, 221-238.	1.0	21

#	ARTICLE	IF	CITATIONS
91	Unraveling DNA helicases from plant cells. , 1997, 33, 947-952.		20
92	Cloning and functional characterization of the promoter of <i>P</i> sSEOF1 gene from <i>Pisum sativum</i> under different stress conditions using <i>Agrobacterium</i> -mediated transient assay. <i>Plant Signaling and Behavior</i> , 2014, 9, e29626.	1.2	20
93	Pea DNA Topoisomerase I Is Phosphorylated and Stimulated by Casein Kinase 2 and Protein Kinase C. <i>Plant Physiology</i> , 2003, 132, 2108-2115.	2.3	19
94	Genetic engineering of crops: a ray of hope for enhanced food security. <i>Plant Signaling and Behavior</i> , 2014, 9, e28545.	1.2	19
95	Prediction and validation of cis-regulatory elements in 5' upstream regulatory regions of lectin receptor-like kinase gene family in rice. <i>Protoplasma</i> , 2017, 254, 669-684.	1.0	19
96	Rice G-protein coupled receptor (GPCR). <i>Plant Signaling and Behavior</i> , 2011, 6, 1079-1086.	1.2	18
97	An efficient and rapid regeneration via multiple shoot induction from mature seed derived embryogenic and organogenic callus of Indian maize ( <i>Zea mays</i> L.). <i>Plant Signaling and Behavior</i> , 2013, 8, e25891.	1.2	18
98	High-frequency regeneration via multiple shoot induction of an elite recalcitrant cotton ( <i>Gossypium hirsutum</i> L. cv Narashima) by using embryo apex. <i>Plant Signaling and Behavior</i> , 2013, 8, e22763.	1.2	18
99	Reproductive Organ and Vascular Specific Promoter of the Rice Plasma Membrane Ca <sup>2+</sup> ATPase Mediates Environmental Stress Responses in Plants. <i>PLoS ONE</i> , 2013, 8, e57803.	1.1	18
100	Marker-free transgenic rice plant overexpressing pea LecRLK imparts salinity tolerance by inhibiting sodium accumulation. <i>Plant Molecular Biology</i> , 2019, 99, 265-281.	2.0	18
101	Translation initiation factor 4A: a prototype member of dead-box protein family. <i>Physiology and Molecular Biology of Plants</i> , 2008, 14, 101-107.	1.4	17
102	In-silico analysis and expression profiling implicate diverse role of EPSPS family genes in regulating developmental and metabolic processes. <i>BMC Research Notes</i> , 2014, 7, 58.	0.6	17
103	<i>PDH45</i> overexpressing transgenic tobacco and rice plants provide salinity stress tolerance via less sodium accumulation. <i>Plant Signaling and Behavior</i> , 2015, 10, e992289.	1.2	17
104	Cold stress-induced pea DNA helicase 47 is homologous to eIF4A and inhibited by DNA-interacting ligands. <i>Archives of Biochemistry and Biophysics</i> , 2005, 440, 79-90.	1.4	16
105	Isolation of high salinity stress tolerant genes from <i>Pisum sativum</i> by random overexpression in <i>Escherichia coli</i> and their functional validation. <i>Plant Signaling and Behavior</i> , 2009, 4, 400-412.	1.2	16
106	Promoter of a salinity and cold stress-induced MCM6 DNA helicase from pea. <i>Plant Signaling and Behavior</i> , 2011, 6, 1006-1008.	1.2	16
107	Isolation and functional characterization of the promoter of a DEAD-box helicase <i>Psp68</i> using <i>Agrobacterium</i> -mediated transient assay. <i>Plant Signaling and Behavior</i> , 2014, 9, e28992.	1.2	16
108	Salinity and drought tolerant OsACA6 enhances cold tolerance in transgenic tobacco by interacting with stress-inducible proteins. <i>Plant Physiology and Biochemistry</i> , 2014, 82, 229-238.	2.8	16

#	ARTICLE	IF	CITATIONS
109	Potent inhibition of DNA unwinding and ATPase activities of pea DNA helicase 45 by DNA-binding agents. <i>Biochemical and Biophysical Research Communications</i> , 2002, 294, 334-339.	1.0	15
110	First evidence of ethylene production by <i>Fusarium mangiferae</i> associated with mango malformation. <i>Plant Signaling and Behavior</i> , 2013, 8, e22673.	1.2	15
111	Effect of salinity tolerant PDH45 transgenic rice on physicochemical properties, enzymatic activities and microbial communities of rhizosphere soils. <i>Plant Signaling and Behavior</i> , 2013, 8, e24950.	1.2	15
112	Isolation, in silico characterization, localization and expression analysis of abiotic stress-responsive rice G-protein $\beta^2$ subunit (RGB1). <i>Plant Signaling and Behavior</i> , 2014, 9, e28890.	1.2	15
113	Salt tolerant SUV3 overexpressing transgenic rice plants conserve physicochemical properties and microbial communities of rhizosphere. <i>Chemosphere</i> , 2015, 119, 1040-1047.	4.2	15
114	Concurrent overexpression of rice G-protein $\beta^2$ and $\beta^3$ subunits provide enhanced tolerance to sheath blight disease and abiotic stress in rice. <i>Planta</i> , 2019, 250, 1505-1520.	1.6	15
115	Silencing of tomato CTR1 provides enhanced tolerance against Tomato leaf curl virus infection. <i>Plant Signaling and Behavior</i> , 2019, 14, e1565595.	1.2	15
116	Acclimation potential of Noni ( <i>Morinda citrifolia</i> L.) plant to temperature stress is mediated through photosynthetic electron transport rate. <i>Plant Signaling and Behavior</i> , 2021, 16, 1865687.	1.2	15
117	The Gly-Arg-rich C-terminal domain of pea nucleolin is a DNA helicase that catalytically translocates in the 5' to 3' direction. <i>Archives of Biochemistry and Biophysics</i> , 2005, 434, 306-315.	1.4	14
118	Wide range of interacting partners of pea $G\beta^2$ subunit of G-proteins suggests its multiple functions in cell signalling. <i>Plant Physiology and Biochemistry</i> , 2012, 58, 1-5.	2.8	14
119	Sequence-specific <sup>1</sup> H, <sup>13</sup> C and <sup>15</sup> N NMR assignments of Cyclophilin A like protein from <i>Piriformospora indica</i> involved in salt stress tolerance. <i>Biomolecular NMR Assignments</i> , 2013, 7, 175-178.	0.4	14
120	Isolation of genes conferring salt tolerance from <i>Piriformospora indica</i> by random overexpression in <i>Escherichia coli</i> . <i>World Journal of Microbiology and Biotechnology</i> , 2015, 31, 1195-1209.	1.7	14
121	Mango ( <i>Mangifera indica</i> L.) malformation: a malady of stress ethylene origin. <i>Physiology and Molecular Biology of Plants</i> , 2015, 21, 1-8.	1.4	14
122	Heterologous expression and biochemical characterization of a highly active and stable chloroplastic CuZn-superoxide dismutase from <i>Pisum sativum</i> . <i>BMC Biotechnology</i> , 2015, 15, 3.	1.7	14
123	Heterologous expression of PDH47 confers drought tolerance in indica rice. <i>Plant Cell, Tissue and Organ Culture</i> , 2017, 130, 577-589.	1.2	14
124	Overexpression of a pea DNA helicase 45 in bacteria confers salinity stress tolerance. <i>Plant Signaling and Behavior</i> , 2011, 6, 1271-1275.	1.2	13
125	Molecular characterization of cyclophilin A-like protein from <i>Piriformospora indica</i> for its potential role to abiotic stress tolerance in <i>E. coli</i> . <i>BMC Research Notes</i> , 2013, 6, 555.	0.6	13
126	Stress-induced <i>Oryza sativa</i> BAT1 dual helicase exhibits unique bipolar translocation. <i>Protoplasma</i> , 2015, 252, 1563-1574.	1.0	13



#	ARTICLE	IF	CITATIONS
127	Ectopic expression of phloem motor protein pea forisome PsSEO-F1 enhances salinity stress tolerance in tobacco. <i>Plant Cell Reports</i> , 2016, 35, 1021-1041.	2.8	13
128	<i>Arabidopsis thaliana</i> MCM3 single subunit of MCM2â€“7 complex functions as 3â€“ to 5â€“ DNA helicase. <i>Protoplasma</i> , 2016, 253, 467-475.	1.0	13
129	Synergistic inoculation of <i>Azotobacter vinelandii</i> and <i>Serendipita indica</i> augmented rice growth. <i>Symbiosis</i> , 2020, 81, 139-148.	1.2	13
130	Plant Cell and Viral Helicases: Essential Enzymes for Nucleic Acid Transactions. <i>Critical Reviews in Plant Sciences</i> , 2000, 19, 449-478.	2.7	12
131	OsBAT1 Augments Salinity Stress Tolerance by Enhancing Detoxification of ROS and Expression of Stress-Responsive Genes in Transgenic Rice. <i>Plant Molecular Biology Reporter</i> , 2015, 33, 1192-1209.	1.0	12
132	Stress-induced <i>Oryza sativa</i> RuvBL1a is DNA-independent ATPase and unwinds DNA duplex in 3â€“ to 5â€“ direction. <i>Protoplasma</i> , 2018, 255, 669-684.	1.0	12
133	Salicylic acid modulates ACS, NHX1, sos1 and HKT1;2 expression to regulate ethylene overproduction and Na <sup>+</sup> ions toxicity that leads to improved physiological status and enhanced salinity stress tolerance in tomato plants cv. Pusa Ruby. <i>Plant Signaling and Behavior</i> , 2021, 16, 1950888.	1.2	12
134	First evidence of putrescine involvement in mitigating the floral malformation in mangoes: A scanning electron microscope study. <i>Protoplasma</i> , 2014, 251, 1255-1261.	1.0	10
135	Simultaneous Expression of PDH45 with EPSPS Gene Improves Salinity and Herbicide Tolerance in Transgenic Tobacco Plants. <i>Frontiers in Plant Science</i> , 2017, 8, 364.	1.7	10
136	Pea GÎ² subunit of G proteins has a role in nitric oxide-induced stomatal closure in response to heat and drought stress. <i>Protoplasma</i> , 2020, 257, 1639-1654.	1.0	10
137	<i>OsRuvB</i> transgene induces salt tolerance in pigeon pea. <i>Journal of Plant Interactions</i> , 2020, 15, 17-26.	1.0	10
138	Plant Cell and Viral Helicases: Essential Enzymes for Nucleic Acid Transactions. , 0, .		10
139	Isolation and in silico analysis of promoter of a high salinity stress-regulated pea DNA helicase 45. <i>Plant Signaling and Behavior</i> , 2011, 6, 1447-1450.	1.2	9
140	OsSUV3 functions in cadmium and zinc stress tolerance in rice ( <i>Oryza sativa</i> L. cv IR64). <i>Plant Signaling and Behavior</i> , 2014, 9, e27389.	1.2	8
141	Overexpression of PDH45 or SUV3 helicases in rice leads to delayed leaf senescence-associated events. <i>Protoplasma</i> , 2017, 254, 1103-1113.	1.0	8
142	Receptor-Like Kinases Control the Development, Stress Response, and Senescence in Plants. , 2019, , 199-210.		8
143	<i>Azotobacter vinelandii</i> helps to combat chromium stress in rice by maintaining antioxidant machinery. <i>3 Biotech</i> , 2021, 11, 275.	1.1	8
144	Prediction of cis-regulatory elements for a detailed insight of RuvB family genes from <i>Oryza sativa</i> . <i>Oryza</i> , 2017, 54, 135.	0.2	8

#	ARTICLE	IF	CITATIONS
145	A pea homologue of human DNA helicase I is localized within the dense fibrillar component of the nucleolus and stimulated by phosphorylation with CK2 and cdc2 protein kinases. <i>Plant Journal</i> , 2001, 25, 9-17.	2.8	7
146	A Method to Confer Salinity Stress Tolerance to Plants by Helicase Overexpression. <i>Methods in Molecular Biology</i> , 2009, 587, 377-387.	0.4	7
147	Differential and temperature dependent regulation of ADP-glucose pyrophosphorylase by specific chromosome in wheat grains. <i>Cereal Research Communications</i> , 2015, 43, 591-603.	0.8	7
148	A new insight into root responses to external cues: Paradigm shift in nutrient sensing. <i>Plant Signaling and Behavior</i> , 2015, 10, e1049791.	1.2	7
149	Response of <i>PiCypA</i> tobacco T2 transgenic matured plant to potential tolerance to salinity stress. <i>Plant Signaling and Behavior</i> , 2014, 9, e27538.	1.2	6
150	Calcium-energized motor protein forisome controls damage in phloem: potential applications as biomimetic smart material. <i>Critical Reviews in Biotechnology</i> , 2015, 35, 173-183.	5.1	6
151	Cyanide produced with ethylene by ACS and its incomplete detoxification by $\beta^2$ -CAS in mango inflorescence leads to malformation. <i>Scientific Reports</i> , 2019, 9, 18361.	1.6	6
152	Introduction to PSB Special Issue. <i>Plant Signaling and Behavior</i> , 2011, 6, 173-174.	1.2	5
153	Analysis of DNA repair helicase UvrD from <i>Arabidopsis thaliana</i> and <i>Oryza sativa</i> . <i>Plant Physiology and Biochemistry</i> , 2013, 71, 254-260.	2.8	5
154	Ethylene mediated physiological response for in vitro development of salinity tolerant tomato. <i>Journal of Plant Interactions</i> , 2020, 15, 406-416.	1.0	5
155	Forisomes as calcium-energized protein complex: A historical perspective. <i>Plant Signaling and Behavior</i> , 2010, 5, 497-500.	1.2	4
156	In vitro: Response of plant growth regulators and antimetabolites on conidia germination of <i>Fusarium mangiferae</i> and incidence of mango malformation. <i>Communicative and Integrative Biology</i> , 2013, 6, e25659.	0.6	4
157	Role of Plant Helicases in Imparting Salinity Stress Tolerance to Plants. , 2019, , 39-52.		4
158	Conserved thioredoxin fold is present in <i>Pisum sativum</i> L. sieve element occlusion-1 protein. <i>Plant Signaling and Behavior</i> , 2010, 5, 623-628.	1.2	3
159	Inhibition of unwinding and ATPase activities of pea MCM6 DNA helicase by actinomycin and nogalamycin. <i>Plant Signaling and Behavior</i> , 2011, 6, 327-329.	1.2	3
160	Calcium powered phloem protein of SEO gene family Forisome functions in wound sealing and act as biomimetic smart materials. <i>Plant Signaling and Behavior</i> , 2014, 9, e29438.	1.2	3
161	Marker-Free Rice ( <i>Oryza sativa</i> L. cv. IR 64) Overexpressing PDH45 Gene Confers Salinity Tolerance by Maintaining Photosynthesis and Antioxidant Machinery. <i>Antioxidants</i> , 2022, 11, 770.	2.2	3
162	Introgression, Generational Expression and Salinity Tolerance Conferred by the Pea DNA Helicase 45 Transgene into Two Commercial Rice Genotypes, BR28 and BR47. <i>Molecular Biotechnology</i> , 2018, 60, 111-123.	1.3	2

#	ARTICLE	IF	CITATIONS
163	Cloning, Sequencing and In Silico Analysis of phbC Gene from Pseudomonas spp.. Indian Journal of Microbiology, 2019, 59, 58-63.	1.5	2
164	An Overview of AAA+ Superfamily Proteins Associated Helicases. , 2019, , 247-264.		2
165	Proteomics for Brassinosteroid signalling: Understanding Brassinosteroids mediated stress responses through advanced proteomics. Plant Gene, 2021, 26, 100282.	1.4	2
166	Helicases and Their Importance in Abiotic Stresses. , 2018, , 119-141.		1
167	The scope of transformation and genome editing for quantitative trait improvements in rice. , 2020, , 23-43.		1
168	Transgenic approach in crop improvement. , 2020, , 329-350.		0