

# Anthony W Addison

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2733175/publications.pdf>

Version: 2024-02-01

104  
papers

13,506  
citations

109321

35  
h-index

36028

97  
g-index

108  
all docs

108  
docs citations

108  
times ranked

11804  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Synthesis, structure, and spectroscopic properties of copper(II) compounds containing nitrogen-sulphur donor ligands; the crystal and molecular structure of aqua[1,7-bis(N-methylbenzimidazol-2-yl)-2,6-dithiaheptane]copper(II) perchlorate. <i>Journal of the Chemical Society Dalton Transactions</i> , 1984, , 1349-1356. | 1.1 | 8,412     |
| 2  | Conversion constants for redox potentials measured versus different reference electrodes in acetonitrile solutions at 25°C. <i>Inorganica Chimica Acta</i> , 2000, 298, 97-102.  | 2.4 | 1,281     |
| 3  | Spectroscopic and redox studies of some copper(II) complexes with biomimetic donor atoms: implications for protein copper centres. <i>Journal of the Chemical Society Dalton Transactions</i> , 1979, , 600.   | 1.1 | 459       |
| 4  | Synthesis of some imidazole- and pyrazole-derived chelating agents. <i>Journal of Heterocyclic Chemistry</i> , 1981, 18, 803-805.  | 2.6 | 320       |
| 5  | Copper complexes of the "tripod" ligand tris(2-benzimidazolymethyl)amine: five- and six-coordinate copper(II) derivatives and some copper(I) derivatives. <i>Inorganic Chemistry</i> , 1981, 20, 103-110.  | 4.0 | 154       |
| 6  | Iridium Dihydroxybipyridine Complexes Show That Ligand Deprotonation Dramatically Speeds Rates of Catalytic Water Oxidation. <i>Inorganic Chemistry</i> , 2013, 52, 9175-9183.   | 4.0 | 142       |
| 7  | Dipicolylamine Complexes of Copper(II): Two Different Coordination Geometries in the Same Unit Cell of Cu(Dipica) <sub>2</sub> (BF <sub>4</sub> ) <sub>2</sub> . <i>Inorganic Chemistry</i> , 1996, 35, 467-471.   | 4.0 | 110       |
| 8  | Nitrosyliron(III) hemoglobin: autoreduction and spectroscopy. <i>Biochemistry</i> , 1986, 25, 4104-4113.   | 2.5 | 105       |
| 9  | Synthesis of some benzimidazole- and benzothiazole-derived ligand systems and their precursory diacids. <i>Journal of Heterocyclic Chemistry</i> , 1983, 20, 1481-1484.  | 2.6 | 100       |
| 10 | New iron(II) spin-crossover complexes with heterocyclic amine-derived ligands and STEPS experiments on photogenerated metastable high-spin states. <i>Journal of the Chemical Society Dalton Transactions</i> , 1987, , 2621.  | 1.1 | 97        |
| 11 | Spectroscopy and structure of thiolate and thioether complexes of copper(II) and the relationship of their redox chemistry to that of certain copper proteins. <i>Inorganic Chemistry</i> , 1984, 23, 1957-1967.   | 4.0 | 96        |
| 12 | Small molecule analogs for the specific iron-binding site of lactoferrin: a single-crystal x-ray structure of bis(methanol)bis[2-(5-methylpyrazol-3-yl)phenolato]iron(III) nitrate-methanol and spectroscopic studies on iron(III) phenolate complexes. <i>Inorganic Chemistry</i> , 1980, 19, 3655-3663.                      | 4.0 | 95        |
| 13 | Pentacoordinate copper complexes of nitrogen-sulfur donors: structural chemistry of two complexes of bis(2-(2-benzimidazolyl)ethyl) sulfide with the sulfur alternatively in equatorial and axial coordination modes. <i>Inorganic Chemistry</i> , 1983, 22, 3645-3653.  | 4.0 | 87        |
| 14 | Is ligand topology an influence on the redox potentials of copper complexes?. <i>Inorganica Chimica Acta</i> , 1989, 162, 217-220.   | 2.4 | 85        |
| 15 | Copper(II) and Nickel(II) Complexes of Dianionic and Tetraanionic Dinucleating Macrocycles. <i>Inorganic Chemistry</i> , 1998, 37, 1028-1036.  | 4.0 | 84        |
| 16 | A stable bis(thiolate) of copper(II) with long axial copper-sulfur linkages: crystal and molecular structure of trans-[Cu(cyclam)(SC <sub>6</sub> F <sub>5</sub> ) <sub>2</sub> ]. <i>Inorganic Chemistry</i> , 1983, 22, 1225-1228.   | 4.0 | 65        |
| 17 | Synthesis and proton transfer-linked redox tuning of ruthenium(II) complexes with tridentate 2,6-bis(benzimidazol-2-yl)pyridine ligands. <i>Journal of the Chemical Society Dalton Transactions</i> , 1993, , 2477.  | 1.1 | 65        |
| 18 | Complexes of structural analogues of terpyridyl with iron and zinc; the x-ray crystal structure of bis[2,6-bis(benzimidazol-2-yl)pyridine]iron(II) trifluoromethylsulphonate bis-ethanol solvate. <i>Polyhedron</i> , 1992, 11, 635-646.   | 2.2 | 62        |

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 19 | Mono- and Trinuclear Nickel(II) Complexes with Sulfur-Containing Oxime Ligands: An Uncommon Templated Coupling of Oxime with Nitrile. <i>Inorganic Chemistry</i> , 1999, 38, 1759-1766.  | 4.0  | 61        |
| 20 | Copper(II) and zinc(II) co-ordination compounds of tridentate bis(benzimidazole)pyridine ligands. Crystal and molecular structures of bis[2,6-bis(1-methylbenzimidazol-2-yl)pyridine]copper(II) diperchlorate monohydrate and (acetonitrile)[2,6-bis(benzimidazol-2-yl)pyridine](perchlorato)copper(II) perchlorate. <i>Journal of the Chemical Society Dalton Transactions</i> , 1988, , 1429-1435.   | 1.1  | 54        |
| 21 | A Tetrameric Nickel(II) Chair with both Antiferromagnetic Internal Coupling and Ferromagnetic Spin Alignment. <i>Angewandte Chemie - International Edition</i> , 2001, 40, 4734-4737.  | 13.8 | 53        |
| 22 | A Triple-Decker Heptadecanuclear (Cu <sup>II</sup> ) <sub>15</sub> (Cr <sup>III</sup> ) <sub>2</sub> Complex Assembled from Pentanuclear Metallacrowns. <i>European Journal of Inorganic Chemistry</i> , 2010, 2010, 4851-4858.  | 2.0  | 51        |
| 23 | Structure of trimeric haemerythrin. <i>Nature</i> , 1983, 303, 86-88.  | 27.8 | 50        |
| 24 | Comparison of hemerythrins from four species of sipunculids by optical absorption, circular dichroism, fluorescence emission, and resonance Raman spectroscopy. <i>Biochemistry</i> , 1977, 16, 1743-1749.   | 2.5  | 49        |
| 25 | Structural Demonstration of the Role of Ligand Framework Conformability in Copper(II)/Copper(I) Redox Potentials. <i>Inorganic Chemistry</i> , 1997, 36, 134-135.  | 4.0  | 49        |
| 26 | Formation of Coordination Polymers or Discrete Adducts via Reactions of Gadolinium(III)–Copper(II) 15-Metallacrown-5 Complexes with Polycarboxylates: Synthesis, Structures and Magnetic Properties. <i>Inorganic Chemistry</i> , 2014, 53, 1320-1330.   | 4.0  | 49        |
| 27 | Nickel(II) complexes with dithiadiminoamine and dithiabis(thiosemicarbazone) ligands. <i>Dalton Transactions RSC</i> , 2000, , 335-341.  | 2.3  | 48        |
| 28 | Manifestation of π–π Stacking Interactions in Luminescence Properties and Energy Transfer in Aromatically-Derived Tb, Eu and Gd Tris(pyrazolyl)borate Complexes. <i>Inorganic Chemistry</i> , 2015, 54, 3125-3133.   | 4.0  | 48        |
| 29 | Mononuclear and Mixed-Valence Binuclear Oxovanadium Complexes with Benzimidazole-Derived Chelating Agents. <i>Inorganic Chemistry</i> , 2002, 41, 2243-2249.   | 4.0  | 47        |
| 30 | Magnetic and Sorption Properties of Supramolecular Systems Based on Pentanuclear Copper(II) 12-Metallacrown-4 Complexes and Isomeric Phthalates: Structural Modeling of the Different Stages of Alcohol Sorption. <i>European Journal of Inorganic Chemistry</i> , 2011, 2011, 4826-4836.  | 2.0  | 47        |
| 31 | Structural aspects of the bis(2,2-dipicolylamine)iron(II) cation. <i>Inorganica Chimica Acta</i> , 1989, 158, 211-215.   | 2.4  | 43        |
| 32 | Structural, magnetic and related attributes of some oximate-bridged tetranuclear nickel(ii) rhombs and a dinuclear congener. Electronic supplementary information (ESI) available: mass spectra, $\chi_T$ vs. T, response of magnetic properties, low-lying spin levels and UV-VIS data. See <a href="http://www.rsc.org/suppdata/dt/b3/b300539a/">http://www.rsc.org/suppdata/dt/b3/b300539a/</a> . <i>Dalton Transactions</i> , 2003, , 1587-1595. | 3.3  | 43        |
| 33 | Helical Antiferromagnetic Copper(II) Chains with a Collagen Structural Motif. <i>Inorganic Chemistry</i> , 1996, 35, 5966-5967.  | 4.0  | 42        |
| 34 | Pentadentate thioether-oxime macrocyclic and quasi-macrocyclic complexes of copper(II) and nickel(II). <i>Inorganica Chimica Acta</i> , 2000, 300-302, 992-1003.   | 2.4  | 41        |
| 35 | Analogues for the specific iron-binding site in the transferrins: molecular structure of a ternary iron(III) model complex and spectroscopic, redox and reactivity properties of related compounds. <i>Inorganic Chemistry</i> , 1990, 29, 3425-3433.  | 4.0  | 37        |
| 36 | Copper(II) complexes of tetradentate thioether-oxime ligands. <i>Inorganica Chimica Acta</i> , 2005, 358, 3449-3456.   | 2.4  | 37        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Iron ligand recognition by monomeric hemoglobins. BBA - Proteins and Proteomics, 1996, 1295, 209-221. Structure and electrochemical properties of antiferromagnetically coupled binuclear hydroxo-bridged copper(II) complexes with pyridazine and phthalazine ligands. Crystal and molecular structures of                    | 2.1 | 35        |
| 38 | [.mu.-3,6-bis(1-pyrazolyl)pyridazine-N2,.mu.-N3,.mu.-N4,N5](.mu.-hydroxo)trichloroaquodicopper-(II)-0.8-water, [.mu.-3,6-bis(1-pyrazolyl)pyridazine-N2,.mu.-N3,.mu.-N4,N5](.+-.-hydroxo)tribromoaquodicopper-(II)-0.6-water, and [.mu.-1,4-bis(1-methyl-2-imidazolyl)phthalazine](.mu.-hydroxo)tribromoaquodicopper-(II)-wa.   | 4.0 | 34        |
| 39 | Inorga<br>Synthesis, structure and magnetic properties of Nd <sup>3+</sup> and Pr <sup>3+</sup> 2D polymers with tetrafluoro-p-phthalate. Dalton Transactions, 2011, 40, 10989.  | 3.3 | 32        |
| 40 | Structure, magnetic and luminescence properties of the lanthanide complexes Ln <sub>2</sub> (Salphen) <sub>3</sub> ·H <sub>2</sub> O (Ln=Pr, Nd, Sm, Eu, Gd, Tb, Dy; H <sub>2</sub> Salphen=N,N- $\epsilon$ -bis(salicylidene)-1,2-phenylenediamine). Inorganica Chimica Acta, 2014, 414, 97-104.                              | 2.4 | 31        |
| 41 | Crystal and molecular structure of dipolar spin-coupled dimers of an irregularly pentacoordinate copper(II) complex, [Cu(5-Melin)(DBM)]. Inorganic Chemistry, 1982, 21, 60-63.   | 4.0 | 29        |
| 42 | Medium effects on the redox properties of tris(2,2- $\epsilon$ -bipyridyl)ruthenium complexes. Inorganica Chimica Acta, 1993, 204, 141-146.  | 2.4 | 29        |
| 43 | Copper co-ordination chemistry of some quadridentate pyridazine and phthalazine (N <sub>4</sub> ) thioether ligands. Binuclear copper(II) complexes exhibiting two-electron reduction at positive potentials. Journal of the Chemical Society Dalton Transactions, 1986, , 2381.   | 1.1 | 28        |
| 44 | Efficient mechanochemical synthesis of tris(pyrazolylborate) complexes of manganese(II), cobalt(II) and nickel(II). Inorganic Chemistry Communication, 2004, 7, 485-488.   | 3.9 | 28        |
| 45 | Some high-potential trithioether chelates of copper. Inorganica Chimica Acta, 1992, 196, 97-103.   | 2.4 | 26        |
| 46 | Copper(II) complexes of new unsymmetrical NSN thioether ligands. Inorganica Chimica Acta, 2001, 324, 123-130.  | 2.4 | 26        |
| 47 | Ruthenium(II) complexes with phenanthroline-, benzimidazole-, benzothiazole-, and pyridine-derived bidentate and tridentate ligands: reactivity and spectroscopic and electrochemical characterization. Transition Metal Chemistry, 1993, 18, 197-204.   | 1.4 | 25        |
| 48 | Heme proteins and metalloporphyrins: Redox chemistry and oxygen binding. International Journal of Quantum Chemistry, 1979, 16, 311-329.  | 2.0 | 23        |
| 49 | Some nickel(II) complexes with pentadentate and tridentate heterocyclic N- and S-donor ligands. Inorganica Chimica Acta, 2000, 308, 22-30.   | 2.4 | 23        |
| 50 | A new class of macrocyclic complexes formed via nickel-promoted macrocyclisation of dioxime with dinitrile. Chemical Communications, 2002, , 468-469.  | 4.1 | 23        |
| 51 | Dinuclear Copper(II) Complexes with Bis-thiocarbohydrazone Ligands. European Journal of Inorganic Chemistry, 2008, 2008, 2530-2536. Structural trends in a series of isostructural lanthanide- $\epsilon$ -copper metallacrown sulfates (Ln <sup>III</sup> = Pr, Nd, Sm, Eu, Gd, Dy and Ho):                                   | 2.0 | 23        |
| 52 | hexaaquapentakis[1/4<sub>3</sub>-glycinehydroxamato(2 $\hat{a}$ )]sulfatopentacopper(II)lanthanide(III) heptaaquapentakis[1/4<sub>3</sub>-glycinehydroxamato(2 $\hat{a}$ )]sulfatopentacopper(II)lanthanide(III) sulfate hexahydrate. Acta Crystallographica Section C: Crystal Structure Communications, 2011, 67, m255-m265. | 0.4 | 23        |
| 53 | Triethylammonium benzene-1,3,5-tricarboxylato(pyridine)zinc(II): a two-dimensional undulating mesh network. Inorganic Chemistry Communication, 2003, 6, 402-404.   | 3.9 | 22        |
| 54 | The 1, 8-bis(2 $\hat{a}$ -pyridyl)-3, 6-dithiaoctane complex of nickel(II): X-ray crystal structure and borohydride adduct formation. Inorganica Chimica Acta, 1998, 278, 217-222.   | 2.4 | 20        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | A nickel(II) dithia dioxime derived macrocycle. <i>Inorganic Chemistry</i> , 1994, 33, 397-399.   | 4.0 | 19        |
| 56 | Crystal structures and intense luminescence of tris(3-(2-pyridyl)-pyrazolyl)borate Tb <sup>3+</sup> and Eu <sup>3+</sup> complexes with carboxylate co-ligands. <i>Dalton Transactions</i> , 2017, 46, 3457-3469.                               | 3.3 | 19        |
| 57 | High Nuclearity Assemblies and One-Dimensional (1D) Coordination Polymers Based on Lanthanide-Copper 15-Metallacrown-5 Complexes (Ln <sup>3+</sup> = Pr, Nd, Sm, Eu). <i>Inorganic Chemistry</i> , 2017, 56, 13152-13165.                       | 4.0 | 19        |
| 58 | Oxo-bridged complexes of iron(III) derived from 2-(2-hydroxyphenyl)-benzothiazole and 2-(2-hydroxyphenyl)benzimidazole ligands. <i>Inorganica Chimica Acta</i> , 1989, 166, 59-69.  | 2.4 | 18        |
| 59 | A novel paradigm for metal-induced ring flipping in the copper(II) complex of 1,2-bis(N) Tj ETQq1 1 0.784314 rgBT/Overlock_10 Tf 505  | 3.9 | 18        |
| 60 | Combination of single-molecule magnet behaviour and luminescence properties in a new series of lanthanide complexes with tris(pyrazolyl)borate and oligo(1 <sup>2</sup> -diketonate) ligands. <i>Dalton Transactions</i> , 2020, 49, 7774-7789. | 3.3 | 17        |
| 61 | The Hexakis(thiocyanato)ferrate(III) Ion: a Coordination Chemistry Classic Reveals an Interesting Geometry Pattern for the Thiocyanate Ligands. <i>European Journal of Inorganic Chemistry</i> , 2005, 2005, 2404-2408.                         | 2.0 | 16        |
| 62 | Ni(II), Co(II) and Mn(II) tris(pyrazolyl)borate complexes with 2,6-di-tert-butyl-4-carboxy-phenol: Formation of coordinated phenoxyl radical. <i>Inorganic Chemistry Communication</i> , 2005, 8, 932-935.                                      | 3.9 | 14        |
| 63 | Autoxidation of Reduced Horse Heart Cytochrome <i>c</i> Catalyzed by Cardiolipin-Containing Membranes. <i>Journal of Physical Chemistry B</i> , 2016, 120, 12219-12231.   | 2.6 | 14        |
| 64 | The metamorphosis of heterometallic trinuclear antiferromagnetic complexes into nano-sized superparamagnetic spinels. <i>Materials Chemistry and Physics</i> , 2010, 121, 47-52.  | 4.0 | 13        |
| 65 | Supramolecular Maleate Adducts of Copper(II) 12-Metallacrown-4: Magnetism, EPR, and Alcohol Sorption Properties. <i>European Journal of Inorganic Chemistry</i> , 2017, 2017, 4866-4878.  | 2.0 | 13        |
| 66 | Magnetic Properties of Ln <sup>3+</sup> -Cu <sup>II</sup> 15-Metallacrown-5 Dimers with Terephthalate (Ln <sup>3+</sup> = Pr, Nd, Sm, Eu). <i>European Journal of Inorganic Chemistry</i> , 2018, 2018, 3504-3511.                              | 2.0 | 13        |
| 67 | Synthesis of some benzimidazole-, pyridine- and imidazole-derived chelating agents. <i>Journal of Heterocyclic Chemistry</i> , 1989, 26, 541-543.   | 2.6 | 12        |
| 68 | A novel copper(II) complex of a tripodal ligand with phenolate-phenol interligand, intramolecular hydrogen bonding. <i>Dalton Transactions</i> , 2009, , 8111.  | 3.3 | 11        |
| 69 | New Pyrazole- and Benzimidazole-derived Ligand Systems. <i>Journal of Heterocyclic Chemistry</i> , 2018, 55, 1291-1307.   | 2.6 | 11        |
| 70 | Synthesis, structure and magnetic properties of oligometallic systems derived from di- and trinuclear copper(ii) amido-oximate complexes. <i>Dalton Transactions</i> , 2008, , 3007.  | 3.3 | 10        |
| 71 | A tetranuclear copper(II) complex with bis(o-aminobenzaldehyde)thiocarbohydrazone. <i>Inorganica Chimica Acta</i> , 2010, 363, 2065-2070.   | 2.4 | 10        |
| 72 | The role of the bridging group in exchange coupling in dinuclear homo- and heterometallic Ni(ii) and Co(ii) complexes with oxalate, oxamidate and dithiooxamidate bridges. <i>Dalton Transactions</i> , 2012, 41, 11319.                        | 3.3 | 10        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | Chemical liquid deposition process for microstructure fabrication. <i>Materials &amp; Design</i> , 2005, 26, 670-679.  | 5.1 | 9         |
| 74 | Effect of Zero-Field Splitting on the Magnetic Susceptibility of Binuclear Complexes of Iron(III). <i>Theoretical and Experimental Chemistry</i> , 2005, 41, 229-234.  | 0.8 | 9         |
| 75 | Synthesis, spectroscopic and redox behaviour of copper(II) complexes with quinquedentate thiaaza ligands. <i>Inorganica Chimica Acta</i> , 1993, 203, 29-35.   | 2.4 | 8         |
| 76 | Some iron(III) complexes with polydentate Schiff base ligands. <i>Inorganica Chimica Acta</i> , 1988, 147, 61-64.  | 2.4 | 7         |
| 77 | Magnetic Characteristics of Trinuclear Complexes $[M_3O(CH_3COO)_6(pz)_3]^+$ (M = Fe, Cr; pz = Pyrazine). <i>Theoretical and Experimental Chemistry</i> , 2004, 40, 214-219.   | 0.8 | 7         |
| 78 | Sorption discrimination between secondary alcohol enantiomers by chiral alkyl-dicarboxylate MOFs. <i>RSC Advances</i> , 2016, 6, 93707-93714.  | 3.6 | 7         |
| 79 | The 1,8-Bis(2- $\pi$ -pyridyl)-3,6-dithiooctane Complex of Rhodium(III). <i>Journal of Chemical Crystallography</i> , 2012, 42, 295-298.   | 1.1 | 6         |
| 80 | Thermochromism of heme adducts of Glycera hemoglobin and some other monomeric heme proteins. <i>Journal of Inorganic Biochemistry</i> , 1990, 39, 351-369.   | 3.5 | 5         |
| 81 | Synthesis of some benzimidazole-, benzothiazole- and pyridine-derived chelating agents. <i>Journal of Heterocyclic Chemistry</i> , 2002, 39, 399-404.  | 2.6 | 5         |
| 82 | Ru(II) thioether complexes with dangling pyridine ligands. <i>Polyhedron</i> , 2014, 68, 70-75.  | 2.2 | 5         |
| 83 | Absorption- and Excitation-Modulated Luminescence of $Pr^{3+}$ , $Nd^{3+}$ , and $Lu^{3+}$ Compounds with Dianions of Tetrafluoroterephthalic and Camphoric Acids. <i>ACS Omega</i> , 2019, 4, 2669-2675.  | 3.5 | 5         |
| 84 | Heme rotational isomerism is not required for the production of Q-band splitting in the spectra of iron-porphyrin proteins. <i>Inorganic Chemistry</i> , 1991, 30, 1151-1153.  | 4.0 | 4         |
| 85 | The crystal structures of $\{LnCu_5\}^{3+}$ ( $Ln = Gd, Dy$ and $Ho$ ) 15-metallacrown-5 complexes and a reevaluation of the isotypic $Eu^{III}$ analogue. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2019, 75, 1215-1223. | 0.5 | 3         |
| 86 | The coelomic hemerythrin of siphonosome cumanense. <i>Comparative Biochemistry and Physiology Part B: Comparative Biochemistry</i> , 1982, 72, 433-438.  | 0.2 | 2         |
| 87 | Spectroscopic and kinetic aspects of <i>Elephas maximus</i> hemoglobin. <i>FEBS Journal</i> , 1990, 189, 185-191.  | 0.2 | 2         |
| 88 | Electron-Counting Rules for Transition Metal-Nitric Oxide Complexes. <i>Journal of Chemical Education</i> , 1997, 74, 1354.  | 2.3 | 2         |
| 89 | Triangular kinetic schemes applied to the stability of a Heme-Globin complex. <i>Journal of Inorganic Biochemistry</i> , 1997, 66, 83-98.  | 3.5 | 2         |
| 90 | Copper(II) Schiff-base complexes and apoglobin stability. <i>Journal of Inorganic Biochemistry</i> , 1999, 73, 137-144.  | 3.5 | 2         |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 91  | 5,7,12,14-Tetrahydro-5,14:7,12-bis([1,2]benzeno)pentacene-6,13-diol dimethylformamide disolvate. IUCrData, 2016, 1, .   | 0.3 | 2         |
| 92  | Chiroptical and physicochemical properties of the extracellular haemoglobin from <i>Cirriformia tentaculata</i> . <i>Comparative Biochemistry and Physiology Part B: Comparative Biochemistry</i> , 1990, 97, 391-399.            | 0.2 | 1         |
| 93  | Ruthenium(II) complexes of some simple classic amine ligands. <i>Inorganica Chimica Acta</i> , 2015, 432, 185-191.  | 2.4 | 1         |
| 94  | New homodinuclear tris(3-alkylpyrazolyl)borate complexes of Colland Nillwith a tetraacetylene dianion as a bridging ligand. <i>Acta Crystallographica Section C, Structural Chemistry</i> , 2016, 72, 777-785.                    | 0.5 | 1         |
| 95  | Crystal structure of 5,7,12,14-tetrahydro-5,14:7,12-bis([1,2]benzeno)pentacene-6,13-dione. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2016, 72, 1734-1738.  | 0.5 | 1         |
| 96  | Lanthanide Complexes with 4,4'-bis(2-sulfonatostyryl)-biphenyl: Crystal Structures and Luminescence Properties. <i>European Journal of Inorganic Chemistry</i> , 2022, 2022, .  | 2.0 | 1         |
| 97  | Crystal structure of (2,2'-bipyridyl)[2,6-bis(1-butyl-1H-benzimidazol-2-yl)pyridine]chloridoiridium(III) trifluoromethanesulfonate. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2017, 73, 127-132. | 0.5 | 0         |
| 98  | Crystal structure of bis(pivaloylhydroxamato) <sup>2-</sup> copper(II). <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2018, 74, 1384-1387.   | 0.5 | 0         |
| 99  | Ruthenium(II) complexes of the tetradentate polypyridyl thioether 1,2-bis[3-(2-pyridyl)-1-thiopropyl]benzene. <i>Polyhedron</i> , 2020, 179, 114367.  | 2.2 | 0         |
| 100 | Crystal structure of a Tb <sup>III</sup> –Cu <sup>II</sup> glycinehydroxamate 15-metallacrown-5 sulfate complex. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2021, 77, 1197-1202.                  | 0.5 | 0         |
| 101 | 1,3-Bis(N-Methylbenzimidazol-2-yl)-2-Phenylpropanedichloridocopper(II). <i>Journal of Chemical Crystallography</i> , 0, , 1.  | 1.1 | 0         |
| 102 | Two Bifunctionalised Derivatives of 1,2,4-Triazole. <i>Journal of Chemical Crystallography</i> , 0, , 1.  | 1.1 | 0         |
| 103 | Bis[3-(anthracen-9-yl)pentane-2,4-dionato] <sup>2-</sup> [tris(pyrazol-1-yl)-N,N'-dimethylformamide-O] [tris(pyrazol-1-yl)-N,N'-dimethylformamide-O] Tj ETQq1 1 0.784314 rgBT /Over<br>Communications. 2022, 78, 103-107.         | 0.5 | 0         |
| 104 | Chlorocobalt complexes with pyridylethyl-derived diazacycloalkanes. <i>Acta Crystallographica Section E: Crystallographic Communications</i> , 2022, 78, 235-243.   | 0.5 | 0         |