

Noroska Gabriela Salazar Mogollon

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2757051/publications.pdf>

Version: 2024-02-01

10
papers

389
citations

1478505

6
h-index

1474206

9
g-index

11
all docs

11
docs citations

11
times ranked

628
citing authors

#	ARTICLE	IF	CITATIONS
1	A First Look at the Inhibitory Potential of <i>Urospatha sagittifolia</i> (Araceae) Ethanolic Extract for <i>Bothrops atrox</i> Snakebite Envenomation. <i>Toxins</i> , 2022, 14, 496.	3.4	5
2	RGCxGC toolbox: An R-package for data processing in comprehensive two-dimensional gas chromatography-mass spectrometry. <i>Microchemical Journal</i> , 2020, 156, 104830.	4.5	23
3	Comprehensive two-dimensional gas chromatography-mass spectrometry combined with multivariate data analysis for pattern recognition in Ecuadorian spirits. <i>Chemistry Central Journal</i> , 2018, 12, 102.	2.6	8
4	New Advances in Toxicological Forensic Analysis Using Mass Spectrometry Techniques. <i>Journal of Analytical Methods in Chemistry</i> , 2018, 2018, 1-17.	1.6	25
5	Genetic and phenotypic diversity of <i>Rhizobium</i> isolates from Southern Ecuador. <i>Ciencia E Agrotecnología</i> , 2017, 41, 634-647.	1.5	4
6	Characterization of crude oil biomarkers using comprehensive two-dimensional gas chromatography coupled to tandem mass spectrometry. <i>Journal of Separation Science</i> , 2016, 39, 3384-3391.	2.5	15
7	Discriminating Brazilian crude oils using comprehensive two-dimensional gas chromatography-mass spectrometry and multiway principal component analysis. <i>Journal of Chromatography A</i> , 2016, 1472, 99-106.	3.7	28
8	Quantitative analysis of biodiesel in blends of biodiesel and conventional diesel by comprehensive two-dimensional gas chromatography and multivariate curve resolution. <i>Analytica Chimica Acta</i> , 2013, 796, 130-136.	5.4	37
9	New materials and trends in sorbents for solid-phase extraction. <i>TrAC - Trends in Analytical Chemistry</i> , 2013, 43, 14-23.	11.4	239
10	Exploratory Analysis of Biodiesel by Combining Comprehensive Two-Dimensional Gas Chromatography and Multiway Principal Component Analysis. <i>Journal of the Brazilian Chemical Society</i> , 0, , .	0.6	5