

# Daniel R Witte

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2763211/publications.pdf>

Version: 2024-02-01

256  
papers

22,296  
citations

15001

68  
h-index

12272

138  
g-index

275  
all docs

275  
docs citations

275  
times ranked

36509  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Obesity and Kidney Function: A Two-Sample Mendelian Randomization Study. <i>Clinical Chemistry</i> , 2022, 68, 461-472.  | 1.5 | 25        |
| 2  | Changes in type 2 diabetes incidence and mortality associated with introduction of HbA1c as diagnostic option: A Danish 24-year population-based study. <i>Lancet Regional Health - Europe</i> , The, 2022, 14, 100291.  | 3.0 | 12        |
| 3  | Rare coding variants in 35 genes associate with circulating lipid levels—A multi-ancestry analysis of 170,000 exomes. <i>American Journal of Human Genetics</i> , 2022, 109, 81-96.  | 2.6 | 24        |
| 4  | The Arg82Cys Polymorphism of the Protein Nepmucin Implies a Role in HDL Metabolism. <i>Journal of the Endocrine Society</i> , 2022, 6, bvac034.  | 0.1 | 1         |
| 5  | Health-related quality of life for normal glycaemia, prediabetes and type 2 diabetes mellitus: Cross-sectional analysis of the ADDITION-EUROPE study. <i>Diabetic Medicine</i> , 2022, 39, e14825.   | 1.2 | 5         |
| 6  | Multi-ancestry genetic study of type 2 diabetes highlights the power of diverse populations for discovery and translation. <i>Nature Genetics</i> , 2022, 54, 560-572.   | 9.4 | 250       |
| 7  | Association of weight loss and weight loss maintenance following diabetes diagnosis by screening and incidence of cardiovascular disease and all-cause mortality: An observational analysis of the ADDITION-EUROPE trial. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 730-741. | 2.2 | 6         |
| 8  | Duration of diabetes-related complications and mortality in type 1 diabetes: a national cohort study. <i>International Journal of Epidemiology</i> , 2021, 50, 1250-1259.  | 0.9 | 9         |
| 9  | Prospects for a Relational Sociology of Islam: Some Remarks on Differentiation Theory, Multiple Modernities, and the Pitfalls of Occidentalism. , 2021, , 65-88.   |     | 1         |
| 10 | Genome-wide association study of circulating levels of glucagon during an oral glucose tolerance test. <i>BMC Medical Genomics</i> , 2021, 14, 3.  | 0.7 | 3         |
| 11 | Heart Rate and Heart Rate Variability Changes Are Not Related to Future Cardiovascular Disease and Death in People With and Without Dysglycemia: A Downfall of Risk Markers? The Whitehall II Cohort Study. <i>Diabetes Care</i> , 2021, 44, 1012-1019.                                | 4.3 | 5         |
| 12 | Spousal concordance in pathophysiological markers and risk factors for type 2 diabetes: a cross-sectional analysis of The Maastricht Study. <i>BMJ Open Diabetes Research and Care</i> , 2021, 9, e001879.   | 1.2 | 2         |
| 13 | Towards precision medicine in diabetes? A critical review of glucoypes. <i>PLoS Biology</i> , 2021, 19, e3000890.  | 2.6 | 4         |
| 14 | Diabetic Polyneuropathy Early in Type 2 Diabetes Is Associated With Higher Incidence Rate of Cardiovascular Disease: Results From Two Danish Cohort Studies. <i>Diabetes Care</i> , 2021, 44, 1714-1721.   | 4.3 | 8         |
| 15 | Objective and subjective sleep measures are associated with HbA1c and insulin sensitivity in the general population: Findings from the ORISCAV-LUX-2 study. <i>Diabetes and Metabolism</i> , 2021, 48, 101263.   | 1.4 | 7         |
| 16 | Plasma lipid metabolites associate with diabetic polyneuropathy in a cohort with type 2 diabetes. <i>Annals of Clinical and Translational Neurology</i> , 2021, 8, 1292-1307.  | 1.7 | 27        |
| 17 | Determinants of penetrance and variable expressivity in monogenic metabolic conditions across 77,184 exomes. <i>Nature Communications</i> , 2021, 12, 3505.  | 5.8 | 49        |
| 18 | Potentially inappropriate medications (PIMs): frequency and extent of GP-related variation in PIMs: a register-based cohort study. <i>BMJ Open</i> , 2021, 11, e046756.  | 0.8 | 4         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Role of fasting duration and weekday in incretin and glucose regulation. <i>Endocrine Connections</i> , 2021, 10, X2-X3.   | 0.8 | 0         |
| 20 | r-cubed: Guiding the overwhelmed scientist from random wrangling to Reproducible Research in R. <i>The Journal of Open Source Education</i> , 2021, 4, 122.  | 0.2 | 0         |
| 21 | Polypharmacy in polymorbid pregnancies and the risk of congenital malformations – a systematic review. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2021, , .   | 1.2 | 3         |
| 22 | Factors associated with attendance at clinical follow-up of a cohort with screen-detected type 2 diabetes: ADDITION-Denmark. <i>Primary Care Diabetes</i> , 2020, 14, 239-245.   | 0.9 | 3         |
| 23 | Discovery of rare variants associated with blood pressure regulation through meta-analysis of 1.3 million individuals. <i>Nature Genetics</i> , 2020, 52, 1314-1332.   | 9.4 | 91        |
| 24 | Das Feld der Macht. , 2020, , .  |     | 6         |
| 25 | Identifying hotspots of cardiometabolic outcomes based on a Bayesian approach: The example of Chile. <i>PLoS ONE</i> , 2020, 15, e0235009.   | 1.1 | 6         |
| 26 | Differences Between Randomized Clinical Trial Participants and Real-World Empagliflozin Users and the Changes in Their Glycated Hemoglobin Levels. <i>JAMA Network Open</i> , 2020, 3, e1920949.   | 2.8 | 13        |
| 27 | Effect of familial diabetes status and age at diagnosis on type 2 diabetes risk: a nation-wide register-based study from Denmark. <i>Diabetologia</i> , 2020, 63, 934-943.   | 2.9 | 4         |
| 28 | Role of fasting duration and weekday in incretin and glucose regulation. <i>Endocrine Connections</i> , 2020, 9, 279-288.  | 0.8 | 5         |
| 29 | Das Feld der Macht als gesellschaftstheoretisches Schlüsselkonzept. , 2020, , 1-13.  |     | 0         |
| 30 | The effect of training GPs in motivational interviewing on incident cardiovascular disease and mortality in people with screen-detected diabetes. Results from the ADDITION-Denmark randomised trial. <i>BJGP Open</i> , 2020, 4, bjgpopen20X101012. | 0.9 | 1         |
| 31 | Das Feld der Macht in der Soziologie Bourdieus. , 2020, , 15-35.   |     | 3         |
| 32 | Vom nationalen zum globalen Feld der Macht. , 2020, , 103-152.   |     | 1         |
| 33 | Zur Pluralisierung der Feldanalyse: Das Feld der Macht als Feld der Felder. , 2020, , 61-102.  |     | 0         |
| 34 | Human pancreatic islet three-dimensional chromatin architecture provides insights into the genetics of type 2 diabetes. <i>Nature Genetics</i> , 2019, 51, 1137-1148.  | 9.4 | 208       |
| 35 | Prospective Association Among Diabetes Diagnosis, HbA1c, Glycemia, and Frailty Trajectories in an Elderly Population. <i>Diabetes Care</i> , 2019, 42, 1903-1911.  | 4.3 | 42        |
| 36 | Prospective Study of Neuropathic Symptoms Preceding Clinically Diagnosed Diabetic Polyneuropathy: ADDITION-Denmark. <i>Diabetes Care</i> , 2019, 42, 2282-2289.  | 4.3 | 13        |

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 37 | Greater glucagon-like peptide-1 responses to oral glucose are associated with lower central and peripheral blood pressures. <i>Cardiovascular Diabetology</i> , 2019, 18, 130.   | 2.7  | 8         |
| 38 | Glucose Measurements at Various Time Points During the OGTT and Their Role in Capturing Glucose Response Patterns. <i>Diabetes Care</i> , 2019, 42, e56-e57.   | 4.3  | 8         |
| 39 | Letter to the Editor: "One-Hour Postload Hyperglycemia: Implications for Prediction and Prevention of Type 2 Diabetes". <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 674-675.  | 1.8  | 0         |
| 40 | Reversion from prediabetes to normoglycaemia and risk of cardiovascular disease and mortality: the Whitehall II cohort study. <i>Diabetologia</i> , 2019, 62, 1385-1390.   | 2.9  | 55        |
| 41 | Exome sequencing of 20,791 cases of type 2 diabetes and 24,440 controls. <i>Nature</i> , 2019, 570, 71-76.   | 13.7 | 248       |
| 42 | Prospective association between late evening food consumption and risk of prediabetes and diabetes: the Whitehall II cohort study. <i>Diabetic Medicine</i> , 2019, 36, 1256-1260.   | 1.2  | 3         |
| 43 | Heart Rate, Autonomic Function, and Future Changes in Glucose Metabolism in Individuals Without Diabetes: The Whitehall II Cohort Study. <i>Diabetes Care</i> , 2019, 42, 867-874.   | 4.3  | 24        |
| 44 | Genetic determinants of blood pressure traits are associated with carotid arterial thickening and plaque formation in patients with type 2 diabetes. <i>Diabetes and Vascular Disease Research</i> , 2019, 16, 13-21.                                      | 0.9  | 3         |
| 45 | Validity of Danish register diagnoses of myocardial infarction and stroke against experts in people with screen-detected diabetes. <i>BMC Public Health</i> , 2019, 19, 228.   | 1.2  | 6         |
| 46 | Long-term effects of intensive multifactorial therapy in individuals with screen-detected type 2 diabetes in primary care: 10-year follow-up of the ADDITION-Europe cluster-randomised trial. <i>Lancet Diabetes and Endocrinology</i> , 2019, 7, 925-937. | 5.5  | 39        |
| 47 | Trajectories of obesity by spousal diabetes status in the English Longitudinal Study of Ageing. <i>Diabetic Medicine</i> , 2019, 36, 105-109.  | 1.2  | 3         |
| 48 | Associations of Mitochondrial and Nuclear Mitochondrial Variants and Genes with Seven Metabolic Traits. <i>American Journal of Human Genetics</i> , 2019, 104, 112-138.  | 2.6  | 106       |
| 49 | Effect of duration and burden of microvascular complications on mortality rate in type 1 diabetes: an observational clinical cohort study. <i>Diabetologia</i> , 2019, 62, 633-643.  | 2.9  | 33        |
| 50 | Netzwerke als transversale Felder. , 2019, , 25-61.  |      | 7         |
| 51 | Der Staat und die gelehrigen Körper. Politologische Aufklärung - Konstruktivistische Perspektiven, 2019, , 211-233.  | 0.4  | 2         |
| 52 | Habitual physical activity is associated with lower fasting and greater glucose-induced GLP-1 response in men. <i>Endocrine Connections</i> , 2019, 8, 1607-1617.  | 0.8  | 5         |
| 53 | Spousal cardiometabolic risk factors and incidence of type 2 diabetes: a prospective analysis from the English Longitudinal Study of Ageing. <i>Diabetologia</i> , 2018, 61, 1572-1580.  | 2.9  | 17        |
| 54 | Risk Factors for Incident Diabetic Polyneuropathy in a Cohort With Screen-Detected Type 2 Diabetes Followed for 13 Years: ADDITION-Denmark. <i>Diabetes Care</i> , 2018, 41, 1068-1075.  | 4.3  | 146       |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Clustering of microvascular complications in Type 1 diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , 2018, 32, 393-399.  | 1.2 | 23        |
| 56 | Can insulin response patterns predict metabolic disease risk in individuals with normal glucose tolerance? Reply to Crofts CAP, Brookler K, Henderson G [letter]. <i>Diabetologia</i> , 2018, 61, 1234-1235.  | 2.9 | 0         |
| 57 | Refining the accuracy of validated target identification through coding variant fine-mapping in type 2 diabetes. <i>Nature Genetics</i> , 2018, 50, 559-571.  | 9.4 | 356       |
| 58 | Risk of Cardiovascular Disease and Death in Individuals With Prediabetes Defined by Different Criteria: The Whitehall II Study. <i>Diabetes Care</i> , 2018, 41, 899-906.   | 4.3 | 116       |
| 59 | Re-analysis of public genetic data reveals a rare X-chromosomal variant associated with type 2 diabetes. <i>Nature Communications</i> , 2018, 9, 321.   | 5.8 | 85        |
| 60 | Evidence of a liver- $\alpha$ cell axis in humans: hepatic insulin resistance attenuates relationship between fasting plasma glucagon and glucagonotropic amino acids. <i>Diabetologia</i> , 2018, 61, 671-680.   | 2.9 | 76        |
| 61 | A Genome-Wide Association Study of Diabetic Kidney Disease in Subjects With Type 2 Diabetes. <i>Diabetes</i> , 2018, 67, 1414-1427.   | 0.3 | 136       |
| 62 | Prevalence of micro- and macrovascular diabetes complications at time of type 2 diabetes diagnosis and associated clinical characteristics: A cross-sectional baseline study of 6958 patients in the Danish DD2 cohort. <i>Journal of Diabetes and Its Complications</i> , 2018, 32, 34-40. | 1.2 | 82        |
| 63 | Glucose patterns during an oral glucose tolerance test and associations with future diabetes, cardiovascular disease and all-cause mortality rate. <i>Diabetologia</i> , 2018, 61, 101-107.   | 2.9 | 43        |
| 64 | Fine-mapping type 2 diabetes loci to single-variant resolution using high-density imputation and islet-specific epigenome maps. <i>Nature Genetics</i> , 2018, 50, 1505-1513.   | 9.4 | 1,331     |
| 65 | First Genome-Wide Association Study of Latent Autoimmune Diabetes in Adults Reveals Novel Insights Linking Immune and Metabolic Diabetes. <i>Diabetes Care</i> , 2018, 41, 2396-2403.   | 4.3 | 99        |
| 66 | Risk Factors for the Presence and Progression of Cardiovascular Autonomic Neuropathy in Type 2 Diabetes: ADDITION-Denmark. <i>Diabetes Care</i> , 2018, 41, 2586-2594.  | 4.3 | 67        |
| 67 | Response to Comment on Andersen et al. Risk-Factor Trajectories Preceding Diabetic Polyneuropathy: ADDITION-Denmark. <i>Diabetes Care</i> 2018;41:1955-1962. <i>Diabetes Care</i> , 2018, 41, e148-e149.  | 4.3 | 0         |
| 68 | The role of physical activity in the development of first cardiovascular disease event: a tree-structured survival analysis of the Danish ADDITION-PRO cohort. <i>Cardiovascular Diabetology</i> , 2018, 17, 126.   | 2.7 | 18        |
| 69 | Pathophysiological Characteristics Underlying Different Glucose Response Curves: A Latent Class Trajectory Analysis From the Prospective EGIR-RISC Study. <i>Diabetes Care</i> , 2018, 41, 1740-1748.   | 4.3 | 52        |
| 70 | Prevalence and geographical distribution of insulin pump therapy in the Central Denmark Region and its association with metabolic parameters. <i>Diabetes Research and Clinical Practice</i> , 2018, 141, 148-155.  | 1.1 | 13        |
| 71 | Risk-Factor Trajectories Preceding Diabetic Polyneuropathy: ADDITION-Denmark. <i>Diabetes Care</i> , 2018, 41, 1955-1962.   | 4.3 | 25        |
| 72 | Common variants in the hERG (KCNH2) voltage-gated potassium channel are associated with altered fasting and glucose-stimulated plasma incretin and glucagon responses. <i>BMC Genetics</i> , 2018, 19, 15.  | 2.7 | 12        |

| #  | ARTICLE  | IF   | CITATIONS |
|----|--|------|-----------|
| 73 | Development of Microvascular Complications and Effect of Concurrent Risk Factors in Type 1 Diabetes: A Multistate Model From an Observational Clinical Cohort Study. <i>Diabetes Care</i> , 2018, 41, 2297-2305.   | 4.3  | 17        |
| 74 | Comparative analysis of the association between 35 frailty scores and cardiovascular events, cancer, and total mortality in an elderly general population in England: An observational study. <i>PLoS Medicine</i> , 2018, 15, e1002543.   | 3.9  | 62        |
| 75 | Protein-altering variants associated with body mass index implicate pathways that control energy intake and expenditure in obesity. <i>Nature Genetics</i> , 2018, 50, 26-41.  | 9.4  | 286       |
| 76 | Rare and low-frequency coding variants alter human adult height. <i>Nature</i> , 2017, 542, 186-190.   | 13.7 | 544       |
| 77 | Nondiabetic Glucometabolic Status and Progression of Aortic Stiffness: The Whitehall II Study. <i>Diabetes Care</i> , 2017, 40, 599-606.   | 4.3  | 33        |
| 78 | Glucose patterns during the OGTT and risk of future diabetes in an urban Indian population: The CARRS study. <i>Diabetes Research and Clinical Practice</i> , 2017, 126, 192-197.  | 1.1  | 22        |
| 79 | Social relations, depressive symptoms, and incident type 2 diabetes mellitus: The English Longitudinal Study of Ageing. <i>Diabetes Research and Clinical Practice</i> , 2017, 126, 86-94.   | 1.1  | 20        |
| 80 | Genetic evidence of a causal effect of insulin resistance on branched-chain amino acid levels. <i>Diabetologia</i> , 2017, 60, 873-878.  | 2.9  | 119       |
| 81 | Trajectories of glycaemia, insulin sensitivity and insulin secretion in South Asian and white individuals before diagnosis of type 2 diabetes: a longitudinal analysis from the Whitehall II cohort study. <i>Diabetologia</i> , 2017, 60, 1252-1260.  | 2.9  | 64        |
| 82 | Does training of general practitioners for intensive treatment of people with screen-detected diabetes have a spillover effect on mortality and cardiovascular morbidity in "at risk" individuals with normoglycaemia? Results from the ADDITION-Denmark cluster-randomised controlled trial. <i>Diabetologia</i> , 2017, 60, 1016-1021. | 2.9  | 5         |
| 83 | Associations between glycaemic deterioration and aortic stiffness and central blood pressure. <i>Journal of Hypertension</i> , 2017, 35, 1832-1840.  | 0.3  | 2         |
| 84 | Pluralizing field analysis: Toward a relational understanding of the field of power. <i>Social Science Information</i> , 2017, 56, 49-73.  | 1.1  | 50        |
| 85 | Effectiveness of Liraglutide and Lixisenatide in the Treatment of Type 2 Diabetes: Real-World Evidence from The Health Improvement Network (THIN) Database in the United Kingdom. <i>Diabetes Therapy</i> , 2017, 8, 417-431.  | 1.2  | 21        |
| 86 | SOS2 and ACP1 Loci Identified through Large-Scale Exome Chip Analysis Regulate Kidney Development and Function. <i>Journal of the American Society of Nephrology: JASN</i> , 2017, 28, 981-994.  | 3.0  | 39        |
| 87 | National study of the prevalence of gestational diabetes mellitus among Danish women from 2004 to 2012. <i>Scandinavian Journal of Public Health</i> , 2017, 45, 811-817.  | 1.2  | 40        |
| 88 | New Blood Pressure-Associated Loci Identified in Meta-Analyses of 475,000 Individuals. <i>Circulation: Cardiovascular Genetics</i> , 2017, 10, .   | 5.1  | 48        |
| 89 | Assessment of time to glucose peak during an oral glucose tolerance test. <i>Clinical Endocrinology</i> , 2017, 87, 879-881.   | 1.2  | 1         |
| 90 | Effect of population screening for type 2 diabetes and cardiovascular risk factors on mortality rate and cardiovascular events: a controlled trial among 1,912,392 Danish adults. <i>Diabetologia</i> , 2017, 60, 2183-2191.   | 2.9  | 35        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 91  | Physical Activity and Improvement of Glycemia in Prediabetes by Different Diagnostic Criteria. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, 3712-3721.   | 1.8 | 14        |
| 92  | Physical Activity Dimensions Associated with Impaired Glucose Metabolism. <i>Medicine and Science in Sports and Exercise</i> , 2017, 49, 2176-2184.  | 0.2 | 8         |
| 93  | Agreement Between 35 Published Frailty Scores in the General Population. <i>American Journal of Epidemiology</i> , 2017, 186, 420-434.   | 1.6 | 193       |
| 94  | Heterogeneity in glucose response curves during an oral glucose tolerance test and associated cardiometabolic risk. <i>Endocrine</i> , 2017, 55, 427-434.  | 1.1 | 21        |
| 95  | Household and familial resemblance in risk factors for type 2 diabetes and related cardiometabolic diseases in rural Uganda: a cross-sectional community sample. <i>BMJ Open</i> , 2017, 7, e015214.   | 0.8 | 6         |
| 96  | Adiponectin, biomarkers of inflammation and changes in cardiac autonomic function: Whitehall II study. <i>Cardiovascular Diabetology</i> , 2017, 16, 153.  | 2.7 | 36        |
| 97  | Die zwei Gesichter der Autonomie. , 2017, , 383-423.   |     | 1         |
| 98  | Does Training and Support of General Practitioners in Intensive Treatment of People with Screen-Detected Diabetes Improve Medication, Morbidity and Mortality in People with Clinically-Diagnosed Diabetes? Investigation of a Spill-Over Effect in a Cluster RCT. <i>PLoS ONE</i> , 2017, 12, e0170697. | 1.1 | 1         |
| 99  | The precarity of critique: Cultures of mistrust and the refusal of justification. <i>Filozofija I Drustvo</i> , 2017, 28, 231-249.   | 0.1 | 1         |
| 100 | Mortality prediction of 35 frailty scores in a 7-years follow-up study in elderly general population. <i>European Journal of Public Health</i> , 2016, 26, .   | 0.1 | 0         |
| 101 | Functional and genetic epidemiological characterisation of the <i>FFAR4</i> ( <i>GPR120</i> ) p.R270H variant in the Danish population. <i>Journal of Medical Genetics</i> , 2016, 53, 616-623.  | 1.5 | 20        |
| 102 | Incidence of register-based diabetes 10 years after a stepwise diabetes screening programme: the ADDITION-Denmark study. <i>Diabetologia</i> , 2016, 59, 989-997.  | 2.9 | 10        |
| 103 | Invasively Measured Aortic Systolic Blood Pressure and Office Systolic Blood Pressure in Cardiovascular Risk Assessment. <i>Hypertension</i> , 2016, 68, 768-774.  | 1.3 | 11        |
| 104 | Insulin Resistance Is Accompanied by Increased Fasting Glucagon and Delayed Glucagon Suppression in Individuals With Normal and Impaired Glucose Regulation. <i>Diabetes</i> , 2016, 65, 3473-3481.  | 0.3 | 137       |
| 105 | Soluble CD163, adiponectin, C-reactive protein and progression of dysglycaemia in individuals at high risk of type 2 diabetes mellitus: the ADDITION-PRO cohort. <i>Diabetologia</i> , 2016, 59, 2467-2476.  | 2.9 | 19        |
| 106 | Biomarkers of subclinical inflammation and increases in glycaemia, insulin resistance and beta-cell function in non-diabetic individuals: the Whitehall II study. <i>European Journal of Endocrinology</i> , 2016, 175, 367-377.   | 1.9 | 52        |
| 107 | Methylglyoxal is associated with changes in kidney function among individuals with screen-detected Type 2 diabetes mellitus. <i>Diabetic Medicine</i> , 2016, 33, 1625-1631.   | 1.2 | 40        |
| 108 | Genome-wide association studies in the Japanese population identify seven novel loci for type 2 diabetes. <i>Nature Communications</i> , 2016, 7, 10531.   | 5.8 | 149       |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 109 | Glucose-Dependent Insulinotropic Polypeptide Is Associated With Lower Low-Density Lipoprotein But Unhealthy Fat Distribution, Independent of Insulin: The ADDITION-PRO Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016, 101, 485-493.       | 1.8 | 46        |
| 110 | Genetic Correlation between Body Fat Percentage and Cardiorespiratory Fitness Suggests Common Genetic Etiology. <i>PLoS ONE</i> , 2016, 11, e0166738.   | 1.1 | 18        |
| 111 | Response to the Letter: Comment on "Abdominal Fat Distribution and Cardiovascular Risk in Men and Women With Different Levels of Glucose Tolerance" by Scheuer S.H., et al. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016, 101, L13-L14.         | 1.8 | 0         |
| 112 | Role of Physical Activity Energy Expenditure versus Estimated Fitness Level in Impaired Glucose Regulation. <i>Medicine and Science in Sports and Exercise</i> , 2015, 47, 675.   | 0.2 | 0         |
| 113 | Heterogeneous effect of gestational weight gain on birth weight: quantile regression analysis from a population-based screening. <i>Annals of Epidemiology</i> , 2015, 25, 133-137.e1.  | 0.9 | 7         |
| 114 | GLP-1 Response to Oral Glucose Is Reduced in Prediabetes, Screen-Detected Type 2 Diabetes, and Obesity and Influenced by Sex: The ADDITION-PRO Study. <i>Diabetes</i> , 2015, 64, 2513-2525.  | 0.3 | 235       |
| 115 | Associations of Objectively Measured Physical Activity and Abdominal Fat Distribution. <i>Medicine and Science in Sports and Exercise</i> , 2015, 47, 983-989.  | 0.2 | 15        |
| 116 | Association of self-perceived body image with body mass index and type 2 diabetes – The ADDITION-PRO study. <i>Preventive Medicine</i> , 2015, 75, 64-69.   | 1.6 | 10        |
| 117 | Impact of intensive treatment on serum methylglyoxal levels among individuals with screen-detected type 2 diabetes: the ADDITION-Denmark study. <i>Acta Diabetologica</i> , 2015, 52, 929-936.  | 1.2 | 8         |
| 118 | Higher Physical Activity Is Associated With Lower Aortic Stiffness but Not With Central Blood Pressure. <i>Medicine (United States)</i> , 2015, 94, e485.   | 0.4 | 19        |
| 119 | Cardiovascular and all-cause mortality in relation to various anthropometric measures of obesity in Europeans. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2015, 25, 295-304.  | 1.1 | 122       |
| 120 | The role of serum methylglyoxal on diabetic peripheral and cardiovascular autonomic neuropathy: the ADDITION Denmark study. <i>Diabetic Medicine</i> , 2015, 32, 778-785.   | 1.2 | 38        |
| 121 | Response to Comment on Fårch et al. GLP-1 Response to Oral Glucose Is Reduced in Prediabetes, Screen-Detected Type 2 Diabetes, and Obesity and Influenced by Sex: The ADDITION-PRO Study. <i>Diabetes</i> 2015;64:2513-2525. <i>Diabetes</i> , 2015, 64, e30-e31. | 0.3 | 1         |
| 122 | Physical activity energy expenditure vs cardiorespiratory fitness level in impaired glucose metabolism. <i>Diabetologia</i> , 2015, 58, 2709-2717.  | 2.9 | 12        |
| 123 | Abdominal Fat Distribution and Cardiovascular Risk in Men and Women With Different Levels of Glucose Tolerance. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, 3340-3347.   | 1.8 | 35        |
| 124 | Relationship Between Insulin Resistance and $\beta$ -Cell Dysfunction in Subphenotypes of Prediabetes and Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, 707-716.  | 1.8 | 41        |
| 125 | Associations between Ultrasound Measures of Abdominal Fat Distribution and Indices of Glucose Metabolism in a Population at High Risk of Type 2 Diabetes: The ADDITION-PRO Study. <i>PLoS ONE</i> , 2015, 10, e0123062.   | 1.1 | 35        |
| 126 | A Combined Analysis of 48 Type 2 Diabetes Genetic Risk Variants Shows No Discriminative Value to Predict Time to First Prescription of a Glucose Lowering Drug in Danish Patients with Screen Detected Type 2 Diabetes. <i>PLoS ONE</i> , 2014, 9, e104837.       | 1.1 | 9         |



| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 127 | Reduction of Specific Circulating Lymphocyte Populations with Metabolic Risk Factors in Patients at Risk to Develop Type 2 Diabetes. <i>PLoS ONE</i> , 2014, 9, e107140.   | 1.1 | 11        |
| 128 | Patterns of Obesity Development before the Diagnosis of Type 2 Diabetes: The Whitehall II Cohort Study. <i>PLoS Medicine</i> , 2014, 11, e1001602.   | 3.9 | 77        |
| 129 | Genetic Determinants of Circulating Interleukin-1 Receptor Antagonist Levels and Their Association With Glycemic Traits. <i>Diabetes</i> , 2014, 63, 4343-4359.  | 0.3 | 40        |
| 130 | Effect of secular trends on age-related trajectories of cardiovascular risk factors: the Whitehall II longitudinal study 1985-2009. <i>International Journal of Epidemiology</i> , 2014, 43, 866-877.  | 0.9 | 27        |
| 131 | Sex differences in glucose and insulin trajectories prior to diabetes diagnosis: the Whitehall II study. <i>Acta Diabetologica</i> , 2014, 51, 315-319.  | 1.2 | 19        |
| 132 | Psychological distress, cardiovascular complications and mortality among people with screen-detected type 2 diabetes: follow-up of the ADDITION-Denmark trial. <i>Diabetologia</i> , 2014, 57, 710-717.  | 2.9 | 39        |
| 133 | Motivational Counseling to Reduce Sitting Time. <i>American Journal of Preventive Medicine</i> , 2014, 47, 576-586.  | 1.6 | 67        |
| 134 | Glycaemic threshold for diabetes-specific retinopathy among individuals from Saudi Arabia, Algeria and Portugal. <i>Diabetes Research and Clinical Practice</i> , 2014, 103, e44-e46.  | 1.1 | 7         |
| 135 | Sex-specific effects of naturally occurring variants in the dopamine receptor D2 locus on insulin secretion and Type 2 diabetes susceptibility. <i>Diabetic Medicine</i> , 2014, 31, 1001-1008.  | 1.2 | 12        |
| 136 | The pro-inflammatory biomarker soluble urokinase plasminogen activator receptor (suPAR) is associated with incident type 2 diabetes among overweight but not obese individuals with impaired glucose regulation: effect modification by smoking and body weight status. <i>Diabetologia</i> , 2013, 56, 1542-1546. | 2.9 | 37        |
| 137 | Effect of time of day and fasting duration on measures of glycaemia: analysis from the Whitehall II Study. <i>Diabetologia</i> , 2013, 56, 294-297.  | 2.9 | 19        |
| 138 | Exome sequencing-driven discovery of coding polymorphisms associated with common metabolic phenotypes. <i>Diabetologia</i> , 2013, 56, 298-310.  | 2.9 | 119       |
| 139 | Association between protein signals and type 2 diabetes incidence. <i>Acta Diabetologica</i> , 2013, 50, 697-704.  | 1.2 | 9         |
| 140 | Studies of association of AGPAT6 variants with type 2 diabetes and related metabolic phenotypes in 12,068 Danes. <i>BMC Medical Genetics</i> , 2013, 14, 113.  | 2.1 | 2         |
| 141 | The frequent UCP2 -866G>A polymorphism protects against insulin resistance and is associated with obesity: a study of obesity and related metabolic traits among 17,636 Danes. <i>International Journal of Obesity</i> , 2013, 37, 175-181.  | 1.6 | 36        |
| 142 | Trajectories of cardiometabolic risk factors before diagnosis of three subtypes of type 2 diabetes: a post-hoc analysis of the longitudinal Whitehall II cohort study. <i>Lancet Diabetes and Endocrinology</i> , 2013, 1, 43-51.  | 5.5 | 87        |
| 143 | Dairy product intake in relation to glucose regulation indices and risk of type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2013, 23, 822-828.  | 1.1 | 72        |
| 144 | Combined Heart Rate and Accelerometer-Assessed Physical Activity Energy Expenditure and Associations With Glucose Homeostasis Markers in a Population at High Risk of Developing Diabetes. <i>Diabetes Care</i> , 2013, 36, 3062-3069.   | 4.3 | 34        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 145 | Impact of early detection and treatment of diabetes on the 6-year prevalence of cardiac autonomic neuropathy in people with screen-detected diabetes: ADDITION-Denmark, a cluster-randomised study. <i>Diabetologia</i> , 2013, 56, 101-108. | 2.9 | 55        |
| 146 | BCG protects against tuberculosis irrespective of HIV status: a matched case-control study in Mwanza, Tanzania: Table A1. <i>Thorax</i> , 2013, 68, 288-289.   | 2.7 | 16        |
| 147 | Causal Relationship between Obesity and Vitamin D Status: Bi-Directional Mendelian Randomization Analysis of Multiple Cohorts. <i>PLoS Medicine</i> , 2013, 10, e1001383.  | 3.9 | 753       |
| 148 | Risk scores for diabetes and impaired glycaemia in the Middle East and North Africa. <i>Diabetic Medicine</i> , 2013, 30, 443-451.   | 1.2 | 14        |
| 149 | Diabetes is a strong predictor of mortality during tuberculosis treatment: a prospective cohort study among tuberculosis patients from Mwanza, Tanzania. <i>Tropical Medicine and International Health</i> , 2013, 18, 822-829.              | 1.0 | 90        |
| 150 | Reproducibility of ultrasonography for assessing abdominal fat distribution in a population at high risk of diabetes. <i>Nutrition and Diabetes</i> , 2013, 3, e82-e82.  | 1.5 | 17        |
| 151 | New loci associated with birth weight identify genetic links between intrauterine growth and adult height and metabolism. <i>Nature Genetics</i> , 2013, 45, 76-82.  | 9.4 | 293       |
| 152 | Solute carrier family 2 member 1 is involved in the development of nonalcoholic fatty liver disease. <i>Hepatology</i> , 2013, 57, 505-514.  | 3.6 | 25        |
| 153 | A Standardized Vascular Disease Health Check in Europe: A Cost-Effectiveness Analysis. <i>PLoS ONE</i> , 2013, 8, e66454.  | 1.1 | 35        |
| 154 | NH2-Terminal Probrain Natriuretic Peptide Is Associated With Diabetes Complications in the EURODIAB Prospective Complications Study. <i>Diabetes Care</i> , 2012, 35, 1931-1936.   | 4.3 | 21        |
| 155 | The role of anthropometric and other predictors for diabetes among urban Tanzanians with tuberculosis. <i>International Journal of Tuberculosis and Lung Disease</i> , 2012, 16, 1680-1685.  | 0.6 | 23        |
| 156 | Adiponectin Trajectories Before Type 2 Diabetes Diagnosis. <i>Diabetes Care</i> , 2012, 35, 2540-2547.   | 4.3 | 48        |
| 157 | Effect of Intensive Multifactorial Treatment Compared With Routine Care on Aortic Stiffness and Central Blood Pressure Among Individuals With Screen-Detected Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 2207-2214.                   | 4.3 | 24        |
| 158 | Severe Hypoglycemia and Cardiovascular Disease Incidence in Type 1 Diabetes. <i>Diabetes Care</i> , 2012, 35, 1598-1604.   | 4.3 | 72        |
| 159 | QTc Interval Prolongation Is Independently Associated With Severe Hypoglycemic Attacks in Type 1 Diabetes From the EURODIAB IDDM Complications Study. <i>Diabetes Care</i> , 2012, 35, 125-127.  | 4.3 | 57        |
| 160 | No Interactions Between Previously Associated 2-Hour Glucose Gene Variants and Physical Activity or BMI on 2-Hour Glucose Levels. <i>Diabetes</i> , 2012, 61, 1291-1296.   | 0.3 | 23        |
| 161 | Apolipoprotein(a) Genetic Sequence Variants Associated With Systemic Atherosclerosis and Coronary Atherosclerotic Burden But Not With Venous Thromboembolism. <i>Journal of the American College of Cardiology</i> , 2012, 60, 722-729.      | 1.2 | 149       |
| 162 | Sleep duration and sleep quality are associated differently with alterations of glucose homeostasis. <i>Diabetic Medicine</i> , 2012, 29, e354-60.   | 1.2 | 65        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 163 | Abdominal obesity has the highest impact on metabolic profile in an overweight African population. <i>Annals of Human Biology</i> , 2012, 39, 530-533.  | 0.4 | 2         |
| 164 | HbA1c, fasting plasma glucose and the prediction of diabetes: Inter99, AusDiab and D.E.S.I.R.. <i>Diabetes Research and Clinical Practice</i> , 2012, 96, 392-399.  | 1.1 | 18        |
| 165 | Glycemic index and glycemic load in relation to glucose intolerance among Greenland's Inuit population. <i>Diabetes Research and Clinical Practice</i> , 2012, 97, 298-305.                                     | 1.1 | 7         |
| 166 | The role of diabetes co-morbidity for tuberculosis treatment outcomes: a prospective cohort study from Mwanza, Tanzania. <i>BMC Infectious Diseases</i> , 2012, 12, 165.  | 1.3 | 26        |
| 167 | The effect of FOXA2rs1209523 on glucose-related phenotypes and risk of type 2 diabetes in Danish individuals. <i>BMC Medical Genetics</i> , 2012, 13, 10.   | 2.1 | 10        |
| 168 | Serum 25(OH)D and Type 2 Diabetes Association in a General Population. <i>Diabetes Care</i> , 2012, 35, 1695-1700.  | 4.3 | 80        |
| 169 | Protocol for ADDITION-PRO: a longitudinal cohort study of the cardiovascular experience of individuals at high risk for diabetes recruited from Danish primary care. <i>BMC Public Health</i> , 2012, 12, 1078. | 1.2 | 45        |
| 170 | Determinants of Aortic Stiffness: 16-Year Follow-Up of the Whitehall II Study. <i>PLoS ONE</i> , 2012, 7, e37165.   | 1.1 | 78        |
| 171 | The PNPLA3 rs738409 G-Allele Associates with Reduced Fasting Serum Triglyceride and Serum Cholesterol in Danes with Impaired Glucose Regulation. <i>PLoS ONE</i> , 2012, 7, e40376.                             | 1.1 | 28        |
| 172 | Diabetes and cancer (1): evaluating the temporal relationship between type 2 diabetes and cancer incidence. <i>Diabetologia</i> , 2012, 55, 1607-1618.  | 2.9 | 191       |
| 173 | Work Stress, Obesity and the Risk of Type 2 Diabetes: Gender-Specific Bidirectional Effect in the Whitehall II Study. <i>Obesity</i> , 2012, 20, 428-433.   | 1.5 | 128       |
| 174 | The role of diabetes on the clinical manifestations of pulmonary tuberculosis. <i>Tropical Medicine and International Health</i> , 2012, 17, 877-883.   | 1.0 | 25        |
| 175 | Diabetes in Greenland and its relationship with urbanization. <i>Diabetic Medicine</i> , 2012, 29, 755-760.   | 1.2 | 53        |
| 176 | Association studies of novel obesity-related gene variants with quantitative metabolic phenotypes in a population-based sample of 6,039 Danish individuals. <i>Diabetologia</i> , 2012, 55, 105-113.            | 2.9 | 32        |
| 177 | Cancer occurrence in Danish diabetic patients: duration and insulin effects. <i>Diabetologia</i> , 2012, 55, 948-958.   | 2.9 | 143       |
| 178 | Antidepressant Medication Use and Risk of Hyperglycemia and Diabetes Mellitus—A Noncausal Association?. <i>Biological Psychiatry</i> , 2011, 70, 978-984.   | 0.7 | 54        |
| 179 | A germline variant in the TP53 polyadenylation signal confers cancer susceptibility. <i>Nature Genetics</i> , 2011, 43, 1098-1103.  | 9.4 | 251       |
| 180 | Genome-Wide Association Identifies Nine Common Variants Associated With Fasting Proinsulin Levels and Provides New Insights Into the Pathophysiology of Type 2 Diabetes. <i>Diabetes</i> , 2011, 60, 2624-2634. | 0.3 | 335       |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 181 | Bioinformatics-Driven Identification and Examination of Candidate Genes for Non-Alcoholic Fatty Liver Disease. PLoS ONE, 2011, 6, e16542.   | 1.1 | 21        |
| 182 | Genome-Wide Association Analysis Identifies Variants Associated with Nonalcoholic Fatty Liver Disease That Have Distinct Effects on Metabolic Traits. PLoS Genetics, 2011, 7, e1001324.   | 1.5 | 796       |
| 183 | The Effect of PCSK1 Variants on Waist, Waist-Hip Ratio and Glucose Metabolism Is Modified by Sex and Glucose Tolerance Status. PLoS ONE, 2011, 6, e23907.   | 1.1 | 23        |
| 184 | Diabetes Is a Risk Factor for Pulmonary Tuberculosis: A Case-Control Study from Mwanza, Tanzania. PLoS ONE, 2011, 6, e24215.  | 1.1 | 96        |
| 185 | Genome-Wide Population-Based Association Study of Extremely Overweight Young Adults – The GOYA Study. PLoS ONE, 2011, 6, e24303.  | 1.1 | 105       |
| 186 | The Birth Weight Lowering C-Allele of rs900400 Near LEKR1 and CCNL1 Associates with Elevated Insulin Release following an Oral Glucose Challenge. PLoS ONE, 2011, 6, e27096.  | 1.1 | 13        |
| 187 | Socioeconomic Status and Reduced Kidney Function in the Whitehall II Study: Role of Obesity and Metabolic Syndrome. American Journal of Kidney Diseases, 2011, 58, 389-397.   | 2.1 | 39        |
| 188 | HbA1c and mean blood glucose show stronger associations with cardiovascular disease risk factors than do postprandial glycaemia or glucose variability in persons with diabetes: the A1C-Derived Average Glucose (ADAG) study. Diabetologia, 2011, 54, 69-72. | 2.9 | 77        |
| 189 | The diabetogenic VPS13C/C2CD4A/C2CD4B rs7172432 variant impairs glucose-stimulated insulin response in 5,722 non-diabetic Danish individuals. Diabetologia, 2011, 54, 789-794.  | 2.9 | 30        |
| 190 | Type 2 diabetes risk allele near CENTD2 is associated with decreased glucose-stimulated insulin release. Diabetologia, 2011, 54, 1052-1056.   | 2.9 | 39        |
| 191 | Homozygous carriers of the G allele of rs4664447 of the glucagon gene (GCG) are characterised by decreased fasting and stimulated levels of insulin, glucagon and glucagon-like peptide (GLP)-1. Diabetologia, 2011, 54, 2820-2831.                           | 2.9 | 16        |
| 192 | The minor C-allele of rs2014355 in ACADSB is associated with reduced insulin release following an oral glucose load. BMC Medical Genetics, 2011, 12, 4.   | 2.1 | 11        |
| 193 | Studies of a genetic variant in HK1 in relation to quantitative metabolic traits and to the prevalence of type 2 diabetes. BMC Medical Genetics, 2011, 12, 99.  | 2.1 | 7         |
| 194 | Estimation of allele frequency and association mapping using next-generation sequencing data. BMC Bioinformatics, 2011, 12, 231.  | 1.2 | 170       |
| 195 | The <i>FOXO3A</i> rs2802292 G-Allele Associates with Improved Peripheral and Hepatic Insulin Sensitivity and Increased Skeletal Muscle- <i>FOXO3A</i> mRNA Expression in Twins. Journal of Clinical Endocrinology and Metabolism, 2011, 96, E119-E124.        | 1.8 | 45        |
| 196 | Associations of Common Genetic Variants With Age-Related Changes in Fasting and Postload Glucose. Diabetes, 2011, 60, 1617-1623.  | 0.3 | 17        |
| 197 | Arterial Stiffness, Physical Function, and Functional Limitation. Hypertension, 2011, 57, 1003-1009.  | 1.3 | 92        |
| 198 | Prevalence of Neuropathy and Peripheral Arterial Disease and the Impact of Treatment in People With Screen-Detected Type 2 Diabetes. Diabetes Care, 2011, 34, 2244-2249.  | 4.3 | 89        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 199 | Natural Selection Affects Multiple Aspects of Genetic Variation at Putatively Neutral Sites across the Human Genome. <i>PLoS Genetics</i> , 2011, 7, e1002326.   | 1.5 | 146       |
| 200 | Studies of the Association of Arg72Pro of Tumor Suppressor Protein p53 with Type 2 Diabetes in a Combined Analysis of 55,521 Europeans. <i>PLoS ONE</i> , 2011, 6, e15813.   | 1.1 | 49        |
| 201 | Implications of Central Obesity-Related Variants in LYPLAL1, NRXN3, MSRA, and TFAP2B on Quantitative Metabolic Traits in Adult Danes. <i>PLoS ONE</i> , 2011, 6, e20640.   | 1.1 | 42        |
| 202 | Studies of Metabolic Phenotypic Correlates of 15 Obesity Associated Gene Variants. <i>PLoS ONE</i> , 2011, 6, e23531.  | 1.1 | 55        |
| 203 | Differential relationship between physical activity and progression to diabetes by glucose tolerance status: the Inter99 Study. <i>Diabetologia</i> , 2010, 53, 70-78.   | 2.9 | 27        |
| 204 | Sex differences in glucose levels: a consequence of physiology or methodological convenience? The Inter99 study. <i>Diabetologia</i> , 2010, 53, 858-865.  | 2.9 | 101       |
| 205 | Real-life glycaemic profiles in non-diabetic individuals with low fasting glucose and normal HbA1c: the A1C-Derived Average Glucose (ADAG) study. <i>Diabetologia</i> , 2010, 53, 1608-1611.   | 2.9 | 52        |
| 206 | Type 2 diabetes risk alleles near ADCY5, CDKAL1 and HHEX-IDE are associated with reduced birthweight. <i>Diabetologia</i> , 2010, 53, 1908-1916.   | 2.9 | 61        |
| 207 | Low birthweight and premature birth are both associated with type 2 diabetes in a random sample of middle-aged Danes. <i>Diabetologia</i> , 2010, 53, 2526-2530.   | 2.9 | 78        |
| 208 | Genome-wide association study identifies a sequence variant within the DAB2IP gene conferring susceptibility to abdominal aortic aneurysm. <i>Nature Genetics</i> , 2010, 42, 692-697.   | 9.4 | 181       |
| 209 | A genome-wide association study in the Japanese population identifies susceptibility loci for type 2 diabetes at UBE2E2 and C2CD4A-C2CD4B. <i>Nature Genetics</i> , 2010, 42, 864-868.   | 9.4 | 245       |
| 210 | Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. <i>Nature Genetics</i> , 2010, 42, 949-960.   | 9.4 | 836       |
| 211 | Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. <i>Nature Genetics</i> , 2010, 42, 937-948.  | 9.4 | 2,634     |
| 212 | Performance of existing risk scores in screening for undiagnosed diabetes: an external validation study. <i>Diabetic Medicine</i> , 2010, 27, 46-53.   | 1.2 | 33        |
| 213 | Do the Joint British Society (JBS2) guidelines on prevention of cardiovascular disease with respect to plasma glucose improve risk stratification in the general population? Prospective cohort study. <i>Diabetic Medicine</i> , 2010, 27, 550-555. | 1.2 | 7         |
| 214 | Comparing risk profiles of individuals diagnosed with diabetes by OGTT and HbA <sub>1c</sub> The Danish Inter99 study. <i>Diabetic Medicine</i> , 2010, 27, 906-910.   | 1.2 | 32        |
| 215 | Moving to an A1C-Based Diagnosis of Diabetes Has a Different Impact on Prevalence in Different Ethnic Groups. <i>Diabetes Care</i> , 2010, 33, 580-582.  | 4.3 | 147       |
| 216 | Associations Between Features of Glucose Exposure and A1C. <i>Diabetes</i> , 2010, 59, 1585-1590.  | 0.3 | 77        |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 217 | Low Peripheral Nerve Conduction Velocities and Amplitudes Are Strongly Related to Diabetic Microvascular Complications in Type 1 Diabetes. <i>Diabetes Care</i> , 2010, 33, 2648-2653.   | 4.3 | 45        |
| 218 | Macrophage inhibitory cytokine-1 is increased in individuals before type 2 diabetes diagnosis but is not an independent predictor of type 2 diabetes: the Whitehall II study. <i>European Journal of Endocrinology</i> , 2010, 162, 913-917. | 1.9 | 62        |
| 219 | Studies of association between LPIN1 variants and common metabolic phenotypes among 17,538 Danes. <i>European Journal of Endocrinology</i> , 2010, 163, 81-87.   | 1.9 | 16        |
| 220 | Incremental Predictive Value of Adding Past Blood Pressure Measurements to the Framingham Hypertension Risk Equation. <i>Hypertension</i> , 2010, 55, 1058-1062.   | 1.3 | 28        |
| 221 | Changes in C-reactive protein levels before type 2 diabetes and cardiovascular death: the Whitehall II study. <i>European Journal of Endocrinology</i> , 2010, 163, 89-95.   | 1.9 | 34        |
| 222 | Antidepressant Use Before and After the Diagnosis of Type 2 Diabetes. <i>Diabetes Care</i> , 2010, 33, 1471-1476.  | 4.3 | 45        |
| 223 | Accelerated Increase in Serum Interleukin-1 Receptor Antagonist Starts 6 Years Before Diagnosis of Type 2 Diabetes. <i>Diabetes</i> , 2010, 59, 1222-1227.   | 0.3 | 117       |
| 224 | New Diagnostic Criteria for Diabetes: Is the Change from Glucose to HbA1c Possible in All Populations?. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010, 95, E333-E336.   | 1.8 | 47        |
| 225 | Do Gene Variants Influencing Adult Adiposity Affect Birth Weight? A Population-Based Study of 24 Loci in 4,744 Danish Individuals. <i>PLoS ONE</i> , 2010, 5, e14190.  | 1.1 | 27        |
| 226 | Hyperglycemia, Type 2 Diabetes, and Depressive Symptoms. <i>Diabetes Care</i> , 2009, 32, 1867-1869.   | 4.3 | 42        |
| 227 | Endothelial Function Predicts Progression of Carotid Intima-Media Thickness. <i>Circulation</i> , 2009, 119, 1005-1012.  | 1.6 | 281       |
| 228 | Anti-HSP60 and Anti-HSP70 Antibody Levels and Diabetes Complications in the EURODIAB Study. <i>Diabetes Care</i> , 2009, 32, 1269-1271.  | 4.3 | 19        |
| 229 | Elevated Levels of the Anti-Inflammatory Interleukin-1 Receptor Antagonist Precede the Onset of Type 2 Diabetes. <i>Diabetes Care</i> , 2009, 32, 421-423.   | 4.3 | 177       |
| 230 | Low Serum Adiponectin Predicts 10-Year Risk of Type 2 Diabetes and HbA1c Independently of Obesity, Lipids, and Inflammation: Whitehall II Study. <i>Hormone and Metabolic Research</i> , 2009, 41, 626-629.                                  | 0.7 | 32        |
| 231 | Anti-HSP60 and Anti-HSP70 antibody levels and micro/ macrovascular complications in type 1 diabetes: the EURODIAB Study. <i>Journal of Internal Medicine</i> , 2009, 266, 527-536.   | 2.7 | 33        |
| 232 | Predictors of future fasting and 2h post-OGTT plasma glucose levels in middle-aged men and women—the Inter99 study. <i>Diabetic Medicine</i> , 2009, 26, 377-383.  | 1.2 | 30        |
| 233 | Psychosocial Stress at Work Doubles the Risk of Type 2 Diabetes in Middle-Aged Women. <i>Diabetes Care</i> , 2009, 32, 2230-2235.  | 4.3 | 183       |
| 234 | Engineering of strained III&#x2013;V heterostructures for high hole mobility. , 2009, , .  |     | 13        |

| #   | ARTICLE   | IF   | CITATIONS |
|-----|---|------|-----------|
| 235 | Trajectories of glycaemia, insulin sensitivity, and insulin secretion before diagnosis of type 2 diabetes: an analysis from the Whitehall II study. <i>Lancet, The</i> , 2009, 373, 2215-2221.  | 6.3  | 692       |
| 236 | The association between n-3 fatty acids in erythrocyte membranes and insulin resistance: The inuit health in transition study. <i>International Journal of Circumpolar Health</i> , 2009, 68, 327-336.  | 0.5  | 38        |
| 237 | The association between father's social class and adult obesity is not explained by educational attainment and an unhealthy lifestyle in adulthood. <i>European Journal of Epidemiology</i> , 2008, 23, 573-579.                                    | 2.5  | 28        |
| 238 | Serum Heat Shock Protein 27 and Diabetes Complications in the EURODIAB Prospective Complications Study. <i>Diabetes</i> , 2008, 57, 1966-1970.  | 0.3  | 75        |
| 239 | Relationship Between Risk Factors and Mortality in Type 1 Diabetic Patients in Europe. <i>Diabetes Care</i> , 2008, 31, 1360-1366.  | 4.3  | 199       |
| 240 | Inflammation, Insulin Resistance, and Diabetes's Mendelian Randomization Using CRP Haplotypes Points Upstream. <i>PLoS Medicine</i> , 2008, 5, e155.  | 3.9  | 136       |
| 241 | Prospective study of coffee and tea consumption in relation to risk of type 2 diabetes mellitus among men and women: The Whitehall II study. <i>British Journal of Nutrition</i> , 2008, 100, 1046-1053.  | 1.2  | 62        |
| 242 | Dietary patterns and 15-y risks of major coronary events, diabetes, and mortality. <i>American Journal of Clinical Nutrition</i> , 2008, 87, 1414-1421.   | 2.2  | 220       |
| 243 | Reply to K Esposito and D Giugliano. <i>American Journal of Clinical Nutrition</i> , 2008, 88, 1180-1181.   | 2.2  | 1         |
| 244 | Dietary glycemic index and glycemic load are associated with high-density-lipoprotein cholesterol at baseline but not with increased risk of diabetes in the Whitehall II study. <i>American Journal of Clinical Nutrition</i> , 2007, 86, 988-994. | 2.2  | 82        |
| 245 | Relation Between Blood Glucose and Coronary Mortality Over 33 Years in the Whitehall Study. <i>Diabetes Care</i> , 2006, 29, 26-31.   | 4.3  | 137       |
| 246 | Early Involvement of the Spinal Cord in Diabetic Peripheral Neuropathy. <i>Diabetes Care</i> , 2006, 29, 2664-2669.   | 4.3  | 141       |
| 247 | Risk factors for cardiac autonomic neuropathy in type 1 diabetes mellitus. <i>Diabetologia</i> , 2005, 48, 164-171.   | 2.9  | 162       |
| 248 | Excess cardiac mortality on Monday: the importance of gender, age and hospitalisation. <i>European Journal of Epidemiology</i> , 2005, 20, 395-399.   | 2.5  | 16        |
| 249 | A Meta-analysis of excess cardiac mortality on Monday. <i>European Journal of Epidemiology</i> , 2005, 20, 401-406.   | 2.5  | 54        |
| 250 | Vascular Risk Factors and Diabetic Neuropathy. <i>New England Journal of Medicine</i> , 2005, 352, 1925-1927.   | 13.9 | 40        |
| 251 | Measurement of flow-mediated dilatation of the brachial artery is affected by local elastic vessel wall properties in high-risk patients. <i>Atherosclerosis</i> , 2005, 182, 323-330.  | 0.4  | 53        |
| 252 | Vascular Risk Factors and Diabetic Neuropathy. <i>New England Journal of Medicine</i> , 2005, 352, 341-350.   | 13.9 | 1,094     |

| #   | ARTICLE  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 253 | Is the Association Between Flow-Mediated Dilatation and Cardiovascular Risk Limited to Low-Risk Populations?. <i>Journal of the American College of Cardiology</i> , 2005, 45, 1987-1993.                      | 1.2 | 162       |
| 254 | Study of agreement between LDL size as measured by nuclear magnetic resonance and gradient gel electrophoresis. <i>Journal of Lipid Research</i> , 2004, 45, 1069-1076.  | 2.0 | 49        |
| 255 | Soluble intercellular adhesion molecule 1 and flow-mediated dilatation are related to the estimated risk of coronary heart disease independently from each other. <i>Atherosclerosis</i> , 2003, 170, 147-153. | 0.4 | 43        |
| 256 | Cardiovascular mortality in Dutch men during 1996 European football championship: longitudinal population study. <i>BMJ: British Medical Journal</i> , 2000, 321, 1552-1554.                                   | 2.4 | 147       |