

Francesca Belleudi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2764990/publications.pdf>

Version: 2024-02-01

26
papers

6,636
citations

567144

15
h-index

552653

26
g-index

26
all docs

26
docs citations

26
times ranked

15216
citing authors

#	ARTICLE	IF	CITATIONS
1	The FGFR2c/PKC μ Axis Controls MCL-1-Mediated Invasion in Pancreatic Ductal Adenocarcinoma Cells: Perspectives for Innovative Target Therapies. <i>Biomedicines</i> , 2022, 10, 1652.	1.4	1
2	The aberrant expression in epithelial cells of the mesenchymal isoform of FGFR2 controls the negative crosstalk between EMT and autophagy. <i>Journal of Cellular and Molecular Medicine</i> , 2021, 25, 4166-4172.	1.6	5
3	Expression Profile of Fibroblast Growth Factor Receptors, Keratinocyte Differentiation Markers, and Epithelial Mesenchymal Transition-Related Genes in Actinic Keratosis: A Possible Predictive Factor for Malignant Progression?. <i>Biology</i> , 2021, 10, 331.	1.3	6
4	Expression of the E5 Oncoprotein of HPV16 Impacts on the Molecular Profiles of EMT-Related and Differentiation Genes in Ectocervical Low-Grade Lesions. <i>International Journal of Molecular Sciences</i> , 2021, 22, 6534.	1.8	6
5	Guidelines for the use and interpretation of assays for monitoring autophagy (4th) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50,582 Tc 1,430	4.3	1,430
6	Role of FGFR2c and Its PKC μ Downstream Signaling in the Control of EMT and Autophagy in Pancreatic Ductal Adenocarcinoma Cells. <i>Cancers</i> , 2021, 13, 4993.	1.7	4
7	Role of PKC μ in the epithelial-mesenchymal transition induced by FGFR2 isoform switch. <i>Cell Communication and Signaling</i> , 2020, 18, 76.	2.7	11
8	The Aberrant Expression of the Mesenchymal Variant of FGFR2 in the Epithelial Context Inhibits Autophagy. <i>Cells</i> , 2019, 8, 653.	1.8	4
9	Role of Fibroblast Growth Factor Receptor 2b in the Cross Talk between Autophagy and Differentiation: Involvement of Jun N-Terminal Protein Kinase Signaling. <i>Molecular and Cellular Biology</i> , 2018, 38, .	1.1	7
10	Interplay between FGFR2b ϵ -induced autophagy and phagocytosis: role of PLC γ -mediated signalling. <i>Journal of Cellular and Molecular Medicine</i> , 2018, 22, 668-683.	1.6	8
11	Expression of the FGFR2c mesenchymal splicing variant in human keratinocytes inhibits differentiation and promotes invasion. <i>Molecular Carcinogenesis</i> , 2018, 57, 272-283.	1.3	18
12	Tumor-Derived Microvesicles Enhance Cross-Processing Ability of Clinical Grade Dendritic Cells. <i>Frontiers in Immunology</i> , 2018, 9, 2481.	2.2	23
13	Role of FGFR2b expression and signaling in keratinocyte differentiation: sequential involvement of PKC γ and PKC δ . <i>Cell Death and Disease</i> , 2018, 9, 565.	2.7	15
14	Tumor-Derived Microvesicles Modulate Antigen Cross-Processing via Reactive Oxygen Species-Mediated Alkalinization of Phagosomal Compartment in Dendritic Cells. <i>Frontiers in Immunology</i> , 2017, 8, 1179.	2.2	21
15	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). <i>Autophagy</i> , 2016, 12, 1-222.	4.3	4,701
16	Expression of the FGFR2 mesenchymal splicing variant in epithelial cells drives epithelial-mesenchymal transition. <i>Oncotarget</i> , 2016, 7, 5440-5460.	0.8	54
17	HPV16 E5 expression induces switching from FGFR2b to FGFR2c and epithelial \rightarrow mesenchymal transition. <i>International Journal of Cancer</i> , 2015, 137, 61-72.	2.3	47
18	HPV16 E5 deregulates the autophagic process in human keratinocytes. <i>Oncotarget</i> , 2015, 6, 9370-9386.	0.8	38

#	ARTICLE	IF	CITATIONS
19	FGF7/KGF regulates autophagy in keratinocytes. <i>Autophagy</i> , 2014, 10, 803-821.	4.3	27
20	HPV16 E5 and KGFR/FGFR2b interplay in differentiating epithelial cells. <i>Oncotarget</i> , 2013, 4, 192-205.	0.8	27
21	The Receptor Tyrosine Kinase FGFR2b/KGFR Controls Early Differentiation of Human Keratinocytes. <i>PLoS ONE</i> , 2011, 6, e24194.	1.1	25
22	Expression and signaling of the tyrosine kinase FGFR2b/KGFR regulates phagocytosis and melanosome uptake in human keratinocytes. <i>FASEB Journal</i> , 2011, 25, 170-181.	0.2	34
23	Internalization and intracellular retention of CD4 are two separate functions of the human immunodeficiency virus type 1 Nef protein. <i>Journal of General Virology</i> , 2007, 88, 3133-3138.	1.3	7
24	Endocytic pathways and biological effects induced by UVB-dependent or ligand-dependent activation of the keratinocyte growth factor receptor. <i>FASEB Journal</i> , 2006, 20, 395-397.	0.2	32
25	Tyrosine 769 of the keratinocyte growth factor receptor is required for receptor signaling but not endocytosis. <i>Biochemical and Biophysical Research Communications</i> , 2005, 327, 523-532.	1.0	26
26	UVB-induced activation and internalization of keratinocyte growth factor receptor. <i>Oncogene</i> , 2003, 22, 2422-2431.	2.6	59