

VÃ©ronique SÃ©bille

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2768626/publications.pdf>

Version: 2024-02-01

111
papers

6,784
citations

147801

31
h-index

62596

80
g-index

114
all docs

114
docs citations

114
times ranked

7339
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Investigation of measurement invariance in longitudinal health-related quality of life in preemptive or previously dialyzed kidney transplant recipients. <i>Quality of Life Research</i> , 2022, 31, 607-620. | 3.1 | 3 |
| 2 | Performance of a Rasch-based method for group comparisons of longitudinal change and response shift at the item level in PRO data: A simulation study. <i>Methods</i> , 2022, 204, 327-339. | 3.8 | 4 |
| 3 | Modeling repeated self-reported outcome data: A continuous-time longitudinal Item Response Theory model. <i>Methods</i> , 2022, 204, 386-395. | 3.8 | 6 |
| 4 | Intermittent theta burst stimulation (iTBS) versus 10 Hz high-frequency repetitive transcranial magnetic stimulation (rTMS) to alleviate treatment-resistant unipolar depression: A randomized controlled trial (THETA-DEP). <i>Brain Stimulation</i> , 2022, 15, 870-880. | 1.6 | 26 |
| 5 | Implementation of French Recommendations for the Prevention and the Treatment of Hospital-acquired Pneumonia: A Cluster-randomized Trial. <i>Clinical Infectious Diseases</i> , 2021, 73, e1601-e1610. | 5.8 | 12 |
| 6 | Response shift in results of patient-reported outcome measures: a commentary to The Response Shiftâ€™in Sync Working Group initiative. <i>Quality of Life Research</i> , 2021, 30, 3299-3308. | 3.1 | 11 |
| 7 | Critical examination of current response shift methods and proposal for advancing new methods. <i>Quality of Life Research</i> , 2021, 30, 3325-3342. | 3.1 | 19 |
| 8 | Higher Negative Self-Reference Level in Patients With Personality Disorders and Suicide Attempt(s) History During Biological Treatment for Major Depressive Disorder: Clinical Implications. <i>Frontiers in Psychology</i> , 2021, 12, 631614. | 2.1 | 6 |
| 9 | Response shift in patient-reported outcomes: definition, theory, and a revised model. <i>Quality of Life Research</i> , 2021, 30, 3309-3322. | 3.1 | 48 |
| 10 | The minimal perceived change: a formal model of the responder definition according to the patientâ€™s meaning of change for patient-reported outcome data analysis and interpretation. <i>BMC Medical Research Methodology</i> , 2021, 21, 128. | 3.1 | 17 |
| 11 | Evaluation of the link between the Guttman errors and response shift at the individual level. <i>Quality of Life Research</i> , 2021, , 1. | 3.1 | 2 |
| 12 | Progress in response-shift research needs diverse approaches. <i>Quality of Life Research</i> , 2021, 30, 3365-3366. | 3.1 | 5 |
| 13 | ABO Blood Group Incompatibility Protects Against SARS-CoV-2 Transmission. <i>Frontiers in Microbiology</i> , 2021, 12, 799519. | 3.5 | 23 |
| 14 | Comparison of structural equation modelling, item response theory and Rasch measurement theory-based methods for response shift detection at item level: A simulation study. <i>Statistical Methods in Medical Research</i> , 2020, 29, 1015-1029. | 1.5 | 16 |
| 15 | Longitudinal associations between coping strategies, locus of control and health-related quality of life in patients with breast cancer or melanoma. <i>Quality of Life Research</i> , 2020, 29, 1271-1279. | 3.1 | 12 |
| 16 | Comparison of longitudinal quality of life outcomes in preemptive and dialyzed patients on waiting list for kidney transplantation. <i>Quality of Life Research</i> , 2020, 29, 959-970. | 3.1 | 8 |
| 17 | Reply to letter to editor: â€œRespOnse Shift ALgorithm in item response theory (ROSALI) for response shift detection with missing data in longitudinal patient-reported outcome studiesâ€: <i>Quality of Life Research</i> , 2020, 29, 2611-2612. | 3.1 | 2 |
| 18 | How to Investigate the Effects of Groups on Changes in Longitudinal Patient-Reported Outcomes and Response Shift Using Rasch Models. <i>Frontiers in Psychology</i> , 2020, 11, 613482. | 2.1 | 5 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Depression Reappraisal and Treatment Effect: Will Response Shift Help Improve the Estimation of Treatment Efficacy in Trials for Mood Disorders?. <i>Frontiers in Psychiatry</i> , 2019, 10, 420. | 2.6 | 8 |
| 20 | Tolerability of high-dose ceftriaxone in CNS infections: a prospective multicentre cohort study. <i>Journal of Antimicrobial Chemotherapy</i> , 2019, 74, 1078-1085. | 3.0 | 21 |
| 21 | High-Dose Ceftriaxone for Bacterial Meningitis and Optimization of Administration Scheme Based on Nomogram. <i>Antimicrobial Agents and Chemotherapy</i> , 2019, 63, . | 3.2 | 21 |
| 22 | Differential Item Functioning (DIF) in composite health measurement scale: Recommendations for characterizing DIF with meaningful consequences within the Rasch model framework. <i>PLoS ONE</i> , 2019, 14, e0215073. | 2.5 | 31 |
| 23 | Clinical and economic aspects of newborn screening for severe combined immunodeficiency: DEPISTREC study results. <i>Clinical Immunology</i> , 2019, 202, 33-39. | 3.2 | 47 |
| 24 | A comparison of posttraumatic growth changes in breast cancer and melanoma.. <i>Health Psychology</i> , 2019, 38, 878-887. | 1.6 | 12 |
| 25 | A systematic review of the quality of reporting of simulation studies about methods for the analysis of complex longitudinal patient-reported outcomes data. <i>Quality of Life Research</i> , 2018, 27, 2507-2516. | 3.1 | 5 |
| 26 | Esmolol indirectly stimulates vagal nerve activity in endotoxemic pigs. <i>Intensive Care Medicine Experimental</i> , 2018, 6, 14. | 1.9 | 4 |
| 27 | A mini-review of quality of life as an outcome in prostate cancer trials: patient-centered approaches are needed to propose appropriate treatments on behalf of patients. <i>Health and Quality of Life Outcomes</i> , 2018, 16, 40. | 2.4 | 8 |
| 28 | Extubation Success Prediction in a Multicentric Cohort of Patients with Severe Brain Injury. <i>Anesthesiology</i> , 2017, 127, 338-346. | 2.5 | 73 |
| 29 | A multi-faceted strategy to reduce ventilation-associated mortality in brain-injured patients. The BI-VILI project: a nationwide quality improvement project. <i>Intensive Care Medicine</i> , 2017, 43, 957-970. | 8.2 | 81 |
| 30 | Does the relationship between health-related quality of life and subjective well-being change over time? An exploratory study among breast cancer patients. <i>Social Science and Medicine</i> , 2017, 174, 96-103. | 3.8 | 24 |
| 31 | Identifying patterns of adaptation in breast cancer patients with cancer-related fatigue using response shift analyses at subgroup level. <i>Cancer Medicine</i> , 2017, 6, 2562-2575. | 2.8 | 22 |
| 32 | Efficacy of intermittent Theta Burst Stimulation (iTBS) and 10-Hz high-frequency repetitive transcranial magnetic stimulation (rTMS) in treatment-resistant unipolar depression: study protocol for a randomised controlled trial. <i>Trials</i> , 2017, 18, 17. | 1.6 | 34 |
| 33 | Sustained quality of life improvement after intracoronary injection of autologous bone marrow cells in the setting of acute myocardial infarction: results from the BONAMI trial. <i>Quality of Life Research</i> , 2017, 26, 121-125. | 3.1 | 11 |
| 34 | Association between continuous hyperosmolar therapy and survival in patients with traumatic brain injury â a multicentre prospective cohort study and systematic review. <i>Critical Care</i> , 2017, 21, 328. | 5.8 | 43 |
| 35 | Partial Credit Model: Estimations and Tests of Fit with Pmodel. <i>The Stata Journal</i> , 2016, 16, 464-481. | 2.2 | 5 |
| 36 | Multicentre study for validation of the French addictovigilance network reports assessment tool. <i>British Journal of Clinical Pharmacology</i> , 2016, 82, 1030-1039. | 2.4 | 4 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | The Guttman errors as a tool for response shift detection at subgroup and item levels. <i>Quality of Life Research</i> , 2016, 25, 1385-1393. | 3.1 | 8 |
| 38 | Baclofen to prevent agitation in alcohol-addicted patients in the ICU: study protocol for a randomised controlled trial. <i>Trials</i> , 2016, 17, 415. | 1.6 | 14 |
| 39 | Prospective, multicenter, controlled study of quality of life, psychological adjustment process and medical outcomes of patients receiving a preemptive kidney transplant compared to a similar population of recipients after a dialysis period of less than three years – The PreKit-QoL study protocol. <i>BMC Nephrology</i> , 2016, 17, 11. | 1.8 | 18 |
| 40 | Rasch-family models are more valuable than score-based approaches for analysing longitudinal patient-reported outcomes with missing data. <i>Statistical Methods in Medical Research</i> , 2016, 25, 2067-2087. | 1.5 | 16 |
| 41 | Changes in quality of life after a diagnosis of cancer: a 2-year study comparing breast cancer and melanoma patients. <i>Quality of Life Research</i> , 2016, 25, 1969-1979. | 3.1 | 18 |
| 42 | Empiric antimicrobial therapy for ventilator-associated pneumonia after brain injury. <i>European Respiratory Journal</i> , 2016, 47, 1219-1228. | 6.7 | 25 |
| 43 | Power and sample size determination for group comparison of patient-reported outcomes using polytomous Rasch models. <i>Statistics in Medicine</i> , 2015, 34, 2444-2455. | 1.6 | 4 |
| 44 | On Comparison of Clustering Methods for Pharmacoepidemiological Data. <i>Journal of Biopharmaceutical Statistics</i> , 2015, 25, 843-856. | 0.8 | 7 |
| 45 | Overall performance of Oort's procedure for response shift detection at item level: a pilot simulation study. <i>Quality of Life Research</i> , 2015, 24, 1799-1807. | 3.1 | 6 |
| 46 | Semantic primes theory may be helpful in designing questionnaires such as to prevent response shift. <i>Journal of Clinical Epidemiology</i> , 2015, 68, 646-654. | 5.0 | 9 |
| 47 | Method variation in the impact of missing data on response shift detection. <i>Quality of Life Research</i> , 2015, 24, 521-528. | 3.1 | 5 |
| 48 | RespOnse Shift ALgorithm in Item response theory (ROSALI) for response shift detection with missing data in longitudinal patient-reported outcome studies. <i>Quality of Life Research</i> , 2015, 24, 553-564. | 3.1 | 37 |
| 49 | Power and sample size determination for the group comparison of patient-reported outcomes using the Rasch model: impact of a misspecification of the parameters. <i>BMC Medical Research Methodology</i> , 2015, 15, 21. | 3.1 | 3 |
| 50 | Assessment of score- and Rasch-based methods for group comparison of longitudinal patient-reported outcomes with intermittent missing data (informative and non-informative). <i>Quality of Life Research</i> , 2015, 24, 19-29. | 3.1 | 14 |
| 51 | Estimation of Parameters of the Rasch Model and Comparison of Groups in Presence of Locally Dependent Items. <i>Journal of Applied Measurement</i> , 2015, 16, 268-77. | 0.3 | 1 |
| 52 | Power and Sample Size Determination in the Rasch Model: Evaluation of the Robustness of a Numerical Method to Non-Normality of the Latent Trait. <i>PLoS ONE</i> , 2014, 9, e83652. | 2.5 | 17 |
| 53 | Multilevel modelling of survey data: impact of the two-level weights used in the pseudolikelihood. <i>Journal of Applied Statistics</i> , 2014, 41, 716-732. | 1.3 | 2 |
| 54 | Sample size used to validate a scale: a review of publications on newly-developed patient reported outcomes measures. <i>Health and Quality of Life Outcomes</i> , 2014, 12, 176. | 2.4 | 601 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 55 | The minimal clinically important difference determined using item response theory models: an attempt to solve the issue of the association with baseline score. <i>Journal of Clinical Epidemiology</i> , 2014, 67, 433-440. | 5.0 | 23 |
| 56 | Source contributions of lead in residential floor dust and within-home variability of dust lead loading. <i>Science of the Total Environment</i> , 2014, 470-471, 768-779. | 8.0 | 23 |
| 57 | Hydrocortisone and fludrocortisone for prevention of hospital-acquired pneumonia in patients with severe traumatic brain injury (Corti-TC): a double-blind, multicentre phase 3, randomised placebo-controlled trial. <i>Lancet Respiratory Medicine</i> , 2014, 2, 706-716. | 10.7 | 98 |
| 58 | A simple ratio-based approach for power and sample size determination for 2-group comparison using Rasch models. <i>BMC Medical Research Methodology</i> , 2014, 14, 87. | 3.1 | 1 |
| 59 | Risk factors for prolonged duration of mechanical ventilation in acute traumatic tetraplegic patientsâa retrospective cohort study. <i>Journal of Critical Care</i> , 2014, 29, 313.e7-313.e13. | 2.2 | 17 |
| 60 | Implementation of an Evidence-based Extubation Readiness Bundle in 499 Brain-injured Patients. A BeforeâAfter Evaluation of a Quality Improvement Project. <i>American Journal of Respiratory and Critical Care Medicine</i> , 2013, 188, 958-966. | 5.6 | 75 |
| 61 | Plasma PCSK9 Is a Late Biomarker of Severity in Patients With Severe Trauma Injury. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, E732-E736. | 3.6 | 32 |
| 62 | Discrimination of psychotropic drugs overâconsumers using a threshold exceedance based approach. <i>Statistical Analysis and Data Mining</i> , 2013, 6, 91-101. | 2.8 | 3 |
| 63 | Power and Sample Size Determination for the Group Comparison of Patient-Reported Outcomes with Rasch Family Models. <i>PLoS ONE</i> , 2013, 8, e57279. | 2.5 | 9 |
| 64 | Etomidate increases susceptibility to pneumonia in trauma patients. <i>Intensive Care Medicine</i> , 2012, 38, 1673-1682. | 8.2 | 37 |
| 65 | Metabolic Syndrome in French HIV-Infected Patients: Prevalence and Predictive Factors After 3 Years of Antiretroviral Therapy. <i>AIDS Research and Human Retroviruses</i> , 2012, 28, 1672-1678. | 1.1 | 27 |
| 66 | Investigation of the impact of HLA-DPB1 matching status in 10/10 HLA matched unrelated hematopoietic stem cell transplantation: Results of a French single center study. <i>Human Immunology</i> , 2012, 73, 711-714. | 2.4 | 7 |
| 67 | Lead contamination in French children's homes and environment. <i>Environmental Research</i> , 2012, 116, 58-65. | 7.5 | 37 |
| 68 | Quality of Life and Coping of Women Treated for Breast Cancer and Their Caregiver. What are the Interactions?. <i>Journal of Clinical Psychology in Medical Settings</i> , 2012, 19, 320-328. | 1.4 | 12 |
| 69 | Towards power and sample size calculations for the comparison of two groups of patients with item response theory models. <i>Statistics in Medicine</i> , 2012, 31, 1277-1290. | 1.6 | 11 |
| 70 | Biases and Power for Groups Comparison on Subjective Health Measurements. <i>PLoS ONE</i> , 2012, 7, e44695. | 2.5 | 8 |
| 71 | Continuous controlled-infusion of hypertonic saline solution in traumatic brain-injured patients: a 9-year retrospective study. <i>Critical Care</i> , 2011, 15, R260. | 5.8 | 67 |
| 72 | Nonparametric Item Response Theory Using Stata. <i>The Stata Journal</i> , 2011, 11, 30-51. | 2.2 | 60 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Ã©valuation prÃ©liminaire de la validitÃ© de construit d'un dispositif cartographique (carte procÃ©durale) pour l'Ã©tude du raisonnement clinique. PÃ©dagogique MÃ©dicale, 2011, 12, 87-101. | 0.1 | 1 |
| 74 | Pharmacoepidemiological characterization of psychotropic drugs consumption using a latent class analysis. International Clinical Psychopharmacology, 2011, 26, 54-62. | 1.7 | 17 |
| 75 | Early rise in circulating endothelial protein C receptor correlates with poor outcome in severe sepsis. Intensive Care Medicine, 2011, 37, 950-956. | 8.2 | 24 |
| 76 | Effects of esmolol on systemic and pulmonary hemodynamics and on oxygenation in pigs with hypodynamic endotoxin shock. Intensive Care Medicine, 2011, 37, 1344-1351. | 8.2 | 64 |
| 77 | Imputation by the mean score should be avoided when validating a Patient Reported Outcomes questionnaire by a Rasch model in presence of informative missing data. BMC Medical Research Methodology, 2011, 11, 105. | 3.1 | 17 |
| 78 | Corticotherapy for traumatic brain-injured Patients - The Corti-TC trial: study protocol for a randomized controlled trial. Trials, 2011, 12, 228. | 1.6 | 9 |
| 79 | Comparison of CTT and Rasch-based approaches for the analysis of longitudinal Patient Reported Outcomes. Statistics in Medicine, 2011, 30, 825-838. | 1.6 | 24 |
| 80 | Hydrocortisone Therapy for Patients With Multiple Trauma. JAMA - Journal of the American Medical Association, 2011, 305, 1201. | 7.4 | 175 |
| 81 | Rural environment and risk factors of amyotrophic lateral sclerosis: a case-control study. Journal of Neurology, 2010, 257, 792-798. | 3.6 | 41 |
| 82 | Methodological issues regarding power of classical test theory (CTT) and item response theory (IRT)-based approaches for the comparison of patient-reported outcomes in two groups of patients - a simulation study. BMC Medical Research Methodology, 2010, 10, 24. | 3.1 | 46 |
| 83 | Analysis of the peripheral T-cell repertoire in kidney transplant patients. European Journal of Immunology, 2010, 40, 3280-3290. | 2.9 | 30 |
| 84 | Exploring how students think: a new method combining think-aloud and concept mapping protocols. Medical Education, 2010, 44, 926-935. | 2.1 | 43 |
| 85 | CpG-ODN and MPLA Prevent Mortality in a Murine Model of Post-Hemorrhage-Staphylococcus aureus Pneumonia. PLoS ONE, 2010, 5, e13228. | 2.5 | 34 |
| 86 | Phenotypic and Functional Analyses of KIR3DL1+and KIR3DS1+NK Cell Subsets Demonstrate Differential Regulation by Bw4 Molecules and Induced KIR3DS1 Expression on Stimulated NK Cells. Journal of Immunology, 2009, 182, 6727-6735. | 0.8 | 30 |
| 87 | Use of 10-point analogue scales to estimate dietary intake: A prospective study in patients nutritionally at-risk. Clinical Nutrition, 2009, 28, 134-140. | 5.0 | 79 |
| 88 | Autologous and allogeneic HLA KIR ligand environments and activating KIR control KIR NK cell functions. European Journal of Immunology, 2008, 38, 3474-3486. | 2.9 | 42 |
| 89 | Inhibition of the interaction between the SARS-CoV Spike protein and its cellular receptor by anti-histo-blood group antibodies. Glycobiology, 2008, 18, 1085-1093. | 2.5 | 306 |
| 90 | Incidence and Prognosis of Sustained Arrhythmias in Critically Ill Patients. American Journal of Respiratory and Critical Care Medicine, 2008, 178, 20-25. | 5.6 | 174 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 91 | Sequential Analysis of Quality-of-Life Measurements Using Mixed Rasch Models. , 2008, , 333-347. | | 0 |
| 92 | Sequential analysis of latent variables using mixed-effect latent variable models: Impact of non-informative and informative missing data. <i>Statistics in Medicine</i> , 2007, 26, 4889-4904. | 1.6 | 6 |
| 93 | Sequential Analysis of Quality of Life Measurements with the Mixed Partial Credit Model. , 2007, , 109-125. | | 2 |
| 94 | Sequential Analysis of Quality of Life Rasch Measurements. , 2006, , 421-439. | | 2 |
| 95 | Exploring the Role of Etomidate in Septic Shock and Acute Respiratory Distress Syndrome. <i>Critical Care Medicine</i> , 2006, 34, 1858-1859. | 0.9 | 3 |
| 96 | Effect of low doses of corticosteroids in septic shock patients with or without early acute respiratory distress syndrome*. <i>Critical Care Medicine</i> , 2006, 34, 22-30. | 0.9 | 303 |
| 97 | Prevention of acquired infections in intubated patients with the combination of two decontamination regimens. <i>Critical Care Medicine</i> , 2005, 33, 307-314. | 0.9 | 95 |
| 98 | Impact of a misspecification of the response rate under standard treatment in sequential clinical trials. <i>Fundamental and Clinical Pharmacology</i> , 2005, 19, 569-578. | 1.9 | 1 |
| 99 | Effect of preoperative oral sustained-release morphine sulfate on postoperative morphine requirements in elective spine surgery. <i>Fundamental and Clinical Pharmacology</i> , 2004, 18, 709-714. | 1.9 | 6 |
| 100 | Sequential methods and group sequential designs for comparative clinical trials. <i>Fundamental and Clinical Pharmacology</i> , 2003, 17, 505-516. | 1.9 | 32 |
| 101 | Effect of Treatment With Low Doses of Hydrocortisone and Fludrocortisone on Mortality in Patients With Septic Shock. <i>JAMA - Journal of the American Medical Association</i> , 2002, 288, 862. | 7.4 | 2,698 |
| 102 | Reduced encephalopathy in pigs with ischemia-induced acute hepatic failure treated with a bioartificial liver containing alginate-entrapped hepatocytes. <i>Critical Care Medicine</i> , 2002, 30, 658-663. | 0.9 | 22 |
| 103 | Comparison of the Two-Sided Single Triangular Test to the Double Triangular Test. <i>Contemporary Clinical Trials</i> , 2001, 22, 503-514. | 1.9 | 18 |
| 104 | Comparison of four sequential methods allowing for early stopping of comparative clinical trials. <i>Clinical Science</i> , 2000, 98, 569-578. | 4.3 | 16 |
| 105 | Comparison of four sequential methods allowing for early stopping of comparative clinical trials. <i>Clinical Science</i> , 2000, 98, 569. | 4.3 | 9 |
| 106 | Compartmentalised inducible nitric-oxide synthase activity in septic shock. <i>Lancet, The</i> , 2000, 355, 1143-1148. | 18.7 | 143 |
| 107 | Safety and efficacy of ritonavir and saquinavir in combination with zidovudine and lamivudine. <i>Clinical Pharmacology and Therapeutics</i> , 1999, 65, 661-671. | 4.7 | 8 |
| 108 | Methodological Issues in Pharmacokinetic-Pharmacodynamic Modelling. <i>Clinical Pharmacokinetics</i> , 1998, 35, 151-166. | 3.5 | 42 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Modeling the Spread of Resistant Nosocomial Pathogens in an Intensive-Care Unit. Infection Control and Hospital Epidemiology, 1997, 18, 84-92. | 1.8 | 86 |
| 110 | A Computer Simulation Model for the Spread of Nosocomial Infections Caused by Multidrug-Resistant Pathogens. Journal of Biomedical Informatics, 1997, 30, 307-322. | 0.7 | 34 |
| 111 | Modeling the Spread of Resistant Nosocomial Pathogens in an Intensive-Care Unit. Infection Control and Hospital Epidemiology, 1997, 18, 84-92. | 1.8 | 60 |