## Yong-Giun Kim

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/2785903/publications.pdf

Version: 2024-02-01

759233 44 501 12 citations h-index papers

20 g-index 44 44 44 806 docs citations times ranked citing authors all docs

752698

#	Article	IF	CITATIONS
1	Triglyceride glucose index is a useful marker for predicting subclinical coronary artery disease in the absence of traditional risk factors. Lipids in Health and Disease, 2020, 19, 7.	3.0	69
2	Association between insulin resistance, hyperglycemia, and coronary artery disease according to the presence of diabetes. Scientific Reports, 2019, 9, 6129.	3.3	65
3	Aortopathy and bicuspid aortic valve: haemodynamic burden is main contributor to aortic dilatation. Heart, 2012, 98, 1822-1827.	2.9	45
4	Trends, Characteristics, and Clinical Outcomes of Patients Undergoing Percutaneous Coronary Intervention in Korea between 2011 and 2015. Korean Circulation Journal, 2018, 48, 310.	1.9	28
5	Metabolic Syndrome and the Risk of New-Onset Atrial Fibrillation in Middle-Aged East Asian Men. Circulation Journal, 2018, 82, 1763-1769.	1.6	23
6	Influence of Diabetes Mellitus on Long-Term (Five-Year) Outcomes of Drug-Eluting Stents and Coronary Artery Bypass Grafting for Multivessel Coronary Revascularization. American Journal of Cardiology, 2012, 109, 1548-1557.	1.6	22
7	Independent role of low-density lipoprotein cholesterol in subclinical coronary atherosclerosis in the absence of traditional cardiovascular risk factors. European Heart Journal Cardiovascular Imaging, 2019, 20, 866-872.	1.2	22
8	Association between nonâ€alcoholic fatty liver disease and risk of newâ€onset atrial fibrillation in healthy adults. Liver International, 2020, 40, 338-346.	3.9	18
9	Higher fatty liver index is associated with increased risk of new onset heart failure in healthy adults: a nationwide population-based study in Korea. BMC Cardiovascular Disorders, 2020, 20, 204.	1.7	18
10	Second coupling interval of nonsustained ventricular tachycardia to distinguish malignant from benign outflow tract ventricular tachycardias. Heart Rhythm, 2014, 11, 2222-2230.	0.7	17
11	Serum uric acid level and subclinical coronary atherosclerosis in asymptomatic individuals: An observational cohort study. Atherosclerosis, 2019, 288, 112-117.	0.8	16
12	Prediabetes is not a risk factor for subclinical coronary atherosclerosis. International Journal of Cardiology, 2017, 243, 479-484.	1.7	14
13	A Close Relationship between Non-Alcoholic Fatty Liver Disease Marker and New-Onset Hypertension in Healthy Korean Adults. Korean Circulation Journal, 2020, 50, 695.	1.9	14
14	Comparison between angiotensin-converting enzyme inhibitor and angiotensin receptor blocker after percutaneous coronary intervention. International Journal of Cardiology, 2020, 306, 35-41.	1.7	12
15	Impact of diabetes mellitus in patients undergoing contemporary percutaneous coronary intervention: Results from a Korean nationwide study. PLoS ONE, 2018, 13, e0208746.	2.5	11
16	Moderate-intensity versus high-intensity statin therapy in Korean patients with angina undergoing percutaneous coronary intervention with drug-eluting stents: A propensity-score matching analysis. PLoS ONE, 2018, 13, e0207889.	2.5	9
17	High-density lipoprotein cholesterol and the risk of obstructive coronary artery disease beyond low-density lipoprotein cholesterol in non-diabetic individuals. European Journal of Preventive Cardiology, 2020, 27, 706-714.	1.8	9
18	Beta-blockers provide a differential survival benefit in patients with coronary artery disease undergoing contemporary post-percutaneous coronary intervention management. Scientific Reports, 2020, 10, 22121.	3.3	9

#	Article	IF	CITATIONS
19	Association between serum phosphorus and subclinical coronary atherosclerosis in asymptomatic Korean individuals without kidney dysfunction. American Journal of Clinical Nutrition, 2020, 112, 66-73.	4.7	9
20	Lipoprotein(a) and subclinical coronary atherosclerosis in asymptomatic individuals. Atherosclerosis, 2022, 349, 190-195.	0.8	8
21	Characteristics of early repolarization parameters and prognostic implications in the general ambulatory Korean population. International Journal of Cardiology, 2017, 227, 571-576.	1.7	7
22	Right bundle branch block–type wide QRS complex tachycardia with a reversed R/S complex in lead V6: Development and validation of electrocardiographic differentiation criteria. Heart Rhythm, 2021, 18, 181-188.	0.7	7
23	Relationship between the optimal cut-off values of anthropometric indices for predicting metabolic syndrome and carotid intima-medial thickness in a Korean population. Medicine (United States), 2019, 98, e17620.	1.0	6
24	Association of gamma-glutamyl transferase with subclinical coronary atherosclerosis and cardiac outcomes in non-alcoholics. Scientific Reports, 2020, 10, 17994.	3.3	5
25	A Case of Perimembranous Ventricular Septal Defect Associated with Sinus of Valsalva Aneurysm Mimicking Membranous Septal Aneurysm. Journal of Cardiovascular Imaging, 2015, 23, 113.	0.8	4
26	High HDL-C levels reduce the risk of obstructive coronary artery disease in asymptomatic diabetics who achieved optimal glycemic control. Scientific Reports, 2019, 9, 15306.	3.3	4
27	Impact of fasciculoventricular bypass tracts on the diagnosis and treatment of concomitant arrhythmias and cardiac diseases. Journal of Electrocardiology, 2019, 55, 34-40.	0.9	4
28	Homocysteine is not a risk factor for subclinical coronary atherosclerosis in asymptomatic individuals. PLoS ONE, 2020, 15, e0231428.	2.5	4
29	Association between nonalcoholic fatty liver disease and cardiovascular disease revealed after comprehensive control of metabolic risk factors: a nationwide population-based study in Korea. European Journal of Gastroenterology and Hepatology, 2021, 33, e375-e382.	1.6	4
30	Radiofrequency catheter ablation for drug-refractory paroxysmal atrial fibrillation in a patient with Ebstein's anomaly. HeartRhythm Case Reports, 2017, 3, 109-111.	0.4	3
31	Pre-existing depression in patients with coronary artery disease undergoing percutaneous coronary intervention. Scientific Reports, 2021, 11, 8600.	3.3	3
32	Cholesterol Control for Subclinical Coronary Atherosclerosis in Subjects Without Indication for Statin Therapy. American Journal of Cardiology, 2021, 153, 51-57.	1.6	3
33	Angiotensin-Converting Enzyme Inhibitor–based Versus Angiotensin Receptor Blocker–based Optimal Medical Therapy After Percutaneous Coronary Intervention: A Nationwide Cohort Study. Journal of Cardiovascular Pharmacology, 2021, 77, 61-68.	1.9	2
34	Ticagrelor versus prasugrel in patients with acute myocardial infarction. International Journal of Cardiology, 2021, 344, 25-30.	1.7	2
35	Changes in Carotid Intima-media Thickness and Left Ventricular Mass by Control of Blood Pressure and Hyperlipidemia in Hypertensive Patients. Journal of the Korean Society of Hypertension, 2011, 17, 177.	0.2	1
36	Effect of atrial fibrillation in Asian patients undergoing percutaneous coronary intervention with drug-eluting stents for stable coronary artery disease. Medicine (United States), 2018, 97, e13488.	1.0	1

3

#	Article	IF	CITATIONS
37	Hospital-level variation in follow-up strategies after percutaneous coronary intervention, revealed in health claims data of Korea. Scientific Reports, 2021, 11, 3322.	3.3	1
38	Comparison of new implantation of cardiac implantable electronic device between tertiary and non-tertiary hospitals: a Korean nationwide study. Scientific Reports, 2021, 11, 3705.	3.3	1
39	Frequency and Significance of Right Bundle Branch Block and Subclinical Coronary Atherosclerosis in Asymptomatic Individuals. American Journal of Cardiology, 2021, 158, 30-36.	1.6	1
40	Left Ventricular Apical Aneurysm: Atypical Feature of Cardiac Sarcoidosis Diagnosed by Multimodality Imaging. Korean Circulation Journal, 2022, 52, 169.	1.9	0
41	Title is missing!. , 2020, 15, e0231428.		O
42	Title is missing!. , 2020, 15, e0231428.		0
43	Title is missing!. , 2020, 15, e0231428.		O
44	Title is missing!. , 2020, 15, e0231428.		0