

Yong-Giun Kim

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2785903/publications.pdf>

Version: 2024-02-01

44
papers

501
citations

759233

12
h-index

752698

20
g-index

44
all docs

44
docs citations

44
times ranked

806
citing authors

#	ARTICLE	IF	CITATIONS
1	Lipoprotein(a) and subclinical coronary atherosclerosis in asymptomatic individuals. <i>Atherosclerosis</i> , 2022, 349, 190-195.	0.8	8
2	Left Ventricular Apical Aneurysm: Atypical Feature of Cardiac Sarcoidosis Diagnosed by Multimodality Imaging. <i>Korean Circulation Journal</i> , 2022, 52, 169.	1.9	0
3	Right bundle branch blockâ€”type wide QRS complex tachycardia with a reversed R/S complex in lead V6: Development and validation of electrocardiographic differentiation criteria. <i>Heart Rhythm</i> , 2021, 18, 181-188.	0.7	7
4	Angiotensin-Converting Enzyme Inhibitorâ€”based Versus Angiotensin Receptor Blockerâ€”based Optimal Medical Therapy After Percutaneous Coronary Intervention: A Nationwide Cohort Study. <i>Journal of Cardiovascular Pharmacology</i> , 2021, 77, 61-68.	1.9	2
5	Hospital-level variation in follow-up strategies after percutaneous coronary intervention, revealed in health claims data of Korea. <i>Scientific Reports</i> , 2021, 11, 3322.	3.3	1
6	Association between nonalcoholic fatty liver disease and cardiovascular disease revealed after comprehensive control of metabolic risk factors: a nationwide population-based study in Korea. <i>European Journal of Gastroenterology and Hepatology</i> , 2021, 33, e375-e382.	1.6	4
7	Comparison of new implantation of cardiac implantable electronic device between tertiary and non-tertiary hospitals: a Korean nationwide study. <i>Scientific Reports</i> , 2021, 11, 3705.	3.3	1
8	Pre-existing depression in patients with coronary artery disease undergoing percutaneous coronary intervention. <i>Scientific Reports</i> , 2021, 11, 8600.	3.3	3
9	Cholesterol Control for Subclinical Coronary Atherosclerosis in Subjects Without Indication for Statin Therapy. <i>American Journal of Cardiology</i> , 2021, 153, 51-57.	1.6	3
10	Frequency and Significance of Right Bundle Branch Block and Subclinical Coronary Atherosclerosis in Asymptomatic Individuals. <i>American Journal of Cardiology</i> , 2021, 158, 30-36.	1.6	1
11	Ticagrelor versus prasugrel in patients with acute myocardial infarction. <i>International Journal of Cardiology</i> , 2021, 344, 25-30.	1.7	2
12	High-density lipoprotein cholesterol and the risk of obstructive coronary artery disease beyond low-density lipoprotein cholesterol in non-diabetic individuals. <i>European Journal of Preventive Cardiology</i> , 2020, 27, 706-714.	1.8	9
13	Comparison between angiotensin-converting enzyme inhibitor and angiotensin receptor blocker after percutaneous coronary intervention. <i>International Journal of Cardiology</i> , 2020, 306, 35-41.	1.7	12
14	Association between nonâ€”alcoholic fatty liver disease and risk of newâ€”onset atrial fibrillation in healthy adults. <i>Liver International</i> , 2020, 40, 338-346.	3.9	18
15	Association of gamma-glutamyl transferase with subclinical coronary atherosclerosis and cardiac outcomes in non-alcoholics. <i>Scientific Reports</i> , 2020, 10, 17994.	3.3	5
16	Beta-blockers provide a differential survival benefit in patients with coronary artery disease undergoing contemporary post-percutaneous coronary intervention management. <i>Scientific Reports</i> , 2020, 10, 22121.	3.3	9
17	Association between serum phosphorus and subclinical coronary atherosclerosis in asymptomatic Korean individuals without kidney dysfunction. <i>American Journal of Clinical Nutrition</i> , 2020, 112, 66-73.	4.7	9
18	Triglyceride glucose index is a useful marker for predicting subclinical coronary artery disease in the absence of traditional risk factors. <i>Lipids in Health and Disease</i> , 2020, 19, 7.	3.0	69

#	ARTICLE	IF	CITATIONS
19	Higher fatty liver index is associated with increased risk of new onset heart failure in healthy adults: a nationwide population-based study in Korea. <i>BMC Cardiovascular Disorders</i> , 2020, 20, 204.	1.7	18
20	Homocysteine is not a risk factor for subclinical coronary atherosclerosis in asymptomatic individuals. <i>PLoS ONE</i> , 2020, 15, e0231428.	2.5	4
21	A Close Relationship between Non-Alcoholic Fatty Liver Disease Marker and New-Onset Hypertension in Healthy Korean Adults. <i>Korean Circulation Journal</i> , 2020, 50, 695.	1.9	14
22	Title is missing!. , 2020, 15, e0231428.		0
23	Title is missing!. , 2020, 15, e0231428.		0
24	Title is missing!. , 2020, 15, e0231428.		0
25	Title is missing!. , 2020, 15, e0231428.		0
26	Serum uric acid level and subclinical coronary atherosclerosis in asymptomatic individuals: An observational cohort study. <i>Atherosclerosis</i> , 2019, 288, 112-117.	0.8	16
27	High HDL-C levels reduce the risk of obstructive coronary artery disease in asymptomatic diabetics who achieved optimal glycemic control. <i>Scientific Reports</i> , 2019, 9, 15306.	3.3	4
28	Association between insulin resistance, hyperglycemia, and coronary artery disease according to the presence of diabetes. <i>Scientific Reports</i> , 2019, 9, 6129.	3.3	65
29	Independent role of low-density lipoprotein cholesterol in subclinical coronary atherosclerosis in the absence of traditional cardiovascular risk factors. <i>European Heart Journal Cardiovascular Imaging</i> , 2019, 20, 866-872.	1.2	22
30	Impact of fasciculoventricular bypass tracts on the diagnosis and treatment of concomitant arrhythmias and cardiac diseases. <i>Journal of Electrocardiology</i> , 2019, 55, 34-40.	0.9	4
31	Relationship between the optimal cut-off values of anthropometric indices for predicting metabolic syndrome and carotid intima-medial thickness in a Korean population. <i>Medicine (United States)</i> , 2019, 98, e17620.	1.0	6
32	Effect of atrial fibrillation in Asian patients undergoing percutaneous coronary intervention with drug-eluting stents for stable coronary artery disease. <i>Medicine (United States)</i> , 2018, 97, e13488.	1.0	1
33	Moderate-intensity versus high-intensity statin therapy in Korean patients with angina undergoing percutaneous coronary intervention with drug-eluting stents: A propensity-score matching analysis. <i>PLoS ONE</i> , 2018, 13, e0207889.	2.5	9
34	Impact of diabetes mellitus in patients undergoing contemporary percutaneous coronary intervention: Results from a Korean nationwide study. <i>PLoS ONE</i> , 2018, 13, e0208746.	2.5	11
35	Trends, Characteristics, and Clinical Outcomes of Patients Undergoing Percutaneous Coronary Intervention in Korea between 2011 and 2015. <i>Korean Circulation Journal</i> , 2018, 48, 310.	1.9	28
36	Metabolic Syndrome and the Risk of New-Onset Atrial Fibrillation in Middle-Aged East Asian Men. <i>Circulation Journal</i> , 2018, 82, 1763-1769.	1.6	23

#	ARTICLE	IF	CITATIONS
37	Radiofrequency catheter ablation for drug-refractory paroxysmal atrial fibrillation in a patient with Ebstein's anomaly. <i>HeartRhythm Case Reports</i> , 2017, 3, 109-111.	0.4	3
38	Prediabetes is not a risk factor for subclinical coronary atherosclerosis. <i>International Journal of Cardiology</i> , 2017, 243, 479-484.	1.7	14
39	Characteristics of early repolarization parameters and prognostic implications in the general ambulatory Korean population. <i>International Journal of Cardiology</i> , 2017, 227, 571-576.	1.7	7
40	A Case of Perimembranous Ventricular Septal Defect Associated with Sinus of Valsalva Aneurysm Mimicking Membranous Septal Aneurysm. <i>Journal of Cardiovascular Imaging</i> , 2015, 23, 113.	0.8	4
41	Second coupling interval of nonsustained ventricular tachycardia to distinguish malignant from benign outflow tract ventricular tachycardias. <i>Heart Rhythm</i> , 2014, 11, 2222-2230.	0.7	17
42	Aortopathy and bicuspid aortic valve: haemodynamic burden is main contributor to aortic dilatation. <i>Heart</i> , 2012, 98, 1822-1827.	2.9	45
43	Influence of Diabetes Mellitus on Long-Term (Five-Year) Outcomes of Drug-Eluting Stents and Coronary Artery Bypass Grafting for Multivessel Coronary Revascularization. <i>American Journal of Cardiology</i> , 2012, 109, 1548-1557.	1.6	22
44	Changes in Carotid Intima-media Thickness and Left Ventricular Mass by Control of Blood Pressure and Hyperlipidemia in Hypertensive Patients. <i>Journal of the Korean Society of Hypertension</i> , 2011, 17, 177.	0.2	1