

Josã© Nã©lio Januãrio

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2831801/publications.pdf>

Version: 2024-02-01

27
papers

664
citations

687363

13
h-index

552781

26
g-index

29
all docs

29
docs citations

29
times ranked

850
citing authors

#	ARTICLE	IF	CITATIONS
1	Congenital Toxoplasmosis in Southeastern Brazil: Results of Early Ophthalmologic Examination of a Large Cohort of Neonates. <i>Ophthalmology</i> , 2009, 116, 2199-2205.e1.	5.2	126
2	Genetic Characterization of <i>Toxoplasma gondii</i> Revealed Highly Diverse Genotypes for Isolates from Newborns with Congenital Toxoplasmosis in Southeastern Brazil. <i>Journal of Clinical Microbiology</i> , 2013, 51, 901-907.	3.9	120
3	Biomarker Analysis Revealed Distinct Profiles of Innate and Adaptive Immunity in Infants with Ocular Lesions of Congenital Toxoplasmosis. <i>Mediators of Inflammation</i> , 2014, 2014, 1-13.	3.0	47
4	Mortality of children with sickle cell disease: a population study. <i>Jornal De Pediatria</i> , 2010, 86, 279-84.	2.0	40
5	Comparison of HPV genotyping by type-specific PCR and sequencing. <i>Memorias Do Instituto Oswaldo Cruz</i> , 2010, 105, 73-78.	1.6	38
6	Hearing loss in congenital toxoplasmosis detected by newborn screening. <i>Brazilian Journal of Otorhinolaryngology</i> , 2008, 74, 21-28.	1.0	36
7	Geographic distribution of human T-lymphotropic virus types 1 and 2 among mothers of newborns tested during neonatal screening, Minas Gerais, Brazil. <i>Revista Panamericana De Salud Publica/Pan American Journal of Public Health</i> , 2010, 27, 330-7.	1.1	33
8	High prevalence of HTLV-1 and 2 viruses in pregnant women in SÃ©o Luis, state of MaranhÃ©o, Brazil. <i>Revista Da Sociedade Brasileira De Medicina Tropical</i> , 2012, 45, 159-162.	0.9	30
9	Blocking Vertical Transmission of Human T Cell Lymphotropic Virus Type 1 and 2 Through Breastfeeding Interruption. <i>Pediatric Infectious Disease Journal</i> , 2012, 31, 1139-1143.	2.0	29
10	Adverse Socioeconomic Conditions and Oocyst-Related Factors Are Associated with Congenital Toxoplasmosis in a Population-Based Study in Minas Gerais, Brazil. <i>PLoS ONE</i> , 2014, 9, e88588.	2.5	27
11	Real-Time PCR as a Prognostic Tool for Human Congenital Toxoplasmosis. <i>Journal of Clinical Microbiology</i> , 2013, 51, 2766-2768.	3.9	19
12	Cytokine Signatures Associated With Early Onset, Active Lesions and Late Cicatricial Events of Retinochoroidal Commitment in Infants With Congenital Toxoplasmosis. <i>Journal of Infectious Diseases</i> , 2016, 213, 1962-1970.	4.0	19
13	Association Between IgG Subclasses Against <i>Toxoplasma gondii</i> and Clinical Signs in Newborns With Congenital Toxoplasmosis. <i>Pediatric Infectious Disease Journal</i> , 2013, 32, 13-16.	2.0	15
14	Early serum biomarker networks in infants with distinct retinochoroidal lesion status of congenital toxoplasmosis. <i>Cytokine</i> , 2017, 95, 102-112.	3.2	10
15	Establishing tools for early diagnosis of congenital toxoplasmosis: Flow cytometric IgG avidity assay as a confirmatory test for neonatal screening. <i>Journal of Immunological Methods</i> , 2017, 451, 37-47.	1.4	9
16	Early diagnosis of congenital toxoplasmosis in newborn infants using IgG subclasses against two <i>Toxoplasma gondii</i> recombinant proteins. <i>Memorias Do Instituto Oswaldo Cruz</i> , 2012, 107, 342-347.	1.6	9
17	High Frequency of Bone Marrow Depression During Congenital Toxoplasmosis Therapy in a Cohort of Children Identified by Neonatal Screening in Minas Gerais, Brazil. <i>Pediatric Infectious Disease Journal</i> , 2017, 36, 1169-1176.	2.0	8
18	Putative biomarkers for early diagnosis and prognosis of congenital ocular toxoplasmosis. <i>Scientific Reports</i> , 2020, 10, 16757.	3.3	7

#	ARTICLE	IF	CITATIONS
19	IgA and IgG1 reactivities assessed by flow cytometry mirror clinical aspects of infants with ocular congenital toxoplasmosis. <i>Journal of Immunological Methods</i> , 2016, 428, 1-8.	1.4	4
20	Frequency of biotinidase gene variants and incidence of biotinidase deficiency in the Newborn Screening Program in Minas Gerais, Brazil. <i>Journal of Medical Screening</i> , 2020, 27, 115-120.	2.3	4
21	Dried blood spots as a practical and inexpensive source for human immunodeficiency virus and hepatitis C virus surveillance. <i>Memorias Do Instituto Oswaldo Cruz</i> , 2005, 100, 265-370.	1.6	4
22	Epidemiological profile of <i>Trypanosoma cruzi</i> -infected mothers and live birth conditions in the state of Minas Gerais, Brazil. <i>Revista Da Sociedade Brasileira De Medicina Tropical</i> , 2013, 46, 196-199.	0.9	2
23	Reply to Gómez Marín. <i>Journal of Infectious Diseases</i> , 2016, 214, 656.2-656.	4.0	2
24	Prognostic value of left ventricular longitudinal strain by speckle-tracking echocardiography in patients with sickle cell disease. <i>International Journal of Cardiovascular Imaging</i> , 2020, 36, 2145-2153.	1.5	1
25	Ação de fatores enzimáticos na obtenção de hidrolisados proteicos de farinha de trigo com baixo teor de fenilalanina. <i>Brazilian Journal of Pharmaceutical Sciences</i> , 2009, 45, 93-98.	1.2	1
26	Comparison between Enzyme Immunoassays Performed on Samples of Dried Blood and Serum for Toxoplasmosis Prenatal Screening: Population-based Study. <i>Revista Brasileira De Ginecologia E Obstetria</i> , 2021, 43, 351-356.	0.8	0
27	Long-term impact of congenital toxoplasmosis on phenotypic and functional features of circulating leukocytes from infants one year after treatment onset. <i>Clinical Immunology</i> , 2021, 232, 108859.	3.2	0