

Kaijun Niu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2839138/publications.pdf>

Version: 2024-02-01

199
papers

3,931
citations

117625

34
h-index

214800

47
g-index

202
all docs

202
docs citations

202
times ranked

5626
citing authors

#	ARTICLE	IF	CITATIONS
1	Blood Neutrophil to Lymphocyte Ratio as a Predictor of Hypertension. <i>American Journal of Hypertension</i> , 2015, 28, 1339-1346.	2.0	106
2	Prevalence and Factors Associated With Sarcopenia in Suburb-dwelling Older Chinese Using the Asian Working Group for Sarcopenia Definition. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2016, 71, 529-535.	3.6	102
3	Effect of beta-hydroxy-beta-methylbutyrate supplementation on muscle loss in older adults: A systematic review and meta-analysis. <i>Archives of Gerontology and Geriatrics</i> , 2015, 61, 168-175.	3.0	101
4	Prospective associations of social isolation and loneliness with poor sleep quality in older adults. <i>Quality of Life Research</i> , 2018, 27, 683-691.	3.1	89
5	Global, regional and national burden of gout, 1990–2017: a systematic analysis of the Global Burden of Disease Study. <i>Rheumatology</i> , 2020, 59, 1529-1538.	1.9	89
6	Dietary long-chain n-3 fatty acids of marine origin and serum C-reactive protein concentrations are associated in a population with a diet rich in marine products. <i>American Journal of Clinical Nutrition</i> , 2006, 84, 223-229.	4.7	84
7	Aspiration pneumonia induces muscle atrophy in the respiratory, skeletal, and swallowing systems. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2018, 9, 643-653.	7.3	77
8	The relationship between thyroid function and the prevalence of type 2 diabetes mellitus in euthyroid subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, jc.2016-2965.	3.6	74
9	Estimated daily quercetin intake and association with the prevalence of type 2 diabetes mellitus in Chinese adults. <i>European Journal of Nutrition</i> , 2019, 58, 819-830.	3.9	72
10	Growth differentiation factor 15 is positively associated with incidence of diabetes mellitus: the Malmö Diet and Cancer Cardiovascular Cohort. <i>Diabetologia</i> , 2019, 62, 78-86.	6.3	71
11	The impact of the 2011 Great East Japan Earthquake on hospitalisation for respiratory disease in a rapidly aging society: a retrospective descriptive and cross-sectional study at the disaster base hospital in Ishinomaki. <i>BMJ Open</i> , 2013, 3, e000865.	1.9	60
12	Timed Up and Go Test can predict recurrent falls: a longitudinal study of the community-dwelling elderly in China. <i>Clinical Interventions in Aging</i> , 2017, Volume 12, 2009-2016.	2.9	58
13	A meta-analysis of the reproducibility of food frequency questionnaires in nutritional epidemiological studies. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2021, 18, 12.	4.6	56
14	The increased risk of sarcopenia in patients with cardiovascular risk factors in Suburb-Dwelling older Chinese using the AWGS definition. <i>Scientific Reports</i> , 2017, 7, 9592.	3.3	55
15	An inverted J-shaped association of serum uric acid with muscle strength among Japanese adult men: a cross-sectional study. <i>BMC Musculoskeletal Disorders</i> , 2013, 14, 258.	1.9	52
16	Neutrophil:lymphocyte ratio is positively related to type 2 diabetes in a large-scale adult population: a Tianjin Chronic Low-Grade Systemic Inflammation and Health cohort study. <i>European Journal of Endocrinology</i> , 2015, 173, 217-225.	3.7	52
17	Dietary patterns are associated with depressive symptoms among Chinese adults: a case-control study with propensity score matching. <i>European Journal of Nutrition</i> , 2017, 56, 2577-2587.	3.9	52
18	Soft drink consumption is associated with depressive symptoms among adults in China. <i>Journal of Affective Disorders</i> , 2015, 172, 422-427.	4.1	50

#	ARTICLE	IF	CITATIONS
19	A Distinct Regulatory Role of Th17 Cytokines IL-17A and IL-17F in Chemokine Secretion from Lung Microvascular Endothelial Cells. <i>Inflammation</i> , 2012, 35, 1119-1131.	3.8	44
20	Elevated Serum Level of Interleukin 17 in a Population With Prehypertension. <i>Journal of Clinical Hypertension</i> , 2015, 17, 770-774.	2.0	42
21	Handgrip strength is inversely associated with metabolic syndrome and its separate components in middle aged and older adults: a large-scale population-based study. <i>Metabolism: Clinical and Experimental</i> , 2019, 93, 61-67.	3.4	42
22	Ultra-processed food consumption and the risk of non-alcoholic fatty liver disease in the Tianjin Chronic Low-grade Systemic Inflammation and Health Cohort Study. <i>International Journal of Epidemiology</i> , 2022, 51, 237-249.	1.9	42
23	Meta-Analysis of Dexmedetomidine on Emergence Agitation and Recovery Profiles in Children after Sevoflurane Anesthesia: Different Administration and Different Dosage. <i>PLoS ONE</i> , 2015, 10, e0123728.	2.5	41
24	Chronic inflammation, lymphangiogenesis, and effect of an anti-VEGFR therapy in a mouse model and in human patients with aspiration pneumonia. <i>Journal of Pathology</i> , 2015, 235, 632-645.	4.5	41
25	Knee loading inhibits osteoclast lineage in a mouse model of osteoarthritis. <i>Scientific Reports</i> , 2016, 6, 24668.	3.3	41
26	Microbiome of Deep Dentinal Caries from Reversible Pulpitis to Irreversible Pulpitis. <i>Journal of Endodontics</i> , 2019, 45, 302-309.e1.	3.1	41
27	A tomato-rich diet is related to depressive symptoms among an elderly population aged 70 years and over: A population-based, cross-sectional analysis. <i>Journal of Affective Disorders</i> , 2013, 144, 165-170.	4.1	40
28	Comparing the inflammatory profiles for incidence of diabetes mellitus and cardiovascular diseases: a prospective study exploring the "common soil" hypothesis. <i>Cardiovascular Diabetology</i> , 2018, 17, 87.	6.8	40
29	Association between nut consumption and non-alcoholic fatty liver disease in adults. <i>Liver International</i> , 2019, 39, 1732-1741.	3.9	40
30	NUT CONSUMPTION IS ASSOCIATED WITH DEPRESSIVE SYMPTOMS AMONG CHINESE ADULTS. <i>Depression and Anxiety</i> , 2016, 33, 1065-1072.	4.1	39
31	Relationship between grip strength and newly diagnosed nonalcoholic fatty liver disease in a large-scale adult population. <i>Scientific Reports</i> , 2016, 6, 33255.	3.3	38
32	Poor Lower Extremity Function Was Associated with Pre-Diabetes and Diabetes in Older Chinese People. <i>PLoS ONE</i> , 2014, 9, e115883.	2.5	38
33	Coffee treatment prevents the progression of sarcopenia in aged mice in vivo and in vitro. <i>Experimental Gerontology</i> , 2014, 50, 1-8.	2.8	37
34	Consumption of low-fat dairy, but not whole-fat dairy, is inversely associated with depressive symptoms in Japanese adults. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2017, 52, 847-853.	3.1	37
35	A dietary pattern rich in animal organ, seafood and processed meat products is associated with newly diagnosed hyperuricaemia in Chinese adults: a propensity score-matched case-control study. <i>British Journal of Nutrition</i> , 2018, 119, 1177-1184.	2.3	37
36	Association of chronic diseases with depression, anxiety and stress in Chinese general population: The CHCN-BTH cohort study. <i>Journal of Affective Disorders</i> , 2021, 282, 1278-1287.	4.1	37

#	ARTICLE	IF	CITATIONS
37	Irritable Bowel Syndrome Is Positively Related to Metabolic Syndrome: A Population-Based Cross-Sectional Study. <i>PLoS ONE</i> , 2014, 9, e112289.	2.5	37
38	Subnormal Peripheral Blood Leukocyte Counts Are Related to the Lowest Prevalence and Incidence of Metabolic Syndrome: Tianjin Chronic Low-Grade Systemic Inflammation and Health Cohort Study. <i>Mediators of Inflammation</i> , 2014, 2014, 1-12.	3.0	36
39	Relationship between daily isoflavone intake and sleep in Japanese adults: a cross-sectional study. <i>Nutrition Journal</i> , 2015, 14, 127.	3.4	36
40	Royal Jelly Prevents the Progression of Sarcopenia in Aged Mice In Vivo and In Vitro. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2013, 68, 1482-1492.	3.6	34
41	Soft drink consumption and risk of nonalcoholic fatty liver disease: results from the Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIH) cohort study. <i>American Journal of Clinical Nutrition</i> , 2021, 113, 1265-1274.	4.7	34
42	Mobility and Muscle Strength Together are More Strongly Correlated with Falls in Suburb-Dwelling Older Chinese. <i>Scientific Reports</i> , 2016, 6, 25420.	3.3	33
43	Association between habitual yogurt consumption and newly diagnosed non-alcoholic fatty liver disease. <i>European Journal of Clinical Nutrition</i> , 2020, 74, 491-499.	2.9	32
44	Association between Complement C3 and Prevalence of Fatty Liver Disease in an Adult Population: A Cross-Sectional Study from the Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIHealth) Cohort Study. <i>PLoS ONE</i> , 2015, 10, e0122026.	2.5	32
45	Muscle strength rather than muscle mass is associated with osteoporosis in older Chinese adults. <i>Journal of the Formosan Medical Association</i> , 2018, 117, 101-108.	1.7	31
46	Association between bedtime and the prevalence of newly diagnosed non-alcoholic fatty liver disease in adults. <i>Liver International</i> , 2018, 38, 2277-2286.	3.9	31
47	White blood cells count as an indicator to identify whether obesity leads to increased risk of type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018, 141, 140-147.	2.8	30
48	Serum Immunoglobulin M Concentration Is Positively Related to Metabolic Syndrome in an Adult Population: Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIH) Cohort Study. <i>PLoS ONE</i> , 2014, 9, e88701.	2.5	29
49	Serum levels of immunoglobulins in an adult population and their relationship with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2016, 115, 76-82.	2.8	28
50	Relationship Between Grip Strength and Prediabetes in a Large-Scale Adult Population. <i>American Journal of Preventive Medicine</i> , 2019, 56, 844-851.	3.0	28
51	Twelve-Month Incidence of Depressive Symptoms in Suburb-Dwelling Chinese Older Adults: Role of Sarcopenia. <i>Journal of the American Medical Directors Association</i> , 2019, 20, 64-69.	2.5	28
52	Insoluble dietary fibre intake is associated with lower prevalence of newly-diagnosed non-alcoholic fatty liver disease in Chinese men: a large population-based cross-sectional study. <i>Nutrition and Metabolism</i> , 2020, 17, 4.	3.0	28
53	Clinical Relevance of Different Handgrip Strength Indexes and Mobility Limitation in the Elderly Adults. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2016, 71, 96-102.	3.6	27
54	Both muscle mass and muscle strength are inversely associated with depressive symptoms in an elderly Chinese population. <i>International Journal of Geriatric Psychiatry</i> , 2017, 32, 769-778.	2.7	27

#	ARTICLE	IF	CITATIONS
55	The traditional Japanese dietary pattern and longitudinal changes in cardiovascular disease risk factors in apparently healthy Japanese adults. <i>European Journal of Nutrition</i> , 2016, 55, 267-279.	3.9	26
56	Association between dietary raw garlic intake and newly diagnosed nonalcoholic fatty liver disease: a population-based study. <i>European Journal of Endocrinology</i> , 2019, 181, 591-602.	3.7	26
57	Dietary patterns and risk of non-alcoholic fatty liver disease in adults: A prospective cohort study. <i>Clinical Nutrition</i> , 2021, 40, 5373-5382.	5.0	25
58	Grip strength and depressive symptoms in a large-scale adult population: The TCLSIH cohort study. <i>Journal of Affective Disorders</i> , 2021, 279, 222-228.	4.1	24
59	Relationship of anabolic and catabolic biomarkers with muscle strength and physical performance in older adults: a population-based cross-sectional study. <i>BMC Musculoskeletal Disorders</i> , 2015, 16, 202.	1.9	23
60	Vitamin D is related to handgrip strength in adult men aged 50 years and over: A population study from the TCLSIH cohort study. <i>Clinical Endocrinology</i> , 2019, 90, 753-765.	2.4	23
61	Intermediary effect of inflammation on the association between dietary patterns and non-alcoholic fatty liver disease. <i>Nutrition</i> , 2020, 71, 110562.	2.4	23
62	Social Isolation and Loneliness as Risk Factors for Grip Strength Decline Among Older Women and Men in China. <i>Journal of the American Medical Directors Association</i> , 2020, 21, 1926-1930.	2.5	23
63	Associations of long-term exposure to ambient air pollution with cardiac conduction abnormalities in Chinese adults: The CHCN-BTH cohort study. <i>Environment International</i> , 2020, 143, 105981.	10.0	23
64	Dietary Patterns are Associated with Helicobacter Pylori Infection in Chinese Adults: A Cross-Sectional Study. <i>Scientific Reports</i> , 2016, 6, 32334.	3.3	22
65	Independent and combined relationship of habitual unhealthy eating behaviors with depressive symptoms: A prospective study. <i>Journal of Epidemiology</i> , 2017, 27, 42-47.	2.4	22
66	Soft drinks consumption is associated with nonalcoholic fatty liver disease independent of metabolic syndrome in Chinese population. <i>European Journal of Nutrition</i> , 2018, 57, 2113-2121.	3.9	22
67	Relationship between thyroid function and elevated blood pressure in euthyroid adults. <i>Journal of Clinical Hypertension</i> , 2018, 20, 1541-1549.	2.0	21
68	Validity of the food frequency questionnaire for adults in nutritional epidemiological studies: A systematic review and meta-analysis. <i>Critical Reviews in Food Science and Nutrition</i> , 2023, 63, 1670-1688.	10.3	21
69	Association between consumption frequency of honey and non-alcoholic fatty liver disease: results from a cross-sectional analysis based on the Tianjin Chronic Low-grade Systemic Inflammation and Health (TCLSIH) Cohort Study. <i>British Journal of Nutrition</i> , 2021, 125, 712-720.	2.3	20
70	Lower Serum Endogenous Secretory Receptor for Advanced Glycation End Product Level as a Risk Factor of Metabolic Syndrome Among Japanese Adult Men: A 2-Year Longitudinal Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014, 99, 587-593.	3.6	18
71	Low serum creatinine predicts risk for type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2018, 34, e3011.	4.0	18
72	A J-shaped association between soy food intake and depressive symptoms in Chinese adults. <i>Clinical Nutrition</i> , 2018, 37, 1013-1018.	5.0	18

#	ARTICLE	IF	CITATIONS
73	Relationship between consumption of raw garlic and handgrip strength in a large-scale adult population. <i>Clinical Nutrition</i> , 2020, 39, 1234-1241.	5.0	18
74	Comparing the diagnostic ability of inflammatory markers in metabolic syndrome. <i>Clinica Chimica Acta</i> , 2017, 475, 1-6.	1.1	17
75	Effects of protease-treated royal jelly on muscle strength in elderly nursing home residents: A randomized, double-blind, placebo-controlled, dose-response study. <i>Scientific Reports</i> , 2017, 7, 11416.	3.3	17
76	Thyroid Function as a Predictor of Handgrip Strength Among Middle-Aged and Older Euthyroid Adults: The TCLSIH Cohort Study. <i>Journal of the American Medical Directors Association</i> , 2019, 20, 1236-1241.	2.5	17
77	Association between edible mushroom intake and the prevalence of newly diagnosed non-alcoholic fatty liver disease: results from the Tianjin Chronic Low-Grade Systemic Inflammation and Health Cohort Study in China. <i>British Journal of Nutrition</i> , 2020, 123, 104-112.	2.3	17
78	Higher serum soluble receptor for advanced glycation end product levels and lower prevalence of metabolic syndrome among Japanese adult men: a cross-sectional study. <i>Diabetology and Metabolic Syndrome</i> , 2014, 6, 33.	2.7	16
79	Dietary myricetin intake is inversely associated with the prevalence of type 2 diabetes mellitus in a Chinese population. <i>Nutrition Research</i> , 2019, 68, 82-91.	2.9	16
80	Antibiotic resistance and molecular characteristics of methicillin-resistant <i>Staphylococcus epidermidis</i> recovered from hospital personnel in China. <i>Journal of Global Antimicrobial Resistance</i> , 2020, 22, 195-201.	2.2	16
81	Serum C-Reactive Protein Even at Very Low (<1.0 mg/l) Concentration Is Associated with Physical Performance in a Community-Based Elderly Population Aged 70 Years and Over. <i>Gerontology</i> , 2008, 54, 260-267.	2.8	15
82	Association of self-reported sleep duration and hypertension: Results of a Chinese prospective cohort study. <i>Clinical and Experimental Hypertension</i> , 2016, 38, 514-519.	1.3	15
83	Elevated serum complement C3 levels are associated with prehypertension in an adult population. <i>Clinical and Experimental Hypertension</i> , 2017, 39, 42-49.	1.3	15
84	The association between thyroid function and incidence of metabolic syndrome in euthyroid subjects: Tianjin chronic low-grade systemic inflammation and health cohort study. <i>Clinical Endocrinology</i> , 2018, 88, 735-743.	2.4	15
85	Relationship of serum immunoglobulin levels to blood pressure and hypertension in an adult population. <i>Journal of Human Hypertension</i> , 2018, 32, 212-218.	2.2	15
86	Habitual yoghurt consumption and depressive symptoms in a general population study of 19,596 adults. <i>European Journal of Nutrition</i> , 2018, 57, 2621-2628.	3.9	15
87	Does a high intake of green leafy vegetables protect from NAFLD? Evidence from a large population study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 1691-1701.	2.6	15
88	Body Mass Index Is Associated with Physical Performance in Suburb-Dwelling Older Chinese: A Cross-Sectional Study. <i>PLoS ONE</i> , 2015, 10, e0119914.	2.5	14
89	Nutritional Biomarkers, Gene-Diet Interaction, and Risk Factors for Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-2.	2.3	14
90	Consumption of preserved egg, a high-lead-containing food, is strongly associated with depressive symptoms in Chinese adults. <i>British Journal of Nutrition</i> , 2017, 118, 737-742.	2.3	14

#	ARTICLE	IF	CITATIONS
91	The association between missing teeth and nonalcoholic fatty liver disease in adults. <i>Journal of Clinical Periodontology</i> , 2018, 45, 941-951.	4.9	14
92	Distinct associations of computer/mobile devices use and TV watching with depressive symptoms in adults: A large population study in China. <i>Depression and Anxiety</i> , 2019, 36, 879-886.	4.1	14
93	Association between neutrophil to lymphocyte ratio and depressive symptoms among Chinese adults: A population study from the TCLSIH cohort study. <i>Psychoneuroendocrinology</i> , 2019, 103, 76-82.	2.7	14
94	Association between consumption of edible seaweeds and newly diagnosed nonalcoholic fatty liver disease: The TCLSIH Cohort Study. <i>Liver International</i> , 2021, 41, 311-320.	3.9	14
95	The predictive power of physical function assessed by questionnaire and physical performance measures for subsequent disability. <i>Aging Clinical and Experimental Research</i> , 2012, 24, 345-53.	2.9	14
96	Impact of Blunted Perception of Dyspnea on Medical Care Use and Expenditure, and Mortality in Elderly People. <i>Frontiers in Physiology</i> , 2012, 3, 238.	2.8	13
97	Frequent alcohol drinking is associated with lower prevalence of self-reported common cold: a retrospective study. <i>BMC Public Health</i> , 2012, 12, 987.	2.9	13
98	Association between dietary patterns and metabolic syndrome in Chinese adults: a propensity score-matched case-control study. <i>Scientific Reports</i> , 2016, 6, 34748.	3.3	13
99	Consumption of chilies, but not sweet peppers, is positively related to handgrip strength in an adult population. <i>Journal of Nutrition, Health and Aging</i> , 2016, 20, 546-552.	3.3	13
100	The relationship between <i>Helicobacter pylori</i> infection and depressive symptoms in the general population in China: The TCLSIH cohort study. <i>Helicobacter</i> , 2019, 24, e12632.	3.5	13
101	Soy Food Intake Is Inversely Associated with Newly Diagnosed Nonalcoholic Fatty Liver Disease in the TCLSIH Cohort Study. <i>Journal of Nutrition</i> , 2020, 150, 3280-3287.	2.9	13
102	Leg Extension Power Is a Pre-Disaster Modifiable Risk Factor for Post-Traumatic Stress Disorder among Survivors of the Great East Japan Earthquake: A Retrospective Cohort Study. <i>PLoS ONE</i> , 2014, 9, e96131.	2.5	12
103	Does elevated serum uric acid level predict the hypertension incidence? A Chinese prospective cohort study. <i>Clinical and Experimental Hypertension</i> , 2015, 37, 498-504.	1.3	12
104	High serum adiponectin levels predict incident falls among middle-aged and older adults: a prospective cohort study. <i>Age and Ageing</i> , 2016, 45, 366-371.	1.6	12
105	Red cell distribution width is associated with hemoglobin A1C elevation, but not glucose elevation. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1544-1548.	2.3	12
106	The relationship between red blood cell distribution width and atrial fibrillation in Asian population: A cross-sectional study. <i>PACE - Pacing and Clinical Electrophysiology</i> , 2019, 42, 1197-1203.	1.2	12
107	Complement C3 and incident hospitalization due to chronic kidney disease: a population-based cohort study. <i>BMC Nephrology</i> , 2019, 20, 61.	1.8	12
108	Daily dietary isoflavone intake in relation to lowered risk of depressive symptoms among men. <i>Journal of Affective Disorders</i> , 2020, 261, 121-125.	4.1	12

#	ARTICLE	IF	CITATIONS
109	Long-term Tai Chi training is related to depressive symptoms among Tai Chi practitioners. <i>Journal of Affective Disorders</i> , 2014, 169, 36-39.	4.1	11
110	The predictive value of mean serum uric acid levels for developing prediabetes. <i>Diabetes Research and Clinical Practice</i> , 2016, 118, 79-89.	2.8	11
111	Cracked Teeth and Poor Oral Masticatory Habits: A Matched Case-control Study in China. <i>Journal of Endodontics</i> , 2017, 43, 885-889.	3.1	11
112	Dietary seaweed intake and depressive symptoms in Japanese adults: a prospective cohort study. <i>Nutrition Journal</i> , 2019, 18, 58.	3.4	11
113	Adjuvant treatment of coronary heart disease angina pectoris with Chinese patent medicine. <i>Medicine (United States)</i> , 2019, 98, e16884.	1.0	11
114	Association between tooth loss and handgrip strength in a general adult population. <i>PLoS ONE</i> , 2020, 15, e0236010.	2.5	11
115	Association between consumption of ultra-processed foods and hyperuricemia: TCLSIH prospective cohort study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 1993-2003.	2.6	11
116	Dietary patterns and risk for hyperuricemia in the general population: Results from the TCLSIH cohort study. <i>Nutrition</i> , 2022, 93, 111501.	2.4	11
117	The overall computer/mobile devices usage time is related to newly diagnosed non-alcoholic fatty liver disease: a population-based study. <i>Annals of Medicine</i> , 2016, 48, 568-576.	3.8	10
118	Raw orange intake is associated with higher prevalence of non-alcoholic fatty liver disease in an adult population. <i>Nutrition</i> , 2019, 60, 252-260.	2.4	10
119	Low-fat dairy, but not whole-/high-fat dairy, consumption is related with higher serum adiponectin levels in apparently healthy adults. <i>European Journal of Nutrition</i> , 2013, 52, 771-778.	3.9	9
120	High-Normal Thyroid Function Predicts Incident Nonalcoholic Fatty Liver Disease Among Middle-Aged and Older Euthyroid Subjects. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2022, 77, 197-203.	3.6	9
121	Staple Foods Consumption and Irritable Bowel Syndrome in Japanese Adults: A Cross-Sectional Study. <i>PLoS ONE</i> , 2015, 10, e0119097.	2.5	9
122	Association between Serum Ferritin Concentrations and Depressive Symptoms among Chinese Adults: A Population Study from the Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIHealth) Cohort Study. <i>PLoS ONE</i> , 2016, 11, e0162682.	2.5	9
123	Ultra-processed food intake is associated with grip strength decline in middle-aged and older adults: a prospective analysis of the TCLSIH study. <i>European Journal of Nutrition</i> , 2022, 61, 1331-1341.	3.9	9
124	Increased serum ferritin levels are independently associated with carotid atherosclerosis in women. <i>British Journal of Nutrition</i> , 2017, 117, 1623-1630.	2.3	8
125	Daily tea drinking is not associated with newly diagnosed non-alcoholic fatty liver disease in Chinese adults: the Tianjin chronic low-grade systemic inflammation and health cohort study. <i>Nutrition Journal</i> , 2019, 18, 71.	3.4	8
126	Reference values for handgrip strength: data from the TCLSIH cohort study. <i>Age and Ageing</i> , 2019, 49, 233-238.	1.6	8

#	ARTICLE	IF	CITATIONS
127	The associations of self-rated health with cardiovascular risk proteins: a proteomics approach. <i>Clinical Proteomics</i> , 2019, 16, 40.	2.1	8
128	Sugar-containing carbonated beverages consumption is associated with hyperuricemia in general adults: A cross-sectional study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020, 30, 1645-1652.	2.6	8
129	Long-term exposure to ambient nitrogen dioxide and ozone modifies systematic low-grade inflammation: The CHCN-BTH study. <i>International Journal of Hygiene and Environmental Health</i> , 2022, 239, 113875.	4.3	8
130	The association between banana consumption and the depressive symptoms in Chinese general adult population: A cross-sectional study. <i>Journal of Affective Disorders</i> , 2020, 264, 1-6.	4.1	7
131	The relationship between dietary patterns and grip strength in the general population: the TCLSIH cohort study. <i>European Journal of Nutrition</i> , 2021, 60, 2409-2421.	3.9	7
132	Associations between different types and sources of dietary fibre intake and depressive symptoms in a general population of adults: a cross-sectional study. <i>British Journal of Nutrition</i> , 2021, 125, 1281-1290.	2.3	7
133	Edible mushroom consumption and incident hyperuricemia: results from the TCLSIH cohort study. <i>Food and Function</i> , 2021, 12, 9178-9187.	4.6	7
134	Handgrip strength as a predictor of incident hypertension in the middle-aged and older population: The TCLSIH cohort study. <i>Maturitas</i> , 2021, 150, 7-13.	2.4	7
135	Association of long-term exposure to ambient particulate pollution with stage 1 hypertension defined by the 2017 ACC/AHA Hypertension Guideline and cardiovascular disease: The CHCN-BTH cohort study. <i>Environmental Research</i> , 2021, 199, 111356.	7.5	7
136	Dietary fibre intake and risk of prediabetes in China: results from the Tianjin Chronic Low-grade Systemic Inflammation and Health (TCLSIH) Cohort Study. <i>British Journal of Nutrition</i> , 2022, 128, 753-761.	2.3	7
137	Inflammatory potential of diet and risk of nonalcoholic fatty liver disease: a prospective cohort study. <i>European Journal of Clinical Nutrition</i> , 2022, 76, 1125-1132.	2.9	7
138	Cortisol is not the primary mediator for augmented CXCR4 expression on natural killer cells after acute exercise. <i>Journal of Applied Physiology</i> , 2014, 117, 199-204.	2.5	6
139	Combined Effect of Obesity and Mobility Limitation with Incidence of Type 2 Diabetes and Mortality in Chinese Elderly. <i>Rejuvenation Research</i> , 2017, 20, 375-382.	1.8	6
140	Association of dietary patterns with depressive symptoms in Chinese postmenopausal women. <i>British Journal of Nutrition</i> , 2019, 122, 1168-1174.	2.3	6
141	Low-Normal Thyroid Function Predicts Incident Anemia in the General Population With Euthyroid Status. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 5693-5702.	3.6	6
142	The Effects of Mobile-App-Based Low-Carbohydrate Dietary Guidance on Postprandial Hyperglycemia in Adults with Prediabetes. <i>Diabetes Therapy</i> , 2020, 11, 2341-2355.	2.5	6
143	Association between soft drink consumption and handgrip strength in middle aged and older adults: the TCLSIH cohort study. <i>International Journal of Food Sciences and Nutrition</i> , 2020, 71, 856-862.	2.8	6
144	Height predict incident non-alcoholic fatty liver disease among general adult population in Tianjin, China, independent of body mass index, waist circumference, waist-to-height ratio, and metabolic syndrome. <i>BMC Public Health</i> , 2020, 20, 388.	2.9	6

#	ARTICLE	IF	CITATIONS
145	Honey consumption is inversely associated with prediabetes among Chinese adults: results from the Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIH) Cohort Study. <i>British Journal of Nutrition</i> , 2020, 124, 112-119.	2.3	6
146	Association between eating speed and newly diagnosed nonalcoholic fatty liver disease among the general population. <i>Nutrition Research</i> , 2020, 80, 78-88.	2.9	6
147	Effect of Al₂O₃ on Evolution of Oxide Inclusions in Tire Cord Steel during Hot Rolling. <i>ISIJ International</i> , 2021, 61, 2605-2612.	1.4	6
148	The occurrence and risk factors associated with post-traumatic stress disorder among discharged COVID-19 patients in Tianjin, China. <i>Brain and Behavior</i> , 2022, 12, e2492.	2.2	6
149	Adherence to a vegetable dietary pattern attenuates the risk of non-alcoholic fatty liver disease in incident type 2 diabetes: The TCLSIH cohort study. <i>Journal of Internal Medicine</i> , 2022, 291, 469-480.	6.0	6
150	High Tomato and Tomato Product Consumption is Protective Against the Decline in Handgrip Strength Among Japanese Adults: The Oroshiso Study. <i>Journal of Epidemiology</i> , 2018, 28, 397-403.	2.4	5
151	Serum levels of immunoglobulins in an adult population and their relationship with nonalcoholic fatty liver disease. <i>Journal of Digestive Diseases</i> , 2018, 19, 498-507.	1.5	5
152	Relationship between high-sensitivity C-reactive protein and subclinical carotid atherosclerosis stratified by glucose metabolic status in Chinese adults. <i>Clinical Cardiology</i> , 2019, 42, 39-46.	1.8	5
153	Low dietary quercetin intake by food frequency questionnaire analysis is not associated with hypertension occurrence. <i>Clinical Nutrition</i> , 2021, 40, 3748-3753.	5.0	5
154	Association between dietary onion intake and subclinical hypothyroidism in adults: a population-based study from an iodine-replete area. <i>Endocrine</i> , 2021, 74, 616-624.	2.3	5
155	Relationship Between Dietary Patterns and Carotid Atherosclerosis Among People Aged 50 Years or Older: A Population-Based Study in China. <i>Frontiers in Nutrition</i> , 2021, 8, 723726.	3.7	5
156	OUP accepted manuscript. <i>Age and Ageing</i> , 2022, 51, .	1.6	5
157	Longitudinal changes in the relationship between serum adiponectin concentration and cardiovascular risk factors among apparently healthy middle-aged adults. <i>International Journal of Cardiology</i> , 2013, 167, 2318-2320.	1.7	4
158	Response to "The Relation Between Neutrophil-Lymphocyte Ratio and Hypertension". <i>American Journal of Hypertension</i> , 2015, 28, 1387-1387.	2.0	4
159	The association between immunoglobulin concentrations and prediabetes prevalence in a large Chinese cohort. <i>Metabolism: Clinical and Experimental</i> , 2017, 73, 77-84.	3.4	4
160	Association of microalbuminuria with diabetes is stronger in people with prehypertension compared to those with ideal blood pressure. <i>Nephrology</i> , 2018, 23, 690-696.	1.6	4
161	Associations between honey consumption and prehypertension in adults aged 40 years and older. <i>Clinical and Experimental Hypertension</i> , 2020, 42, 420-427.	1.3	4
162	Consumption of chilies and sweet peppers is associated with lower risk of sarcopenia in older adults. <i>Aging</i> , 2021, 13, 9135-9142.	3.1	4

#	ARTICLE	IF	CITATIONS
163	The consumption of wholegrain is related to depressive symptoms among Chinese adults: a cross-sectional study. <i>European Journal of Clinical Nutrition</i> , 2022, 76, 126-133.	2.9	4
164	Environmental and Genetic Determinants of Major Chronic Disease in Beijingâ€“Tianjinâ€“Hebei Region: Protocol for a Community-Based Cohort Study. <i>Frontiers in Public Health</i> , 2021, 9, 659701.	2.7	4
165	Associations of Nutritional, Lifestyle, and Metabolic Factors With Non-alcoholic Fatty Liver Disease: An Umbrella Review With More Than 380,000 Participants. <i>Frontiers in Nutrition</i> , 2021, 8, 642509.	3.7	4
166	Longitudinal Associations Between Hand Grip Strength and Non-Alcoholic Fatty Liver Disease in Adults: A Prospective Cohort Study. <i>Frontiers in Medicine</i> , 2021, 8, 752999.	2.6	4
167	Relationship between the consumption of wholegrain and nonalcoholic fatty liver disease: The TCLSIH cohort study. <i>Clinical Nutrition</i> , 2022, 41, 1483-1490.	5.0	4
168	Sex-Specific Associations Between Complement Component 3 and Component 4 Levels and Metabolic Syndrome in an Adult Population. <i>Metabolic Syndrome and Related Disorders</i> , 2018, 16, 143-149.	1.3	3
169	Relationship between serum levels of immunoglobulins and metabolic syndrome in an adult population: A population study from the TCLSIH cohort study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2019, 29, 916-922.	2.6	3
170	Complex Dietary Topologies in Non-alcoholic Fatty Liver Disease: A Network Science Analysis. <i>Frontiers in Nutrition</i> , 2020, 7, 579086.	3.7	3
171	Thyroid function and lipid profile in euthyroid adults: the TCLSIH cohort study. <i>Endocrine</i> , 2020, 70, 107-114.	2.3	3
172	Serum Immunoglobulin M Concentration Varies with Triglyceride Levels in an Adult Population: Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIHealth) Cohort Study. <i>PLoS ONE</i> , 2015, 10, e0124255.	2.5	3
173	Relationship between seaweeds consumption and hyperuricaemia in general adults: a Population-based study from the Tianjin Chronic Low-grade Systemic Inflammation and Health (TCLSIH) cohort study. <i>British Journal of Nutrition</i> , 2022, 127, 369-376.	2.3	2
174	Predictive Value of Thyroid Hormones for Incident Hyperuricemia in Euthyroid Subjects: The Tianjin Chronic Low-Grade Systemic Inflammation and Health Cohort Study. <i>Endocrine Practice</i> , 2021, 27, 291-297.	2.1	2
175	Higher plain water intake is related to lower newly diagnosed nonalcoholic fatty liver disease risk: a population-based study. <i>European Journal of Clinical Nutrition</i> , 2021, 75, 1801-1808.	2.9	2
176	Consumption of Preserved Egg Is Associated with Modestly Increased Risk of Nonalcoholic Fatty Liver Disease in Chinese Adults. <i>Journal of Nutrition</i> , 2021, 151, 2741-2748.	2.9	2
177	Dietary patterns and sarcopenia in elderly adults: the TCLSIH study. <i>British Journal of Nutrition</i> , 2021, , 1-26.	2.3	2
178	Ultra-processed food consumption and the risk of subclinical thyroid dysfunction: a prospective cohort study. <i>Food and Function</i> , 2022, 13, 3431-3440.	4.6	2
179	The longitudinal associations between sweet potato intake and the risk of non-alcoholic fatty liver disease: the TCLSIH cohort study. <i>International Journal of Food Sciences and Nutrition</i> , 2022, , 1-12.	2.8	2
180	Green tea consumption and risk of depressive symptoms: Results from the TCLSIH Cohort Study. <i>Journal of Affective Disorders</i> , 2022, 310, 183-188.	4.1	2

#	ARTICLE	IF	CITATIONS
181	Raw garlic consumption is inversely associated with prehypertension in a large-scale adult population. <i>Journal of Human Hypertension</i> , 2020, 34, 59-67.	2.2	1
182	Saltwater fish but not freshwater fish consumption is positively related to handgrip strength: The TCLSIH Cohort Study. <i>Nutrition Research</i> , 2021, 90, 46-54.	2.9	1
183	Association between edible mushrooms consumption and handgrip strength: A large-scale population based on the TCLSIH cohort study. <i>Clinical Nutrition</i> , 2022, 41, 1197-1207.	5.0	1
184	Green Tea and Feeling Depressed. , 2013, , 1385-1398.		0
185	Frequent coffee drinking is associated with higher leg extension power among adult men : a cross-sectional study. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2014, 63, 383-389.	0.0	0
186	Response to "Risk factors for developing prediabetes". <i>Diabetes Research and Clinical Practice</i> , 2018, 135, 233-234.	2.8	0
187	Effects of self-monitoring devices on blood pressure in older adults with hypertension and diabetes: a randomised controlled trial. <i>Journal of Epidemiology and Community Health</i> , 2020, 74, 137-143.	3.7	0
188	Response to "Red cell distribution width as an emerging screening tool in atrial fibrillation". <i>PACE - Pacing and Clinical Electrophysiology</i> , 2020, 43, 154-155.	1.2	0
189	The Association between Leukocyte and Its Subtypes and Benign Breast Disease: The TCLSIH Cohort Study. <i>Journal of Oncology</i> , 2020, 2020, 1-7.	1.3	0
190	The association between longitudinal trends of thyroid hormones levels and incident hypertension in a euthyroid population. <i>Journal of Human Hypertension</i> , 2021, 35, 1159-1169.	2.2	0
191	Relative Handgrip Strength But Not Absolute Handgrip Strength is a Determinant of Incident Type 2 Diabetes: A Prospective Cohort Study and Systematic Review with Meta-Analysis. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
192	Light-To-Moderate Raw Garlic Consumption Frequency Is Inversely Associated With Thickened Carotid Intima-Media Thickness: A Population-Based Study. <i>Frontiers in Nutrition</i> , 2021, 8, 648821.	3.7	0
193	Association of Appendicular Skeletal Muscle to Trunk Fat Ratio with Type 2 Diabetes Mellitus in Older Adults. <i>Gerontology</i> , 2021, , 1-8.	2.8	0
194	Non-Linear Associations of Total and Differential Leukocyte Counts with Incident Nonalcoholic Fatty Liver Disease. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
195	Serum Immunoglobulin Concentrations and Risk of Type 2 Diabetes Mellitus in Adults: A Prospective Cohort Study. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
196	Serum Immunoglobulin Concentrations and Risk of Type 2 Diabetes Mellitus in Adults: A Prospective Cohort Study. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
197	Association between Consumption of Edible Seaweeds and Newly Diagnosed Non-Alcohol Fatty Liver Disease: The TCLSIH Cohort Study. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
198	The association between daily yogurt consumption and serum lipid profiles in the general adult population: the TCLSIH cohort study. <i>International Journal of Food Sciences and Nutrition</i> , 2021, , 1-9.	2.8	0

#	ARTICLE	IF	CITATIONS
199	Organ meat consumption and risk of nonalcoholic fatty liver disease: the TCLSIH study. British Journal of Nutrition, 2022, , 1-21.	2.3	0