

Kaijun Niu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2839138/publications.pdf>

Version: 2024-02-01

199
papers

3,931
citations

117625

34
h-index

214800

47
g-index

202
all docs

202
docs citations

202
times ranked

5626
citing authors

#	ARTICLE	IF	CITATIONS
1	Validity of the food frequency questionnaire for adults in nutritional epidemiological studies: A systematic review and meta-analysis. <i>Critical Reviews in Food Science and Nutrition</i> , 2023, 63, 1670-1688.	10.3	21
2	High-Normal Thyroid Function Predicts Incident Nonalcoholic Fatty Liver Disease Among Middle-Aged and Older Euthyroid Subjects. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2022, 77, 197-203.	3.6	9
3	Relationship between seaweeds consumption and hyperuricaemia in general adults: a Population-based study from the Tianjin Chronic Low-grade Systemic Inflammation and Health (TCLSIH) cohort study. <i>British Journal of Nutrition</i> , 2022, 127, 369-376.	2.3	2
4	The consumption of wholegrain is related to depressive symptoms among Chinese adults: a cross-sectional study. <i>European Journal of Clinical Nutrition</i> , 2022, 76, 126-133.	2.9	4
5	Dietary fibre intake and risk of prediabetes in China: results from the Tianjin Chronic Low-grade Systemic Inflammation and Health (TCLSIH) Cohort Study. <i>British Journal of Nutrition</i> , 2022, 128, 753-761.	2.3	7
6	Dietary patterns and risk for hyperuricemia in the general population: Results from the TCLSIH cohort study. <i>Nutrition</i> , 2022, 93, 111501.	2.4	11
7	Ultra-processed food consumption and the risk of non-alcoholic fatty liver disease in the Tianjin Chronic Low-grade Systemic Inflammation and Health Cohort Study. <i>International Journal of Epidemiology</i> , 2022, 51, 237-249.	1.9	42
8	Long-term exposure to ambient nitrogen dioxide and ozone modifies systematic low-grade inflammation: The CHCN-BTH study. <i>International Journal of Hygiene and Environmental Health</i> , 2022, 239, 113875.	4.3	8
9	Ultra-processed food intake is associated with grip strength decline in middle-aged and older adults: a prospective analysis of the TCLSIH study. <i>European Journal of Nutrition</i> , 2022, 61, 1331-1341.	3.9	9
10	OUP accepted manuscript. <i>Age and Ageing</i> , 2022, 51, .	1.6	5
11	The occurrence and risk factors associated with post-traumatic stress disorder among discharged COVID-19 patients in Tianjin, China. <i>Brain and Behavior</i> , 2022, 12, e2492.	2.2	6
12	Inflammatory potential of diet and risk of nonalcoholic fatty liver disease: a prospective cohort study. <i>European Journal of Clinical Nutrition</i> , 2022, 76, 1125-1132.	2.9	7
13	Ultra-processed food consumption and the risk of subclinical thyroid dysfunction: a prospective cohort study. <i>Food and Function</i> , 2022, 13, 3431-3440.	4.6	2
14	Organ meat consumption and risk of nonalcoholic fatty liver disease: the TCLSIH study. <i>British Journal of Nutrition</i> , 2022, , 1-21.	2.3	0
15	The longitudinal associations between sweet potato intake and the risk of non-alcoholic fatty liver disease: the TCLSIH cohort study. <i>International Journal of Food Sciences and Nutrition</i> , 2022, , 1-12.	2.8	2
16	Adherence to a vegetable dietary pattern attenuates the risk of non-alcoholic fatty liver disease in incident type 2 diabetes: The TCLSIH cohort study. <i>Journal of Internal Medicine</i> , 2022, 291, 469-480.	6.0	6
17	Association between edible mushrooms consumption and handgrip strength: A large-scale population based on the TCLSIH cohort study. <i>Clinical Nutrition</i> , 2022, 41, 1197-1207.	5.0	1
18	Green tea consumption and risk of depressive symptoms: Results from the TCLSIH Cohort Study. <i>Journal of Affective Disorders</i> , 2022, 310, 183-188.	4.1	2

#	ARTICLE	IF	CITATIONS
19	Relationship between the consumption of wholegrain and nonalcoholic fatty liver disease: The TCLSIH cohort study. <i>Clinical Nutrition</i> , 2022, 41, 1483-1490.	5.0	4
20	The relationship between dietary patterns and grip strength in the general population: the TCLSIH cohort study. <i>European Journal of Nutrition</i> , 2021, 60, 2409-2421.	3.9	7
21	Association between consumption frequency of honey and non-alcoholic fatty liver disease: results from a cross-sectional analysis based on the Tianjin Chronic Low-grade Systemic Inflammation and Health (TCLSIH) Cohort Study. <i>British Journal of Nutrition</i> , 2021, 125, 712-720.	2.3	20
22	Association between consumption of edible seaweeds and newly diagnosed non-alcohol fatty liver disease: The TCLSIH Cohort Study. <i>Liver International</i> , 2021, 41, 311-320.	3.9	14
23	Associations between different types and sources of dietary fibre intake and depressive symptoms in a general population of adults: a cross-sectional study. <i>British Journal of Nutrition</i> , 2021, 125, 1281-1290.	2.3	7
24	Edible mushroom consumption and incident hyperuricemia: results from the TCLSIH cohort study. <i>Food and Function</i> , 2021, 12, 9178-9187.	4.6	7
25	Grip strength and depressive symptoms in a large-scale adult population: The TCLSIH cohort study. <i>Journal of Affective Disorders</i> , 2021, 279, 222-228.	4.1	24
26	The association between longitudinal trends of thyroid hormones levels and incident hypertension in a euthyroid population. <i>Journal of Human Hypertension</i> , 2021, 35, 1159-1169.	2.2	0
27	A meta-analysis of the reproducibility of food frequency questionnaires in nutritional epidemiological studies. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2021, 18, 12.	4.6	56
28	Soft drink consumption and risk of nonalcoholic fatty liver disease: results from the Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIH) cohort study. <i>American Journal of Clinical Nutrition</i> , 2021, 113, 1265-1274.	4.7	34
29	Light-To-Moderate Raw Garlic Consumption Frequency Is Inversely Associated With Thickened Carotid Intima-Media Thickness: A Population-Based Study. <i>Frontiers in Nutrition</i> , 2021, 8, 648821.	3.7	0
30	Consumption of chilies and sweet peppers is associated with lower risk of sarcopenia in older adults. <i>Aging</i> , 2021, 13, 9135-9142.	3.1	4
31	Association of chronic diseases with depression, anxiety and stress in Chinese general population: The CHCN-BTH cohort study. <i>Journal of Affective Disorders</i> , 2021, 282, 1278-1287.	4.1	37
32	Predictive Value of Thyroid Hormones for Incident Hyperuricemia in Euthyroid Subjects: The Tianjin Chronic Low-Grade Systemic Inflammation and Health Cohort Study. <i>Endocrine Practice</i> , 2021, 27, 291-297.	2.1	2
33	Higher plain water intake is related to lower newly diagnosed nonalcoholic fatty liver disease risk: a population-based study. <i>European Journal of Clinical Nutrition</i> , 2021, 75, 1801-1808.	2.9	2
34	Consumption of Preserved Egg Is Associated with Modestly Increased Risk of Nonalcoholic Fatty Liver Disease in Chinese Adults. <i>Journal of Nutrition</i> , 2021, 151, 2741-2748.	2.9	2
35	Low dietary quercetin intake by food frequency questionnaire analysis is not associated with hypertension occurrence. <i>Clinical Nutrition</i> , 2021, 40, 3748-3753.	5.0	5
36	Environmental and Genetic Determinants of Major Chronic Disease in Beijing-Tianjin-Hebei Region: Protocol for a Community-Based Cohort Study. <i>Frontiers in Public Health</i> , 2021, 9, 659701.	2.7	4

#	ARTICLE	IF	CITATIONS
37	Association between consumption of ultra-processed foods and hyperuricemia: TCLSIH prospective cohort study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 1993-2003.	2.6	11
38	Saltwater fish but not freshwater fish consumption is positively related to handgrip strength: The TCLSIH Cohort Study. <i>Nutrition Research</i> , 2021, 90, 46-54.	2.9	1
39	Association of Appendicular Skeletal Muscle to Trunk Fat Ratio with Type 2 Diabetes Mellitus in Older Adults. <i>Gerontology</i> , 2021, , 1-8.	2.8	0
40	Does a high intake of green leafy vegetables protect from NAFLD? Evidence from a large population study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021, 31, 1691-1701.	2.6	15
41	Association between dietary onion intake and subclinical hypothyroidism in adults: a population-based study from an iodine-replete area. <i>Endocrine</i> , 2021, 74, 616-624.	2.3	5
42	Handgrip strength as a predictor of incident hypertension in the middle-aged and older population: The TCLSIH cohort study. <i>Maturitas</i> , 2021, 150, 7-13.	2.4	7
43	Association of long-term exposure to ambient particulate pollution with stage 1 hypertension defined by the 2017 ACC/AHA Hypertension Guideline and cardiovascular disease: The CHCN-BTH cohort study. <i>Environmental Research</i> , 2021, 199, 111356.	7.5	7
44	Dietary patterns and sarcopenia in elderly adults: the TCLSIH study. <i>British Journal of Nutrition</i> , 2021, , 1-26.	2.3	2
45	Associations of Nutritional, Lifestyle, and Metabolic Factors With Non-alcoholic Fatty Liver Disease: An Umbrella Review With More Than 380,000 Participants. <i>Frontiers in Nutrition</i> , 2021, 8, 642509.	3.7	4
46	Dietary patterns and risk of non-alcoholic fatty liver disease in adults: A prospective cohort study. <i>Clinical Nutrition</i> , 2021, 40, 5373-5382.	5.0	25
47	Effect of Al₂O₃ on Evolution of Oxide Inclusions in Tire Cord Steel during Hot Rolling. <i>ISIJ International</i> , 2021, 61, 2605-2612.	1.4	6
48	Longitudinal Associations Between Hand Grip Strength and Non-Alcoholic Fatty Liver Disease in Adults: A Prospective Cohort Study. <i>Frontiers in Medicine</i> , 2021, 8, 752999.	2.6	4
49	The association between daily yogurt consumption and serum lipid profiles in the general adult population: the TCLSIH cohort study. <i>International Journal of Food Sciences and Nutrition</i> , 2021, , 1-9.	2.8	0
50	Relationship Between Dietary Patterns and Carotid Atherosclerosis Among People Aged 50 Years or Older: A Population-Based Study in China. <i>Frontiers in Nutrition</i> , 2021, 8, 723726.	3.7	5
51	Relationship between consumption of raw garlic and handgrip strength in a large-scale adult population. <i>Clinical Nutrition</i> , 2020, 39, 1234-1241.	5.0	18
52	Raw garlic consumption is inversely associated with prehypertension in a large-scale adult population. <i>Journal of Human Hypertension</i> , 2020, 34, 59-67.	2.2	1
53	Association between edible mushroom intake and the prevalence of newly diagnosed non-alcoholic fatty liver disease: results from the Tianjin Chronic Low-Grade Systemic Inflammation and Health Cohort Study in China. <i>British Journal of Nutrition</i> , 2020, 123, 104-112.	2.3	17
54	Global, regional and national burden of gout, 1990â€“2017: a systematic analysis of the Global Burden of Disease Study. <i>Rheumatology</i> , 2020, 59, 1529-1538.	1.9	89

#	ARTICLE	IF	CITATIONS
55	Daily dietary isoflavone intake in relation to lowered risk of depressive symptoms among men. <i>Journal of Affective Disorders</i> , 2020, 261, 121-125.	4.1	12
56	Intermediary effect of inflammation on the association between dietary patterns and non-alcoholic fatty liver disease. <i>Nutrition</i> , 2020, 71, 110562.	2.4	23
57	Association between habitual yogurt consumption and newly diagnosed non-alcoholic fatty liver disease. <i>European Journal of Clinical Nutrition</i> , 2020, 74, 491-499.	2.9	32
58	Effects of self-monitoring devices on blood pressure in older adults with hypertension and diabetes: a randomised controlled trial. <i>Journal of Epidemiology and Community Health</i> , 2020, 74, 137-143.	3.7	0
59	Response to "Red cell distribution width as an emerging screening tool in atrial fibrillation" PACE - <i>Pacing and Clinical Electrophysiology</i> , 2020, 43, 154-155.	1.2	0
60	The association between banana consumption and the depressive symptoms in Chinese general adult population: A cross-sectional study. <i>Journal of Affective Disorders</i> , 2020, 264, 1-6.	4.1	7
61	Associations between honey consumption and prehypertension in adults aged 40 years and older. <i>Clinical and Experimental Hypertension</i> , 2020, 42, 420-427.	1.3	4
62	Complex Dietary Topologies in Non-alcoholic Fatty Liver Disease: A Network Science Analysis. <i>Frontiers in Nutrition</i> , 2020, 7, 579086.	3.7	3
63	The Association between Leukocyte and Its Subtypes and Benign Breast Disease: The TCLSIH Cohort Study. <i>Journal of Oncology</i> , 2020, 2020, 1-7.	1.3	0
64	Association between tooth loss and handgrip strength in a general adult population. <i>PLoS ONE</i> , 2020, 15, e0236010.	2.5	11
65	Social Isolation and Loneliness as Risk Factors for Grip Strength Decline Among Older Women and Men in China. <i>Journal of the American Medical Directors Association</i> , 2020, 21, 1926-1930.	2.5	23
66	Associations of long-term exposure to ambient air pollution with cardiac conduction abnormalities in Chinese adults: The CHCN-BTH cohort study. <i>Environment International</i> , 2020, 143, 105981.	10.0	23
67	Soy Food Intake Is Inversely Associated with Newly Diagnosed Nonalcoholic Fatty Liver Disease in the TCLSIH Cohort Study. <i>Journal of Nutrition</i> , 2020, 150, 3280-3287.	2.9	13
68	The Effects of Mobile-App-Based Low-Carbohydrate Dietary Guidance on Postprandial Hyperglycemia in Adults with Prediabetes. <i>Diabetes Therapy</i> , 2020, 11, 2341-2355.	2.5	6
69	Sugar-containing carbonated beverages consumption is associated with hyperuricemia in general adults: A cross-sectional study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020, 30, 1645-1652.	2.6	8
70	Association between soft drink consumption and handgrip strength in middle aged and older adults: the TCLSIH cohort study. <i>International Journal of Food Sciences and Nutrition</i> , 2020, 71, 856-862.	2.8	6
71	Height predict incident non-alcoholic fatty liver disease among general adult population in Tianjin, China, independent of body mass index, waist circumference, waist-to-height ratio, and metabolic syndrome. <i>BMC Public Health</i> , 2020, 20, 388.	2.9	6
72	Honey consumption is inversely associated with prediabetes among Chinese adults: results from the Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIH) Cohort Study. <i>British Journal of Nutrition</i> , 2020, 124, 112-119.	2.3	6

#	ARTICLE	IF	CITATIONS
73	Association between eating speed and newly diagnosed nonalcoholic fatty liver disease among the general population. <i>Nutrition Research</i> , 2020, 80, 78-88.	2.9	6
74	Antibiotic resistance and molecular characteristics of methicillin-resistant <i>Staphylococcus epidermidis</i> recovered from hospital personnel in China. <i>Journal of Global Antimicrobial Resistance</i> , 2020, 22, 195-201.	2.2	16
75	Insoluble dietary fibre intake is associated with lower prevalence of newly-diagnosed non-alcoholic fatty liver disease in Chinese men: a large population-based cross-sectional study. <i>Nutrition and Metabolism</i> , 2020, 17, 4.	3.0	28
76	Thyroid function and lipid profile in euthyroid adults: the TCLSIH cohort study. <i>Endocrine</i> , 2020, 70, 107-114.	2.3	3
77	Estimated daily quercetin intake and association with the prevalence of type 2 diabetes mellitus in Chinese adults. <i>European Journal of Nutrition</i> , 2019, 58, 819-830.	3.9	72
78	Dietary myricetin intake is inversely associated with the prevalence of type 2 diabetes mellitus in a Chinese population. <i>Nutrition Research</i> , 2019, 68, 82-91.	2.9	16
79	The relationship between red blood cell distribution width and atrial fibrillation in Asian population: A cross-sectional study. <i>PACE - Pacing and Clinical Electrophysiology</i> , 2019, 42, 1197-1203.	1.2	12
80	Thyroid Function as a Predictor of Handgrip Strength Among Middle-Aged and Older Euthyroid Adults: The TCLSIH Cohort Study. <i>Journal of the American Medical Directors Association</i> , 2019, 20, 1236-1241.	2.5	17
81	Relationship between serum levels of immunoglobulins and metabolic syndrome in an adult population: A population study from the TCLSIH cohort study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2019, 29, 916-922.	2.6	3
82	The relationship between <i>Helicobacter pylori</i> infection and depressive symptoms in the general population in China: The TCLSIH cohort study. <i>Helicobacter</i> , 2019, 24, e12632.	3.5	13
83	Association of dietary patterns with depressive symptoms in Chinese postmenopausal women. <i>British Journal of Nutrition</i> , 2019, 122, 1168-1174.	2.3	6
84	Distinct associations of computer/mobile devices use and TV watching with depressive symptoms in adults: A large population study in China. <i>Depression and Anxiety</i> , 2019, 36, 879-886.	4.1	14
85	Dietary seaweed intake and depressive symptoms in Japanese adults: a prospective cohort study. <i>Nutrition Journal</i> , 2019, 18, 58.	3.4	11
86	Low-Normal Thyroid Function Predicts Incident Anemia in the General Population With Euthyroid Status. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 5693-5702.	3.6	6
87	Daily tea drinking is not associated with newly diagnosed non-alcoholic fatty liver disease in Chinese adults: the Tianjin chronic low-grade systemic inflammation and health cohort study. <i>Nutrition Journal</i> , 2019, 18, 71.	3.4	8
88	Handgrip strength is inversely associated with metabolic syndrome and its separate components in middle aged and older adults: a large-scale population-based study. <i>Metabolism: Clinical and Experimental</i> , 2019, 93, 61-67.	3.4	42
89	Association between nut consumption and non-alcoholic fatty liver disease in adults. <i>Liver International</i> , 2019, 39, 1732-1741.	3.9	40
90	Relationship Between Grip Strength and Prediabetes in a Large-Scale Adult Population. <i>American Journal of Preventive Medicine</i> , 2019, 56, 844-851.	3.0	28

#	ARTICLE	IF	CITATIONS
91	Microbiome of Deep Dentinal Caries from Reversible Pulpitis to Irreversible Pulpitis. <i>Journal of Endodontics</i> , 2019, 45, 302-309.e1.	3.1	41
92	Complement C3 and incident hospitalization due to chronic kidney disease: a population-based cohort study. <i>BMC Nephrology</i> , 2019, 20, 61.	1.8	12
93	Vitamin D is related to handgrip strength in adult men aged 50 years and over: A population study from the TCLSIH cohort study. <i>Clinical Endocrinology</i> , 2019, 90, 753-765.	2.4	23
94	Reference values for handgrip strength: data from the TCLSIH cohort study. <i>Age and Ageing</i> , 2019, 49, 233-238.	1.6	8
95	The associations of self-rated health with cardiovascular risk proteins: a proteomics approach. <i>Clinical Proteomics</i> , 2019, 16, 40.	2.1	8
96	Adjuvant treatment of coronary heart disease angina pectoris with Chinese patent medicine. <i>Medicine (United States)</i> , 2019, 98, e16884.	1.0	11
97	Association between neutrophil to lymphocyte ratio and depressive symptoms among Chinese adults: A population study from the TCLSIH cohort study. <i>Psychoneuroendocrinology</i> , 2019, 103, 76-82.	2.7	14
98	Growth differentiation factor 15 is positively associated with incidence of diabetes mellitus: the Malmö Diet and Cancer Cardiovascular Cohort. <i>Diabetologia</i> , 2019, 62, 78-86.	6.3	71
99	Relationship between high-sensitivity C-reactive protein and subclinical carotid atherosclerosis stratified by glucose metabolic status in Chinese adults. <i>Clinical Cardiology</i> , 2019, 42, 39-46.	1.8	5
100	Raw orange intake is associated with higher prevalence of non-alcoholic fatty liver disease in an adult population. <i>Nutrition</i> , 2019, 60, 252-260.	2.4	10
101	Twelve-Month Incidence of Depressive Symptoms in Suburb-Dwelling Chinese Older Adults: Role of Sarcopenia. <i>Journal of the American Medical Directors Association</i> , 2019, 20, 64-69.	2.5	28
102	Association between dietary raw garlic intake and newly diagnosed nonalcoholic fatty liver disease: a population-based study. <i>European Journal of Endocrinology</i> , 2019, 181, 591-602.	3.7	26
103	Sex-Specific Associations Between Complement Component 3 and Component 4 Levels and Metabolic Syndrome in an Adult Population. <i>Metabolic Syndrome and Related Disorders</i> , 2018, 16, 143-149.	1.3	3
104	Low serum creatinine predicts risk for type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2018, 34, e3011.	4.0	18
105	The association between thyroid function and incidence of metabolic syndrome in euthyroid subjects: Tianjin chronic low-grade systemic inflammation and health cohort study. <i>Clinical Endocrinology</i> , 2018, 88, 735-743.	2.4	15
106	Relationship of serum immunoglobulin levels to blood pressure and hypertension in an adult population. <i>Journal of Human Hypertension</i> , 2018, 32, 212-218.	2.2	15
107	White blood cells count as an indicator to identify whether obesity leads to increased risk of type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2018, 141, 140-147.	2.8	30
108	High Tomato and Tomato Product Consumption is Protective Against the Decline in Handgrip Strength Among Japanese Adults: The Oroshiso Study. <i>Journal of Epidemiology</i> , 2018, 28, 397-403.	2.4	5

#	ARTICLE	IF	CITATIONS
109	Muscle strength rather than muscle mass is associated with osteoporosis in older Chinese adults. <i>Journal of the Formosan Medical Association</i> , 2018, 117, 101-108.	1.7	31
110	A J-shaped association between soy food intake and depressive symptoms in Chinese adults. <i>Clinical Nutrition</i> , 2018, 37, 1013-1018.	5.0	18
111	Response to "Risk factors for developing prediabetes". <i>Diabetes Research and Clinical Practice</i> , 2018, 135, 233-234.	2.8	0
112	Association of microalbuminuria with diabetes is stronger in people with prehypertension compared to those with ideal blood pressure. <i>Nephrology</i> , 2018, 23, 690-696.	1.6	4
113	Soft drinks consumption is associated with nonalcoholic fatty liver disease independent of metabolic syndrome in Chinese population. <i>European Journal of Nutrition</i> , 2018, 57, 2113-2121.	3.9	22
114	Habitual yoghurt consumption and depressive symptoms in a general population study of 19,596 adults. <i>European Journal of Nutrition</i> , 2018, 57, 2621-2628.	3.9	15
115	Prospective associations of social isolation and loneliness with poor sleep quality in older adults. <i>Quality of Life Research</i> , 2018, 27, 683-691.	3.1	89
116	Relationship between thyroid function and elevated blood pressure in euthyroid adults. <i>Journal of Clinical Hypertension</i> , 2018, 20, 1541-1549.	2.0	21
117	A dietary pattern rich in animal organ, seafood and processed meat products is associated with newly diagnosed hyperuricaemia in Chinese adults: a propensity score-matched case-control study. <i>British Journal of Nutrition</i> , 2018, 119, 1177-1184.	2.3	37
118	The association between missing teeth and nonalcoholic fatty liver disease in adults. <i>Journal of Clinical Periodontology</i> , 2018, 45, 941-951.	4.9	14
119	Aspiration pneumonia induces muscle atrophy in the respiratory, skeletal, and swallowing systems. <i>Journal of Cachexia, Sarcopenia and Muscle</i> , 2018, 9, 643-653.	7.3	77
120	Association between bedtime and the prevalence of newly diagnosed nonalcoholic fatty liver disease in adults. <i>Liver International</i> , 2018, 38, 2277-2286.	3.9	31
121	Comparing the inflammatory profiles for incidence of diabetes mellitus and cardiovascular diseases: a prospective study exploring the "common soil" hypothesis. <i>Cardiovascular Diabetology</i> , 2018, 17, 87.	6.8	40
122	Serum levels of immunoglobulins in an adult population and their relationship with nonalcoholic fatty liver disease. <i>Journal of Digestive Diseases</i> , 2018, 19, 498-507.	1.5	5
123	The relationship between thyroid function and the prevalence of type 2 diabetes mellitus in euthyroid subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, jc.2016-2965.	3.6	74
124	Both muscle mass and muscle strength are inversely associated with depressive symptoms in an elderly Chinese population. <i>International Journal of Geriatric Psychiatry</i> , 2017, 32, 769-778.	2.7	27
125	Independent and combined relationship of habitual unhealthy eating behaviors with depressive symptoms: A prospective study. <i>Journal of Epidemiology</i> , 2017, 27, 42-47.	2.4	22
126	Consumption of low-fat dairy, but not whole-fat dairy, is inversely associated with depressive symptoms in Japanese adults. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2017, 52, 847-853.	3.1	37

#	ARTICLE	IF	CITATIONS
127	Cracked Teeth and Poor Oral Masticatory Habits: A Matched Case-control Study in China. <i>Journal of Endodontics</i> , 2017, 43, 885-889.	3.1	11
128	Combined Effect of Obesity and Mobility Limitation with Incidence of Type 2 Diabetes and Mortality in Chinese Elderly. <i>Rejuvenation Research</i> , 2017, 20, 375-382.	1.8	6
129	The association between immunoglobulin concentrations and prediabetes prevalence in a large Chinese cohort. <i>Metabolism: Clinical and Experimental</i> , 2017, 73, 77-84.	3.4	4
130	Elevated serum complement C3 levels are associated with prehypertension in an adult population. <i>Clinical and Experimental Hypertension</i> , 2017, 39, 42-49.	1.3	15
131	Comparing the diagnostic ability of inflammatory markers in metabolic syndrome. <i>Clinica Chimica Acta</i> , 2017, 475, 1-6.	1.1	17
132	Effects of protease-treated royal jelly on muscle strength in elderly nursing home residents: A randomized, double-blind, placebo-controlled, dose-response study. <i>Scientific Reports</i> , 2017, 7, 11416.	3.3	17
133	The increased risk of sarcopenia in patients with cardiovascular risk factors in Suburb-Dwelling older Chinese using the AWGS definition. <i>Scientific Reports</i> , 2017, 7, 9592.	3.3	55
134	Increased serum ferritin levels are independently associated with carotid atherosclerosis in women. <i>British Journal of Nutrition</i> , 2017, 117, 1623-1630.	2.3	8
135	Red cell distribution width is associated with hemoglobin A1C elevation, but not glucose elevation. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1544-1548.	2.3	12
136	Consumption of preserved egg, a high-lead-containing food, is strongly associated with depressive symptoms in Chinese adults. <i>British Journal of Nutrition</i> , 2017, 118, 737-742.	2.3	14
137	Dietary patterns are associated with depressive symptoms among Chinese adults: a case-control study with propensity score matching. <i>European Journal of Nutrition</i> , 2017, 56, 2577-2587.	3.9	52
138	Timed Up and Go Test can predict recurrent falls: a longitudinal study of the community-dwelling elderly in China. <i>Clinical Interventions in Aging</i> , 2017, Volume 12, 2009-2016.	2.9	58
139	Nutritional Biomarkers, Gene-Diet Interaction, and Risk Factors for Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2016, 2016, 1-2.	2.3	14
140	NUT CONSUMPTION IS ASSOCIATED WITH DEPRESSIVE SYMPTOMS AMONG CHINESE ADULTS. <i>Depression and Anxiety</i> , 2016, 33, 1065-1072.	4.1	39
141	Association between dietary patterns and metabolic syndrome in Chinese adults: a propensity score-matched case-control study. <i>Scientific Reports</i> , 2016, 6, 34748.	3.3	13
142	Mobility and Muscle Strength Together are More Strongly Correlated with Falls in Suburb-Dwelling Older Chinese. <i>Scientific Reports</i> , 2016, 6, 25420.	3.3	33
143	Dietary Patterns are Associated with Helicobacter Pylori Infection in Chinese Adults: A Cross-Sectional Study. <i>Scientific Reports</i> , 2016, 6, 32334.	3.3	22
144	Relationship between grip strength and newly diagnosed nonalcoholic fatty liver disease in a large-scale adult population. <i>Scientific Reports</i> , 2016, 6, 33255.	3.3	38

#	ARTICLE	IF	CITATIONS
145	Knee loading inhibits osteoclast lineage in a mouse model of osteoarthritis. <i>Scientific Reports</i> , 2016, 6, 24668.	3.3	41
146	Consumption of chilies, but not sweet peppers, is positively related to handgrip strength in an adult population. <i>Journal of Nutrition, Health and Aging</i> , 2016, 20, 546-552.	3.3	13
147	High serum adiponectin levels predict incident falls among middle-aged and older adults: a prospective cohort study. <i>Age and Ageing</i> , 2016, 45, 366-371.	1.6	12
148	Serum levels of immunoglobulins in an adult population and their relationship with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2016, 115, 76-82.	2.8	28
149	Association of self-reported sleep duration and hypertension: Results of a Chinese prospective cohort study. <i>Clinical and Experimental Hypertension</i> , 2016, 38, 514-519.	1.3	15
150	The overall computer/mobile devices usage time is related to newly diagnosed non-alcoholic fatty liver disease: a population-based study. <i>Annals of Medicine</i> , 2016, 48, 568-576.	3.8	10
151	The predictive value of mean serum uric acid levels for developing prediabetes. <i>Diabetes Research and Clinical Practice</i> , 2016, 118, 79-89.	2.8	11
152	The traditional Japanese dietary pattern and longitudinal changes in cardiovascular disease risk factors in apparently healthy Japanese adults. <i>European Journal of Nutrition</i> , 2016, 55, 267-279.	3.9	26
153	Prevalence and Factors Associated With Sarcopenia in Suburb-dwelling Older Chinese Using the Asian Working Group for Sarcopenia Definition. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2016, 71, 529-535.	3.6	102
154	Clinical Relevance of Different Handgrip Strength Indexes and Mobility Limitation in the Elderly Adults. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2016, 71, 96-102.	3.6	27
155	Association between Serum Ferritin Concentrations and Depressive Symptoms among Chinese Adults: A Population Study from the Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIHealth) Cohort Study. <i>PLoS ONE</i> , 2016, 11, e0162682.	2.5	9
156	Elevated Serum Level of Interleukin 17 in a Population With Prehypertension. <i>Journal of Clinical Hypertension</i> , 2015, 17, 770-774.	2.0	42
157	Relationship between daily isoflavone intake and sleep in Japanese adults: a cross-sectional study. <i>Nutrition Journal</i> , 2015, 14, 127.	3.4	36
158	Body Mass Index Is Associated with Physical Performance in Suburb-Dwelling Older Chinese: A Cross-Sectional Study. <i>PLoS ONE</i> , 2015, 10, e0119914.	2.5	14
159	Meta-Analysis of Dexmedetomidine on Emergence Agitation and Recovery Profiles in Children after Sevoflurane Anesthesia: Different Administration and Different Dosage. <i>PLoS ONE</i> , 2015, 10, e0123728.	2.5	41
160	Neutrophil:lymphocyte ratio is positively related to type 2 diabetes in a large-scale adult population: a Tianjin Chronic Low-Grade Systemic Inflammation and Health cohort study. <i>European Journal of Endocrinology</i> , 2015, 173, 217-225.	3.7	52
161	Soft drink consumption is associated with depressive symptoms among adults in China. <i>Journal of Affective Disorders</i> , 2015, 172, 422-427.	4.1	50
162	Does elevated serum uric acid level predict the hypertension incidence? A Chinese prospective cohort study. <i>Clinical and Experimental Hypertension</i> , 2015, 37, 498-504.	1.3	12

#	ARTICLE	IF	CITATIONS
163	Blood Neutrophil to Lymphocyte Ratio as a Predictor of Hypertension. <i>American Journal of Hypertension</i> , 2015, 28, 1339-1346.	2.0	106
164	Response to “The Relation Between Neutrophil-Lymphocyte Ratio and Hypertension”. <i>American Journal of Hypertension</i> , 2015, 28, 1387-1387.	2.0	4
165	Effect of beta-hydroxy-beta-methylbutyrate supplementation on muscle loss in older adults: A systematic review and meta-analysis. <i>Archives of Gerontology and Geriatrics</i> , 2015, 61, 168-175.	3.0	101
166	Relationship of anabolic and catabolic biomarkers with muscle strength and physical performance in older adults: a population-based cross-sectional study. <i>BMC Musculoskeletal Disorders</i> , 2015, 16, 202.	1.9	23
167	Chronic inflammation, lymphangiogenesis, and effect of an anti-VEGFR therapy in a mouse model and in human patients with aspiration pneumonia. <i>Journal of Pathology</i> , 2015, 235, 632-645.	4.5	41
168	Staple Foods Consumption and Irritable Bowel Syndrome in Japanese Adults: A Cross-Sectional Study. <i>PLoS ONE</i> , 2015, 10, e0119097.	2.5	9
169	Association between Complement C3 and Prevalence of Fatty Liver Disease in an Adult Population: A Cross-Sectional Study from the Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIHealth) Cohort Study. <i>PLoS ONE</i> , 2015, 10, e0122026.	2.5	32
170	Serum Immunoglobulin M Concentration Varies with Triglyceride Levels in an Adult Population: Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIHealth) Cohort Study. <i>PLoS ONE</i> , 2015, 10, e0124255.	2.5	3
171	Leg Extension Power Is a Pre-Disaster Modifiable Risk Factor for Post-Traumatic Stress Disorder among Survivors of the Great East Japan Earthquake: A Retrospective Cohort Study. <i>PLoS ONE</i> , 2014, 9, e96131.	2.5	12
172	Frequent coffee drinking is associated with higher leg extension power among adult men : a cross-sectional study. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2014, 63, 383-389.	0.0	0
173	Cortisol is not the primary mediator for augmented CXCR4 expression on natural killer cells after acute exercise. <i>Journal of Applied Physiology</i> , 2014, 117, 199-204.	2.5	6
174	Subnormal Peripheral Blood Leukocyte Counts Are Related to the Lowest Prevalence and Incidence of Metabolic Syndrome: Tianjin Chronic Low-Grade Systemic Inflammation and Health Cohort Study. <i>Mediators of Inflammation</i> , 2014, 2014, 1-12.	3.0	36
175	Lower Serum Endogenous Secretory Receptor for Advanced Glycation End Product Level as a Risk Factor of Metabolic Syndrome Among Japanese Adult Men: A 2-Year Longitudinal Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014, 99, 587-593.	3.6	18
176	Long-term Tai Chi training is related to depressive symptoms among Tai Chi practitioners. <i>Journal of Affective Disorders</i> , 2014, 169, 36-39.	4.1	11
177	Higher serum soluble receptor for advanced glycation end product levels and lower prevalence of metabolic syndrome among Japanese adult men: a cross-sectional study. <i>Diabetology and Metabolic Syndrome</i> , 2014, 6, 33.	2.7	16
178	Coffee treatment prevents the progression of sarcopenia in aged mice in vivo and in vitro. <i>Experimental Gerontology</i> , 2014, 50, 1-8.	2.8	37
179	Serum Immunoglobulin M Concentration Is Positively Related to Metabolic Syndrome in an Adult Population: Tianjin Chronic Low-Grade Systemic Inflammation and Health (TCLSIH) Cohort Study. <i>PLoS ONE</i> , 2014, 9, e88701.	2.5	29
180	Irritable Bowel Syndrome Is Positively Related to Metabolic Syndrome: A Population-Based Cross-Sectional Study. <i>PLoS ONE</i> , 2014, 9, e112289.	2.5	37

#	ARTICLE	IF	CITATIONS
181	Poor Lower Extremity Function Was Associated with Pre-Diabetes and Diabetes in Older Chinese People. PLoS ONE, 2014, 9, e115883.	2.5	38
182	Low-fat dairy, but not whole-/high-fat dairy, consumption is related with higher serum adiponectin levels in apparently healthy adults. European Journal of Nutrition, 2013, 52, 771-778.	3.9	9
183	A tomato-rich diet is related to depressive symptoms among an elderly population aged 70 years and over: A population-based, cross-sectional analysis. Journal of Affective Disorders, 2013, 144, 165-170.	4.1	40
184	An inverted J-shaped association of serum uric acid with muscle strength among Japanese adult men: a cross-sectional study. BMC Musculoskeletal Disorders, 2013, 14, 258.	1.9	52
185	Longitudinal changes in the relationship between serum adiponectin concentration and cardiovascular risk factors among apparently healthy middle-aged adults. International Journal of Cardiology, 2013, 167, 2318-2320.	1.7	4
186	The impact of the 2011 Great East Japan Earthquake on hospitalisation for respiratory disease in a rapidly aging society: a retrospective descriptive and cross-sectional study at the disaster base hospital in Ishinomaki. BMJ Open, 2013, 3, e000865.	1.9	60
187	Royal Jelly Prevents the Progression of Sarcopenia in Aged Mice In Vivo and In Vitro. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2013, 68, 1482-1492.	3.6	34
188	Green Tea and Feeling Depressed. , 2013, , 1385-1398.		0
189	Impact of Blunted Perception of Dyspnea on Medical Care Use and Expenditure, and Mortality in Elderly People. Frontiers in Physiology, 2012, 3, 238.	2.8	13
190	Frequent alcohol drinking is associated with lower prevalence of self-reported common cold: a retrospective study. BMC Public Health, 2012, 12, 987.	2.9	13
191	A Distinct Regulatory Role of Th17 Cytokines IL-17A and IL-17F in Chemokine Secretion from Lung Microvascular Endothelial Cells. Inflammation, 2012, 35, 1119-1131.	3.8	44
192	The predictive power of physical function assessed by questionnaire and physical performance measures for subsequent disability. Aging Clinical and Experimental Research, 2012, 24, 345-53.	2.9	14
193	Serum C-Reactive Protein Even at Very Low ($\leq 1.0\text{ mg/l}$) Concentration Is Associated with Physical Performance in a Community-Based Elderly Population Aged 70 Years and Over. Gerontology, 2008, 54, 260-267.	2.8	15
194	Dietary long-chain ω -3 fatty acids of marine origin and serum C-reactive protein concentrations are associated in a population with a diet rich in marine products. American Journal of Clinical Nutrition, 2006, 84, 223-229.	4.7	84
195	Relative Handgrip Strength But Not Absolute Handgrip Strength is a Determinant of Incident Type 2 Diabetes: A Prospective Cohort Study and Systematic Review with Meta-Analysis. SSRN Electronic Journal, 0, , .	0.4	0
196	Non-Linear Associations of Total and Differential Leukocyte Counts with Incident Nonalcoholic Fatty Liver Disease. SSRN Electronic Journal, 0, , .	0.4	0
197	Serum Immunoglobulin Concentrations and Risk of Type 2 Diabetes Mellitus in Adults: A Prospective Cohort Study. SSRN Electronic Journal, 0, , .	0.4	0
198	Serum Immunoglobulin Concentrations and Risk of Type 2 Diabetes Mellitus in Adults: A Prospective Cohort Study. SSRN Electronic Journal, 0, , .	0.4	0

#	ARTICLE	IF	CITATIONS
199	Association between Consumption of Edible Seaweeds and Newly Diagnosed Non-Alcohol Fatty Liver Disease: The TCLSIH Cohort Study. SSRN Electronic Journal, 0, , .	0.4	0