

Martin L Kieffer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2867762/publications.pdf>

Version: 2024-02-01

21
papers

2,800
citations

516710

16
h-index

677142

22
g-index

24
all docs

24
docs citations

24
times ranked

3894
citing authors

#	ARTICLE	IF	CITATIONS
1	New fluorescent auxin probes visualise tissue-specific and subcellular distributions of auxin in Arabidopsis. <i>New Phytologist</i> , 2021, 230, 535-549.	7.3	15
2	Selective auxin agonists induce specific AUX/IAA protein degradation to modulate plant development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019, 116, 6463-6472.	7.1	23
3	The Tetrazole Analogue of the Auxin Indole-3-acetic Acid Binds Preferentially to TIR1 and Not AFB5. <i>ACS Chemical Biology</i> , 2018, 13, 2585-2594.	3.4	13
4	The developmental and environmental regulation of gravitropic setpoint angle in Arabidopsis and bean. <i>Scientific Reports</i> , 2017, 7, 42664.	3.3	44
5	The Arabidopsis JAGGED LATERAL ORGANS (JLO) gene sensitizes plants to auxin. <i>Journal of Experimental Botany</i> , 2017, 68, 2741-2755.	4.8	11
6	HSP90 regulates temperature-dependent seedling growth in Arabidopsis by stabilizing the auxin co-receptor F-box protein TIR1. <i>Nature Communications</i> , 2016, 7, 10269.	12.8	210
7	Auxin Controls Gravitropic Setpoint Angle in Higher Plant Lateral Branches. <i>Current Biology</i> , 2013, 23, 1497-1504.	3.9	116
8	The Arabidopsis O-Linked N-Acetylglucosamine Transferase SPINDLY Interacts with Class I TCPs to Facilitate Cytokinin Responses in Leaves and Flowers. <i>Plant Cell</i> , 2012, 24, 96-108.	6.6	142
9	In Vitro Propagation of Cauliflower Using Curd Microexplants. <i>Methods in Molecular Biology</i> , 2012, 11013, 329-339.	0.9	1
10	TCP14 and TCP15 affect internode length and leaf shape in Arabidopsis. <i>Plant Journal</i> , 2011, 68, 147-158.	5.7	261
11	Defining auxin response contexts in plant development. <i>Current Opinion in Plant Biology</i> , 2010, 13, 12-20.	7.1	125
12	Analysis of the Transcription Factor WUSCHEL and Its Functional Homologue in Antirrhinum Reveals a Potential Mechanism for Their Roles in Meristem Maintenance. <i>Plant Cell</i> , 2006, 18, 560-573.	6.6	203
13	UPF1 is required for nonsense-mediated mRNA decay (NMD) and RNAi in Arabidopsis. <i>Plant Journal</i> , 2006, 47, 480-489.	5.7	183
14	Comprehensive Interaction Map of the Arabidopsis MADS Box Transcription Factors. <i>Plant Cell</i> , 2005, 17, 1424-1433.	6.6	528
15	Molecular and Phylogenetic Analyses of the Complete MADS-Box Transcription Factor Family in Arabidopsis. <i>Plant Cell</i> , 2003, 15, 1538-1551.	6.6	758
16	PLANT BIOLOGY: MADS-Box Genes Reach Maturity. <i>Science</i> , 2002, 296, 275-276.	12.6	62
17	Developmental programmes in floral organ formation. <i>Seminars in Cell and Developmental Biology</i> , 2001, 12, 373-380.	5.0	22
18	A cost effective protocol for in vitro mass propagation of cauliflower. <i>Plant Science</i> , 2001, 160, 1015-1024.	3.6	18

#	ARTICLE	IF	CITATIONS
19	Explaining curd and spear geometry in broccoli, cauliflower and 'romanesco': quantitative variation in activity of primary meristems. <i>Planta</i> , 1998, 206, 34-43.	3.2	25
20	Rapid mass production of cauliflower propagules from fractionated and graded curd. <i>Plant Science</i> , 1995, 107, 229-235.	3.6	18
21	Anther culture of kale (<i>Brassica oleracea</i> L. convar.acephala (DC.) Alef.). <i>Plant Cell, Tissue and Organ Culture</i> , 1993, 33, 303-313.	2.3	9