

Chun-fa Chen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2935335/publications.pdf>

Version: 2024-02-01

25
papers

845
citations

567281

15
h-index

552781

26
g-index

27
all docs

27
docs citations

27
times ranked

1311
citing authors

#	ARTICLE	IF	CITATIONS
1	Lymph Node Ratio Is an Independent Prognostic Factor for Patients with Siewert Type II Adenocarcinoma of Esophagogastric Junction: Results from a 10-Year Follow-up Study. <i>Journal of Gastrointestinal Cancer</i> , 2021, 52, 983-992.	1.3	8
2	Notch3 inhibits cell proliferation and tumorigenesis and predicts better prognosis in breast cancer through transactivating PTEN. <i>Cell Death and Disease</i> , 2021, 12, 502.	6.3	19
3	CXCL9 Is a Potential Biomarker of Immune Infiltration Associated With Favorable Prognosis in ER-Negative Breast Cancer. <i>Frontiers in Oncology</i> , 2021, 11, 710286.	2.8	30
4	O-6-methylguanine DNA methyltransferase is a favorable biomarker with proliferation suppressive potential in Breast Cancer. <i>Journal of Cancer</i> , 2020, 11, 6326-6336.	2.5	4
5	A negative binomial regression model for risk estimation of 2 axillary lymph node metastases in breast cancer patients. <i>Scientific Reports</i> , 2020, 10, 21856.	3.3	2
6	Knockdown of nucleophosmin 1 suppresses proliferation of triple-negative breast cancer cells through activating CDH1/Skp2/p27kip1 pathway. <i>Cancer Management and Research</i> , 2019, Volume 11, 143-156.	1.9	19
7	TOX3 is a favorable prognostic indicator and potential immunomodulatory factor in lung adenocarcinoma. <i>Oncology Letters</i> , 2019, 18, 4144-4152.	1.8	8
8	Performing Data Mining And Integrative Analysis Of Biomarker in Breast Cancer Using Multiple Publicly Accessible Databases. <i>Journal of Visualized Experiments</i> , 2019, , .	0.3	1
9	<p>Long noncoding RNA H19 is a critical oncogenic driver and contributes to epithelial-mesenchymal transition in papillary thyroid carcinoma</p>. <i>Cancer Management and Research</i> , 2019, Volume 11, 2059-2072.	1.9	26
10	circTADA2As suppress breast cancer progression and metastasis via targeting miR-203a-3p/SOCS3 axis. <i>Cell Death and Disease</i> , 2019, 10, 175.	6.3	187
11	GATA Binding Protein 3 Boosts Extracellular ATP Hydrolysis and Inhibits Metastasis of Breast Cancer by Up-regulating Ectonucleoside Triphosphate Diphosphohydrolase 3. <i>International Journal of Biological Sciences</i> , 2019, 15, 2522-2537.	6.4	6
12	Decrease in the Ki67 index during neoadjuvant chemotherapy predicts favorable relapse-free survival in patients with locally advanced breast cancer. <i>Cancer Biology and Medicine</i> , 2019, 16, 575-586.	3.0	14
13	Prognostic values of the inhibitor of DNA-binding family members in breast cancer. <i>Oncology Reports</i> , 2018, 40, 1897-1906.	2.6	9
14	MiR-221/222 promote epithelial-mesenchymal transition by targeting Notch3 in breast cancer cell lines. <i>Npj Breast Cancer</i> , 2018, 4, 20.	5.2	52
15	Notch3 inhibits epithelial-mesenchymal transition in breast cancer via a novel mechanism, upregulation of GATA-3 expression. <i>Oncogenesis</i> , 2018, 7, 59.	4.9	31
16	Predictive Value of Preoperative Multidetector-Row Computed Tomography for Axillary Lymph Nodes Metastasis in Patients With Breast Cancer. <i>Frontiers in Oncology</i> , 2018, 8, 666.	2.8	17
17	FoxM1 is a promising candidate target in the treatment of breast cancer. <i>Oncotarget</i> , 2018, 9, 842-852.	1.8	31
18	CCR10 activation stimulates the invasion and migration of breast cancer cells through the ERK1/2/MMP-7 signaling pathway. <i>International Immunopharmacology</i> , 2017, 51, 124-130.	3.8	29

#	ARTICLE	IF	CITATIONS
19	MCAM/CD146 promotes tamoxifen resistance in breast cancer cells through induction of epithelialâ€“mesenchymal transition, decreased ER α expression and AKT activation. <i>Cancer Letters</i> , 2017, 386, 65-76.	7.2	54
20	GATA3 and TRPS1 are distinct biomarkers and prognostic factors in breast cancer: database mining for GATA family members in malignancies. <i>Oncotarget</i> , 2017, 8, 34750-34761.	1.8	39
21	Notch3 Maintains Luminal Phenotype and Suppresses Tumorigenesis and Metastasis of Breast Cancer via Trans-Activating Estrogen Receptor- α . <i>Theranostics</i> , 2017, 7, 4041-4056.	10.0	48
22	Prognostic values of distinct CBX family members in breast cancer. <i>Oncotarget</i> , 2017, 8, 92375-92387.	1.8	67
23	Notch3 overexpression causes arrest of cell cycle progression by inducing Cdh1 expression in human breast cancer cells. <i>Cell Cycle</i> , 2016, 15, 432-440.	2.6	36
24	A case of primary neuroendocrine breast carcinoma that responded to neo-adjuvant chemotherapy. <i>Frontiers of Medicine</i> , 2015, 9, 112-116.	3.4	15
25	Cytoplasmic Skp2 Expression Is Associated with p-Akt1 and Predicts Poor Prognosis in Human Breast Carcinomas. <i>PLoS ONE</i> , 2012, 7, e52675.	2.5	49