

Peter Ladlow

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2967230/publications.pdf>

Version: 2024-02-01

13
papers

313
citations

933447

10
h-index

1199594

12
g-index

13
all docs

13
docs citations

13
times ranked

248
citing authors

#	ARTICLE	IF	CITATIONS
1	Integration of strength training into UK Defence Rehabilitation practice: current trends and future challenges. <i>BMJ Military Health</i> , 2022, 168, 314-319.	0.9	12
2	Effect of a physical activity and behaviour maintenance programme on functional mobility decline in older adults: the REACT (Retirement in Action) randomised controlled trial. <i>Lancet Public Health</i> , The, 2022, 7, e316-e326.	10.0	26
3	Dysautonomia following COVID-19 is not associated with subjective limitations or symptoms but is associated with objective functional limitations. <i>Heart Rhythm</i> , 2022, 19, 613-620.	0.7	60
4	Cognitive functional therapy (CFT)-based rehabilitation improves clinical outcomes in UK military personnel with persistent low back pain. <i>BMJ Military Health</i> , 2020, 166, 336-341.	0.9	10
5	The Tribulations of Trials: Lessons Learnt Recruiting 777 Older Adults Into REtirement in ACTion (REACT), a Trial of a Community, Group-Based Active Aging Intervention Targeting Mobility Disability. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2020, 75, 2387-2395.	3.6	13
6	Predicting ambulatory energy expenditure in lower limb amputees using multi-sensor methods. <i>PLoS ONE</i> , 2019, 14, e0209249.	2.5	9
7	Passive-dynamic ankle-foot orthosis improves medium-term clinical outcomes after severe lower extremity trauma. <i>Journal of the Royal Army Medical Corps</i> , 2019, 165, 330-337.	0.8	8
8	Low-Load Resistance Training With Blood Flow Restriction Improves Clinical Outcomes in Musculoskeletal Rehabilitation: A Single-Blind Randomized Controlled Trial. <i>Frontiers in Physiology</i> , 2018, 9, 1269.	2.8	76
9	The effects of low-intensity blood flow restricted exercise compared with conventional resistance training on the clinical outcomes of active UK military personnel following a 3-week in-patient rehabilitation programme: protocol for a randomized controlled feasibility study. <i>Pilot and Feasibility Studies</i> , 2017, 3, 71.	1.2	10
10	Impact of anatomical placement of an accelerometer on prediction of physical activity energy expenditure in lower-limb amputees. <i>PLoS ONE</i> , 2017, 12, e0185731.	2.5	14
11	Influence of Immediate and Delayed Lower-Limb Amputation Compared with Lower-Limb Salvage on Functional and Mental Health Outcomes Post-Rehabilitation in the U.K. Military. <i>Journal of Bone and Joint Surgery - Series A</i> , 2016, 98, 1996-2005.	3.0	36
12	Validity Of The Actiheart Monitor For Predicting Physical Activity Energy Expenditure In Lower-limb Amputees.. <i>Medicine and Science in Sports and Exercise</i> , 2016, 48, 982-983.	0.4	0
13	Functional and Mental Health Status of United Kingdom Military Amputees Postrehabilitation. <i>Archives of Physical Medicine and Rehabilitation</i> , 2015, 96, 2048-2054.	0.9	39