

# Josã Rioja Villodres

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2967574/publications.pdf>

Version: 2024-02-01

29  
papers

665  
citations

567144

15  
h-index

580701

25  
g-index

30  
all docs

30  
docs citations

30  
times ranked

1192  
citing authors

#	ARTICLE	IF	CITATIONS
1	Postprandial Hyperlipidemia: Association with Inflammation and Subclinical Atherosclerosis in Patients with Rheumatoid Arthritis. <i>Biomedicines</i> , 2022, 10, 133.	1.4	6
2	Genetic variants in the LPL and GPIHBP1 genes, in patients with severe hypertriglyceridaemia, detected with high resolution melting analysis. <i>Clinica Chimica Acta</i> , 2020, 500, 163-171.	0.5	8
3	Postprandial Apolipoprotein B48 is Associated with Subclinical Atherosclerosis in Patients with Rheumatoid Arthritis. <i>Journal of Clinical Medicine</i> , 2020, 9, 2483.	1.0	11
4	Evaluation of the chylomicronâ€ƒTG to VLDLâ€ƒTG ratio for type I hyperlipoproteinemia diagnostic. <i>European Journal of Clinical Investigation</i> , 2020, 50, e13345.	1.7	16
5	Subclinical Arteriosclerosis is Associated with Common Vascular Risk Factors in Long-Term Survivors of Testicular Cancer. <i>Journal of Clinical Medicine</i> , 2020, 9, 971.	1.0	0
6	Fractional excretion of phosphorus and vascular calcification in stage 3 chronic kidney disease. <i>Journal of Investigative Medicine</i> , 2019, 67, 674-680.	0.7	8
7	Evidence for increased chylomicron remnants in rheumatoid arthritis. <i>European Journal of Clinical Investigation</i> , 2018, 48, e12873.	1.7	8
8	High risk of subclinical atherosclerosis in COPD exacerbator phenotype. <i>Respiratory Medicine</i> , 2018, 141, 165-171.	1.3	8
9	The Effects of Parenteral K1 Administration in Pseudoxanthoma Elasticum Patients Versus Controls. A Pilot Study. <i>Frontiers in Medicine</i> , 2018, 5, 86.	1.2	6
10	Molecular basis of the familial chylomicronemia syndrome in patients from the National Dyslipidemia Registry of the Spanish Atherosclerosis Society. <i>Journal of Clinical Lipidology</i> , 2018, 12, 1482-1492.e3.	0.6	22
11	Alzheimerâ€™s like brain changes correlate with low adiponectin plasma levels in type 2 diabetic patients. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 281-286.	1.2	33
12	Insulin resistance and levels of adipokines in patients with untreated early rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2016, 35, 43-53.	1.0	29
13	Novel mutations in the GPIHBP1 gene identified in 2 patients with recurrent acute pancreatitis. <i>Journal of Clinical Lipidology</i> , 2016, 10, 92-100.e1.	0.6	27
14	Inâ€švivo evidence for chylomicrons as mediators of postprandial inflammation. <i>Atherosclerosis</i> , 2015, 243, 540-545.	0.4	18
15	Structural and Functional Brain Changes in Middle-Aged Type 2 Diabetic Patients: A Cross-Sectional Study. <i>Journal of Alzheimer's Disease</i> , 2014, 40, 375-386.	1.2	58
16	Cognitive Dysfunctions in Middle-Aged Type 2 Diabetic Patients and Neuroimaging Correlations: A Cross-Sectional Study. <i>Journal of Alzheimer's Disease</i> , 2014, 42, 1337-1346.	1.2	66
17	Exploring the value of apoB48 as a marker for atherosclerosis in clinical practice. <i>European Journal of Clinical Investigation</i> , 2012, 42, 702-708.	1.7	35
18	Análisis de la influencia de polimorfismos en APOE, APOA5, LPL, LIPC y CETP sobre los niveles de triglicéridos en población laboral malagueña. <i>Clínica E Investigaci3n En Arteriosclerosis</i> , 2011, 23, 62-71.	0.4	1

#	ARTICLE	IF	CITATIONS
19	Additive effects of LPL, APOA5 and APOE variant combinations on triglyceride levels and hypertriglyceridemia: results of the ICARIA genetic sub-study. <i>BMC Medical Genetics</i> , 2010, 11, 66.	2.1	55
20	Lipoprotein lipase activity and mass, apolipoprotein C-II mass and polymorphisms of apolipoproteins E and A5 in subjects with prior acute hypertriglyceridaemic pancreatitis. <i>BMC Gastroenterology</i> , 2009, 9, 46.	0.8	26
21	Omega 3 fatty acids induce a marked reduction of apolipoprotein B48 when added to fluvastatin in patients with type 2 diabetes and mixed hyperlipidemia: a preliminary report. <i>Cardiovascular Diabetology</i> , 2009, 8, 1.	2.7	55
22	Plasma redox status is impaired in the portacaval shunted rat – the risk of the reduced antioxidant ability. <i>Comparative Hepatology</i> , 2008, 7, 1.	0.9	7
23	Association of the -250G/A promoter polymorphism of the hepatic lipase gene with the risk of peripheral arterial disease in type 2 diabetic patients. <i>Journal of Diabetes and Its Complications</i> , 2008, 22, 273-277.	1.2	13
24	Polimorfismo –250G/A de LIPC y su asociaci3n con la enfermedad arterial perif3rica en pacientes con diabetes tipo 2. <i>Cl3nica E Investigaci3n En Arteriosclerosis</i> , 2008, 20, 22-27.	0.4	0
25	Respuesta posprandial de la apolipoprote3na B-48 y del perfil lip3dico en pacientes con diabetes mellitus tipo 2. <i>Cl3nica E Investigaci3n En Arteriosclerosis</i> , 2008, 20, 145-150.	0.4	0
26	Low Doses of Insulin-Like Growth Factor-I Induce Mitochondrial Protection in Aging Rats. <i>Endocrinology</i> , 2008, 149, 2620-2627.	1.4	103
27	5-HT1A receptor activation counteracted the effect of acute immobilization of noradrenergic neurons in the rat locus coeruleus. <i>Neuroscience Letters</i> , 2007, 412, 84-88.	1.0	8
28	Smoking and postprandial triglycerides are associated with vascular disease in patients with type 2 diabetes. <i>Atherosclerosis</i> , 2007, 194, 391-396.	0.4	19
29	5-HT1A receptor activation counteracts c-Fos immunoreactivity induced in serotonin neurons of the raphe nuclei after immobilization stress in the male rat. <i>Neuroscience Letters</i> , 2006, 397, 190-195.	1.0	17