

Tatsunori Maekawa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2976724/publications.pdf>

Version: 2024-02-01

16
papers

426
citations

840585

11
h-index

1058333

14
g-index

16
all docs

16
docs citations

16
times ranked

679
citing authors

#	ARTICLE	IF	CITATIONS
1	Histamine H2-Receptor Antagonists Improve Non-Steroidal Anti-Inflammatory Drug-Induced Intestinal Dysbiosis. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8166.	1.8	11
2	Leucine-Rich Repeat Kinase 2 Controls Inflammatory Cytokines Production through NF- κ B Phosphorylation and Antigen Presentation in Bone Marrow-Derived Dendritic Cells. <i>International Journal of Molecular Sciences</i> , 2020, 21, 1890.	1.8	7
3	LRRK2 Inhibition Ameliorates Dexamethasone-Induced Glucose Intolerance <i>via</i> Prevents Impairment in GLUT4 Membrane Translocation in Adipocytes. <i>Biological and Pharmaceutical Bulletin</i> , 2020, 43, 1660-1668.	0.6	6
4	Leucine-Rich Repeat Kinase 2 Is Associated With Activation of the Paraventricular Nucleus of the Hypothalamus and Stress-Related Gastrointestinal Dysmotility. <i>Frontiers in Neuroscience</i> , 2019, 13, 905.	1.4	10
5	LRRK2: An Emerging New Molecule in the Enteric Neuronal System That Quantitatively Regulates Neuronal Peptides and IgA in the Gut. <i>Digestive Diseases and Sciences</i> , 2017, 62, 903-912.	1.1	17
6	Leucine-rich repeat kinase 2 (LRRK2) regulates α -synuclein clearance in microglia. <i>BMC Neuroscience</i> , 2016, 17, 77.	0.8	48
7	Leucine-rich repeat kinase 2 is a regulator of B cell function, affecting homeostasis, BCR signaling, IgA production, and TI antigen responses. <i>Journal of Neuroimmunology</i> , 2016, 292, 1-8.	1.1	16
8	Influence of H2-receptor antagonists on intestinal mucositis induced by 5-fluorouracil in rats. <i>Cancer Research Frontiers</i> , 2016, 2, 33-42.	0.2	0
9	Elemental diet moderates 5-fluorouracil-induced gastrointestinal mucositis through mucus barrier alteration. <i>Cancer Chemotherapy and Pharmacology</i> , 2015, 76, 269-277.	1.1	13
10	Leucine-rich repeat kinase 2 regulates tau phosphorylation through direct activation of glycogen synthase kinase-3 β . <i>FEBS Journal</i> , 2014, 281, 3-13.	2.2	53
11	Tu1242 Changes in the Mucus Barrier Are the Causes or Results of Chemotherapy-Induced Mucositis?. <i>Gastroenterology</i> , 2014, 146, S-793.	0.6	0
12	The I2020T Leucine-rich repeat kinase 2 transgenic mouse exhibits impaired locomotive ability accompanied by dopaminergic neuron abnormalities. <i>Molecular Neurodegeneration</i> , 2012, 7, 15.	4.4	36
13	LRRK2 Phosphorylates Tubulin-Associated Tau but Not the Free Molecule: LRRK2-Mediated Regulation of the Tau-Tubulin Association and Neurite Outgrowth. <i>PLoS ONE</i> , 2012, 7, e30834.	1.1	104
14	LRRK2 is expressed in B-2 but not in B-1 B cells, and downregulated by cellular activation. <i>Journal of Neuroimmunology</i> , 2010, 229, 123-128.	1.1	42
15	Age-dependent and cell-population-restricted LRRK2 expression in normal mouse spleen. <i>Biochemical and Biophysical Research Communications</i> , 2010, 392, 431-435.	1.0	40
16	I2020T leucine-rich repeat kinase 2, the causative mutant molecule of familial Parkinson's disease, has a higher intracellular degradation rate than the wild-type molecule. <i>Biochemical and Biophysical Research Communications</i> , 2009, 390, 710-715.	1.0	23