

Honghao Wang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2977021/publications.pdf>

Version: 2024-02-01

62
papers

5,209
citations

304743

22
h-index

133252

59
g-index

68
all docs

68
docs citations

68
times ranked

5828
citing authors

#	ARTICLE	IF	CITATIONS
1	HMG-1 as a Late Mediator of Endotoxin Lethality in Mice. <i>Science</i> , 1999, 285, 248-251.	12.6	3,807
2	Interleukin-17-secreting T cells in neuromyelitis optica and multiple sclerosis during relapse. <i>Journal of Clinical Neuroscience</i> , 2011, 18, 1313-1317.	1.5	141
3	HLA-DPB1*0501 is associated with susceptibility to anti-aquaporin-4 antibodies positive neuromyelitis optica in Southern Han Chinese. <i>Journal of Neuroimmunology</i> , 2011, 233, 181-184.	2.3	105
4	Increased memory Th17 cells in patients with neuromyelitis optica and multiple sclerosis. <i>Journal of Neuroimmunology</i> , 2011, 234, 155-160.	2.3	90
5	Cerebrospinal Fluid BAFF and APRIL Levels in Neuromyelitis Optica and Multiple Sclerosis Patients During Relapse. <i>Journal of Clinical Immunology</i> , 2012, 32, 1007-1011.	3.8	72
6	Genome-wide association analyses in Han Chinese identify two new susceptibility loci for amyotrophic lateral sclerosis. <i>Nature Genetics</i> , 2013, 45, 697-700.	21.4	67
7	IL-22 secreting CD4 + T cells in the patients with neuromyelitis optica and multiple sclerosis. <i>Journal of Neuroimmunology</i> , 2013, 261, 87-91.	2.3	65
8	Interleukin 17 gene polymorphism is associated with anti-aquaporin 4 antibody-positive neuromyelitis optica in the Southern Han Chinese – A case control study. <i>Journal of the Neurological Sciences</i> , 2012, 314, 26-28.	0.6	41
9	Notable Increased Cerebrospinal Fluid Levels of Soluble Interleukin-6 Receptors in Neuromyelitis Optica. <i>NeuroImmunoModulation</i> , 2012, 19, 304-308.	1.8	39
10	Interleukin-1 receptor associated kinase (IRAK)-M -mediated type 2 microglia polarization ameliorates the severity of experimental autoimmune encephalomyelitis (EAE). <i>Journal of Autoimmunity</i> , 2019, 102, 77-88.	6.5	37
11	Reduced Serum Levels of Triglyceride, Very Low Density Lipoprotein Cholesterol and Apolipoprotein B in Parkinson's Disease Patients. <i>PLoS ONE</i> , 2013, 8, e75743.	2.5	36
12	Elevation of YKL-40 in the CSF of Anti-NMDAR Encephalitis Patients Is Associated With Poor Prognosis. <i>Frontiers in Neurology</i> , 2018, 9, 727.	2.4	33
13	Cerebrospinal fluid levels of CXCL13 are elevated in neuromyelitis optica. <i>Journal of Neuroimmunology</i> , 2011, 240-241, 104-108.	2.3	32
14	Increased plasma levels of pentraxin 3 in patients with multiple sclerosis and neuromyelitis optica. <i>Multiple Sclerosis Journal</i> , 2013, 19, 926-931.	3.0	30
15	Cerebrospinal fluid β -synuclein levels are elevated in multiple sclerosis and neuromyelitis optica patients during relapse. <i>Journal of Neurochemistry</i> , 2012, 122, 19-23.	3.9	29
16	Increased Soluble α -Syn in CSF of Neuromyelitis Optica. <i>Scandinavian Journal of Immunology</i> , 2014, 79, 127-130.	2.7	29
17	Aquaporin 4 Antibodies in the Cerebrospinal Fluid Are Helpful in Diagnosing Chinese Patients with Neuromyelitis Optica. <i>NeuroImmunoModulation</i> , 2012, 19, 96-102.	1.8	28
18	Cerebrospinal Fluid High-Mobility Group Box Protein 1 in Neuromyelitis Optica and Multiple Sclerosis. <i>NeuroImmunoModulation</i> , 2013, 20, 113-118.	1.8	27

#	ARTICLE	IF	CITATIONS
19	Cerebrospinal Fluid IL-21 Levels in Neuromyelitis Optica and Multiple Sclerosis. <i>Canadian Journal of Neurological Sciences</i> , 2012, 39, 813-820.	0.5	26
20	Pregnancy in neuromyelitis optica spectrum disorder: A multicenter study from South China. <i>Journal of the Neurological Sciences</i> , 2017, 372, 152-156.	0.6	26
21	Cerebrospinal fluid light and heavy neurofilament level increased in anti-N-methyl-D-aspartate receptor encephalitis. <i>Brain and Behavior</i> , 2019, 9, e01354.	2.2	26
22	Cell-Free Mitochondrial DNA in the CSF: A Potential Prognostic Biomarker of Anti-NMDAR Encephalitis. <i>Frontiers in Immunology</i> , 2019, 10, 103.	4.8	26
23	Increased Plasma Interleukin-32 Expression in Patients with Neuromyelitis Optica. <i>Journal of Clinical Immunology</i> , 2013, 33, 666-670.	3.8	22
24	The HMGB1 is increased in CSF of patients with an Anti-NMDAR encephalitis. <i>Acta Neurologica Scandinavica</i> , 2018, 137, 277-282.	2.1	20
25	Elevated Soluble Fas and FasL in Cerebrospinal Fluid and Serum of Patients With Anti-N-methyl-D-aspartate Receptor Encephalitis. <i>Frontiers in Neurology</i> , 2018, 9, 904.	2.4	20
26	Elevated neuron-specific enolase and S100 calcium-binding protein B concentrations in cerebrospinal fluid of patients with anti-N-methyl-d-aspartate receptor encephalitis. <i>Clinica Chimica Acta</i> , 2018, 480, 79-83.	1.1	19
27	Higher CSF Levels of NLRP3 Inflammasome Is Associated With Poor Prognosis of Anti-N-methyl-D-Aspartate Receptor Encephalitis. <i>Frontiers in Immunology</i> , 2019, 10, 905.	4.8	19
28	Renal impairment in different phenotypes of Wilson disease. <i>Neurological Sciences</i> , 2015, 36, 2111-2115.	1.9	17
29	Anti-N-methyl-d-aspartate receptor encephalitis associated with intracranial <i>Angiostrongylus cantonensis</i> infection: a case report. <i>Neurological Sciences</i> , 2017, 38, 703-706.	1.9	17
30	Anti-NMDAR encephalitis induced in mice by active immunization with a peptide from the amino-terminal domain of the GluN1 subunit. <i>Journal of Neuroinflammation</i> , 2021, 18, 53.	7.2	17
31	Cerebrospinal fluid light and heavy neurofilaments in neuromyelitis optica. <i>Neurochemistry International</i> , 2013, 63, 805-808.	3.8	15
32	Cerebrospinal fluid pentraxin 3 and CD40 ligand in anti-N-methyl-d-aspartate receptor encephalitis. <i>Journal of Neuroimmunology</i> , 2018, 315, 40-44.	2.3	15
33	Serum lipoprotein levels in patients with neuromyelitis optica elevated but had little correlation with clinical presentations. <i>Clinical Neurology and Neurosurgery</i> , 2010, 112, 478-481.	1.4	14
34	Suppression of lncRNA RMRP ameliorates oxygen-glucose deprivation/re-oxygenation-induced neural cells injury by inhibiting autophagy and PI3K/Akt/mTOR-mediated apoptosis. <i>Bioscience Reports</i> , 2019, 39, .	2.4	14
35	Plasma sCD28, sCTLA-4 levels in neuromyelitis optica and multiple sclerosis during relapse. <i>Journal of Neuroimmunology</i> , 2012, 243, 52-55.	2.3	13
36	Elevated soluble syndecan-1 levels in neuromyelitis optica are associated with disease severity. <i>Cytokine</i> , 2018, 111, 140-145.	3.2	12

#	ARTICLE	IF	CITATIONS
37	Anti-N-methyl-D-aspartate receptor (NMDAR) encephalitis is associated with IRF7 , BANK1 and TBX21 polymorphisms in two populations. <i>European Journal of Neurology</i> , 2021, 28, 595-601.	3.3	12
38	Mitofusin 2 confers the suppression of microglial activation by cannabidiol: Insights from in vitro and in vivo models. <i>Brain, Behavior, and Immunity</i> , 2022, 104, 155-170.	4.1	12
39	Elevated fibrinogen levels in neuromyelitis optica is associated with severity of disease. <i>Neurological Sciences</i> , 2016, 37, 1823-1829.	1.9	11
40	The CSF Levels of Neutrophil-Related Chemokines in Patients with Neuromyelitis Optica. <i>Annals of Clinical and Translational Neurology</i> , 2020, 7, 1245-1251.	3.7	11
41	Elevated Levels of NLRP3 in Cerebrospinal Fluid of Patients With Autoimmune GFAP Astrocytopathy. <i>Frontiers in Neurology</i> , 2019, 10, 1019.	2.4	10
42	NLRP3 level in cerebrospinal fluid of patients with neuromyelitis optica spectrum disorders: Increased levels and association with disease severity. <i>Multiple Sclerosis and Related Disorders</i> , 2020, 39, 101888.	2.0	10
43	A refractory anti-NMDA receptor encephalitis successfully treated by bilateral salpingo-oophorectomy and intrathecal injection of methotrexate and dexamethasone: a case report. <i>Journal of International Medical Research</i> , 2020, 48, 030006052092566.	1.0	10
44	Serum concentration of CD40L is elevated in inflammatory demyelinating diseases. <i>Journal of Neuroimmunology</i> , 2016, 299, 66-69.	2.3	9
45	Status of Immunotherapy Acceptance in Chinese Patients With Multiple Sclerosis: Analysis of Multiple Sclerosis Patient Survival Report 2018. <i>Frontiers in Neurology</i> , 2021, 12, 651511.	2.4	9
46	Comparison of magnetic resonance spectroscopy (MRS) with arterial spin labeling (ASL) in the differentiation between mitochondrial encephalomyopathy, lactic Acidosis, plus stroke-like episodes (MELAS) and acute ischemic stroke (AIS). <i>Journal of Clinical Neuroscience</i> , 2018, 55, 65-70.	1.5	9
47	Cerebrospinal Fluid Level of Soluble CD27 Is Associated with Disease Severity in Neuromyelitis Optica Spectrum Disorder. <i>NeuroImmunoModulation</i> , 2018, 25, 185-192.	1.8	8
48	Elevated Serum and Cerebrospinal Fluid CD138 in Patients With Anti-N-Methyl-d-Aspartate Receptor Encephalitis. <i>Frontiers in Molecular Neuroscience</i> , 2019, 12, 116.	2.9	8
49	Elevated serum brain natriuretic peptide and matrix metalloproteinases 2 and 9 in Wilson's disease. <i>Metabolic Brain Disease</i> , 2015, 30, 1087-1091.	2.9	7
50	Cerebrospinal Fluid Osteopontin and Inflammation-Associated Cytokines in Patients With Anti-N-Methyl-D-Aspartate Receptor Encephalitis. <i>Frontiers in Neurology</i> , 2020, 11, 519692.	2.4	6
51	Predictors of caregiver burden in patients with neurologic Wilson disease. <i>Journal of International Medical Research</i> , 2020, 48, 030006052093015.	1.0	6
52	Therapeutic Response and Possible Biomarkers in Acute Attacks of Neuromyelitis Optica Spectrum Disorders: A Prospective Observational Study. <i>Frontiers in Immunology</i> , 2021, 12, 720907.	4.8	6
53	Clinical significance of soluble adhesion molecules in anti-NMDAR encephalitis patients. <i>Annals of Clinical and Translational Neurology</i> , 2019, 6, 945-953.	3.7	4
54	High Level of Soluble CD146 In Cerebrospinal Fluid Might be a Biomarker of Severity of Anti-N-Methyl-D-Aspartate Receptor Encephalitis. <i>Frontiers in Immunology</i> , 2021, 12, 680424.	4.8	4

#	ARTICLE	IF	CITATIONS
55	High Level of Serum and Cerebrospinal Fluid of Heparan Sulfate and Hyaluronic Acid Might Be a Biomarker of Severity of Neuromyelitis Optica. <i>Frontiers in Immunology</i> , 2021, 12, 705536.	4.8	3
56	Serum pentraxin 3 is elevated in patients with neurological Wilson's disease. <i>Clinica Chimica Acta</i> , 2016, 462, 178-182.	1.1	2
57	Developing normal number of small segmentsâ€activity clouds of the electromyography interference pattern. <i>Muscle and Nerve</i> , 2020, 61, 485-490.	2.2	2
58	Electroconvulsive Therapy and Klinefelter Syndrome. <i>Journal of ECT</i> , 2013, 29, e36-e37.	0.6	1
59	Sensitivity and specificity of single and combined clouds analyses compared with quantitative motor unit potential analysis. <i>Muscle and Nerve</i> , 2021, 63, 225-230.	2.2	1
60	Association of Polymorphisms in Inflammatory Cytokines Encoding Genes With Anti-N-methyl-D-Aspartate Receptor Encephalitis in the Southern Han Chinese. <i>Frontiers in Neurology</i> , 2020, 11, 553355.	2.4	1
61	Outcome Prediction by 40-Hz Steady-State Response After Large Hemispheric Infarction. <i>Frontiers in Neurology</i> , 2018, 9, 1093.	2.4	0
62	Autoantibodies detection in anti-N-methyl-D-aspartate receptor encephalitis. <i>Annals of Translational Medicine</i> , 2021, .	1.7	0